Statistics Tutorial

Robert Les

ATLAS Open Data Tutorial, Nov 25 2025



Introduction



Statistics



In high-energy physics (and all of science) statistics are needed to address some fundamental questions of using data

- Experiment design:
  - What cuts are “best” to separate my signal and background?
  - Can I model the data? What parameters can I estimate?
  - How do sources of uncertainty effect my results?
- Interpretation:
  - Is my result “significant” or could this be by chance?
  - How uncertain am I on this?
  - Which model best describes the data?

Statistics is the field of interpreting data

Gambler's fallacy: If something happened less frequently, it must occur more often in the future

Persecutor's fallacy: Ignoring base rate when claiming something unlikely

Fallacy of division: Something that is true for a group, is not true for each individual

Correlation=Causation: Correlation could be accidental, or due to third true cause

Berkson's paradox: Biased selection of data can lead to correlations









Outline and Extra Resources

This material can be found in technical detail in:
Cowan - Statistical Data Analysis

For all levels recommend the channel:
StatQuest with Josh Starmer

A bit impossible to cover a whole field but will follow
- Fundamental Concepts
- Estimators and Likelihood Models
- Hypothesis Testing: ATLAS Cheat Sheet
- Hands-on tutorials with ATLAS open-data




MODIFIED BAYES' THEOREM:
$P(H|X) = P(H) \times \left(1 + P(C) \times \left(\frac{P(X|H)}{P(X)} - 1\right)\right)$
H: HYPOTHESIS
X: OBSERVATION
P(H): PRIOR PROBABILITY THAT H IS TRUE
P(X): PRIOR PROBABILITY OF OBSERVING X
P(C): PROBABILITY THAT YOU'RE USING BAYESIAN STATISTICS CORRECTLY

Probability and definitions

Probability P is a way of describing how likely some event X is to happen

- $P(X) = 0$: Never, $P(X) = 1$: Certainly, $0 < P(X) < 1$: Maybe
- Often convert $P(X) = 0.12$ to percentage 12%
- There may be many X possible but the sum of all probabilities $= 1$



Examples:

- A fair coin has 2 sides $\rightarrow$ 1 out of 2 outcomes $\rightarrow$ 50% probability of heads.
- Under similar weather conditions there is a 30% probability of rain
- There was a 5% chance this effect in our data could be cause by random chance

A probability distribution function (pdf) is a continuous version of this: any interval corresponds to a probability

The Gaussian distribution



The Gaussian/Normal distribution is the most recurring distribution in statistics

- Has two parameters:
  - the mean (location)$=\mu$
  - the variance (scale)$=\sigma$
- Many useful mathematical properties make it easy to work with
- Many other distributions become gaussian in certain regimes

The central limit theorem: Under most conditions the average/sum of many random variables becomes gaussian as the number increases

- We often measure averages value
- Uncertainties are often “sums” of smaller effects
- Can extrapolate from a sub-sample to the whole if theorem holds

Classic example: Sum of dice rolls



Original probability for a dice is not Gaussian, but many summed valued are

Would be true for weighted dice or any probability distribution

*Under mostly general conditions

The Poisson distribution





The Poison distribution is especially relevant for high-energy physics

- Has one parameters:
  - the expected value (location)$=\lambda$
- Always predict integer values: $1, 2, 3, \ldots$
- Not symmetric
- Around $\lambda \sim 8$ starts to look Gaussian

Expresses the probability of number of occurrences in an interval. Aka describes "counting experiments"

- How many emails I get per day
- How many rain-drops in an area
- How many events I see in my detector element

But what does a probability “mean”? More then just an abstract philosophical question

Most common definition is that of **frequentist statistics**

- The subsets X are called events
- The probability is the long-term relative frequency of the event

$$P(X) = \lim_{n \to \inf} \frac{\text{occurances of event } X}{n}$$



Interpretation of Probability: Bayesian

Another common approach is the **Bayesian statistics**

- The subsets refer to hypothesis/prepositions that are either true/false
- Then the probability is:

$$P(X) = \text{degree of belief proposition } X \text{ is true}$$

Bayes theorem is central as can say:

$$P(\text{theory}|\text{data}) \propto P(\text{data}|\text{theory}) \cdot P(\text{theory})$$



Interpretation of Probability: Comparison

Frequentist

- The theory parameter **is a constant of nature**
  - Say probability my interval covers the theory parameter
- Assumes you repeated an experiment infinite times
- **No prior assumption**
- Decision rules on the compatibility of an alt model

Bayesian

- The theory parameter **is a random variable**
  - Can say $p(\text{theory}|\text{data})$ in an interval
- Clear for a single experiment case
- **Need to make an assumption on theory prior** $p(\text{theory})$
- Decision rules on the comparison of the theory probabilities

One not "more correct" then the other, but affects methodology
ATLAS uses mainly frequentist statistics, so will phrase the rest of the slides around that


R²=0.06
REXTHOR, THE DOG-BEARER
I DON'T TRUST LINEAR REGRESSIONS WHEN IT'S HARDER TO GUESS THE DIRECTION OF THE CORRELATION FROM THE SCATTER PLOT THAN TO FIND NEW CONSTELLATIONS ON IT.

Estimators

But you don't seee $P(X)$ directly in experiments! Just some sample of data $X_1, X_2, ..., X_N$

An estimator $\hat{\theta}$ is a function of data used to approximate a model property θ

Example estimators for μ and σ of a Gaussian

- Sample mean $\hat{\mu} = \frac{\sum_i^N X_i}{N}$
- Sample variance $\hat{\sigma}^2 = \frac{\sum_i^N (X_i - \hat{\mu})^2}{N-1}$

A histogram of X divided by N is also an estimate of $P(X)$

Note that estimators are a function of data and so are random variables too

- They have some uncertainty on them too!



We call the process of getting estimators from data as "fitting" $\rightarrow$ In almost all cases it is done numerically

Functional form fitting

An estimator example: Estimating a function f which models your data

Have measured values y_i at point x_i

Assume your data follows:

$$f(x|\boldsymbol{\theta}) = \sum_j^N f_j(x)\theta_j$$
$$= \theta_0 + \theta_1 x + \theta_2 x^2 + \ldots$$



Define the best estimator $\hat{\theta}$ as the one which minimizes the "least squares":

$$r = \sum_i (y_i - f(x_i|\boldsymbol{\theta}))^2$$

Functional form fitting



The function $f(x|\theta)$ is an arbitrary choice.

Compromise on the number of paramaters

- Not enough $\rightarrow$ bad fit
- Too many $\rightarrow$ overfitting

"Goodness of fit": $\chi^2 = \sum_i \frac{|y_i - f(x_i|\boldsymbol{\theta})|^2}{\sigma_i^2} \sim dof$

Degree of freedom (dof) $=$ number of fit parameters - number of points

A likelihood encodes the compatibility of data with a model as the product of individual probabilities for x

$$L(\boldsymbol{\theta}|\boldsymbol{x}) = \prod_i P(x|\boldsymbol{\theta})$$



It's to be thought of as a function of parameters while P is a function of data

The method of maximum likelihood says the best estimated values $\hat{\boldsymbol{\theta}}$ from your data is the value which maximizes likelihood

The uncertainity on $\hat{\boldsymbol{\theta}}$ can be estimated from the curvature of $L(\boldsymbol{\theta}|\boldsymbol{x})$



Nuisance Parameters

Very often there are parameters in your model you care about:

- Parameters of Interest (POI): $\boldsymbol{\mu}$

And parameters you don't care about:

- Nuisance Parameters: $\boldsymbol{\theta}$

E.x. I may only care about the μ of a Gaussian and not it's σ (or vice-versa)

In frequenist statistics we will sometimes employ the technique of "profiling" to remove these degrees of freedom by replacing $p(x|\boldsymbol{\mu},\boldsymbol{\theta}) \to p(x|\boldsymbol{\mu},\hat{\boldsymbol{\theta}})$

Let's denote the the best fit value of $\boldsymbol{\theta}$ at a specific $\boldsymbol{\mu}$ assumptions as $\hat{\hat{\boldsymbol{\theta}}}$

ATLAS Likelihood Models

Very common in ATLAS analysis, the use binned likelihood of the form:

$$L(\mu, \boldsymbol{\theta}|\boldsymbol{x}) = \prod_i Pois(x_i|\mu \cdot s_i(\boldsymbol{\theta}) + b_i(\boldsymbol{\theta})) \prod_j C(\alpha_j|\theta_j)$$

Each bin of your histogram is a counting experiment, where we expect b_i background and s_i signal events

$\implies \; Pois(x_i|\mu \cdot s_i + b_i)$

Where μ is the signal strength

Systematics encompass some possible shift to the predictions

$\implies \; s_i \rightarrow s_i(\boldsymbol{\theta}) \qquad b_i \rightarrow b_i(\boldsymbol{\theta})$

$C(\alpha_j|\theta_j)$ represents an external constraint on your nuisance parameters



Each histogram bin you are looking at is an individual counting experiment where you see x events and expect $\mu s + b$ events

$$Pois(x|\lambda = \mu s + b) = \frac{(\mu s + b)^x e^{-(\mu s + b)}}{x!}$$

Notes:

- μ is a handle on our signal assumption
- x is an integer, while $\mu s + b$ can be a float
- There isn't knowledge of the bin edges
  - Each bin is like a separate region

$\Longrightarrow$ Product each bin terms in the likelihood





Building the ATLAS model: Constraint terms

For systematics usually know the "size" of these variations beforehand, and so we include that knowledge in the model itself

$\implies$ Add constraint term $C(\alpha_j|\theta_j)$

Example the jet-energy scale uncertainty:

- Measured in dedicated analysis
- But still allow our analysis to vary it (θ_j) if improves modelling
- The constraint term tells the fit that we already know the systematics is this size (or smaller)
- The α_j is known as the global/auxiliary measurement data



- Usually assume and scale things so $C(\alpha_j|\theta_j) = Gauss(0|\theta_i, 1)$
- c.f MC stat uncertainity which is Poisson

Binned likelihood:

$$L(\mu, \boldsymbol{\theta}|\boldsymbol{x}) = \prod_i Pois(x_i|\mu \cdot s_i(\boldsymbol{\theta}) + b_i(\boldsymbol{\theta})) \prod_j C(\alpha_j|\theta_j)$$

To actually make the model you provide:

1) The nominal expected signal and background histograms
   - Define s, b
2) Alternative histograms templates for each possible systematic source
   - Define θ dependence of s, b
   - Add in a constraint term if needed
3) The data histogram
   - Define x to evaluate on



Understanding pulls

When we "run a fit" we evaluate the point and interval estimate for all parameters

A "pull" for a systematic source $= \frac{\hat{\theta} - \theta_{prefit}}{\sigma_\theta}$

By construction a constrained nuisance parameter will should have pull of zero and have uncertainty ± 1

- Pulls/constraints can tell you real physics effects
  - A pull means the data prefers a different value then the nominal
  - A constraint say your analysis can constrain this systematics more then the CP measurement
  - Need to understand both of these effects
- If pulls/constraint everywhere $\rightarrow$ bad fit model
- If underconstrained $\rightarrow$ broken fit model

The correlations of parameters and shapes of ± 1 variations can give insight




STATS EXAM GRADE
100
90
80
70
STUDENT B
STUDENT C
STUDENT A
β=1.94
p=0.586
86 88 90 92 94
SCREAM LOUDNESS (DECIBELS)
DARN, NOT SIGNIFICANT.
WE NEED MORE DATA. HAVE THEM EACH TRY YELLING INTO THE MIC A FEW MORE TIMES.
STATS EXAM GRADE
100
90
80
70
STUDENT B
STUDENT C
STUDENT A
β=1.94
p=0.037*
*SIGNIFICANT!
86 88 90 92 94
SCREAM LOUDNESS (DECIBELS)
PERFECT!
ARE YOU SURE WE'RE DOING SLOPE HYPOTHESIS TESTING RIGHT?

What if I have multiple competing models $\boldsymbol{\theta_i}$ that I want to make a statement on?

Example: counting experiment

- $\boldsymbol{\theta_0}$ =Background-only model of expected 10 events
- $\boldsymbol{\theta_1}$ =Signal+background model of expected 15.5 events
- Observe 11 events



Can I say something about which model the data favors or excludes?

* c.f. In Bayesian setting can measure $p(\boldsymbol{\theta_i})$, in frequentist $\boldsymbol{\theta_i}$ either true/false objectively

Hypothesis Testing

In frequentist setting, make statements on a hypothesis by defining a quantity of the data "test-statistic" and a critical region

1) Define null/alt hypotheses $\boldsymbol{\theta_0}/\boldsymbol{\theta_1}$
   - For a SM measurement: null=background, alternative=signal+background

2) Define a test-statistic t
   - E.x. the number of observed events

3) Derive the distribution of the test-static under the hypotheses $p(t|\boldsymbol{\theta_i})$

4) Define a threshold t_{thres} for a specific false negative rate α
   - Often $\alpha = 0.05$ in science



5) Only now look at data t_{obs} and compare to t_{cut}
   - Often quote the p-value: the probability **to observe** t_{obs} **or more extreme** under the null hypothesis

Hypothesis testing "colloquially"

The core point is condensing all of your data into one number: the test statistic t

- If know how it's distributed under you assumption, can evaluate how likely the observed value was to happen
  - Why you see p-values everywhere in science
- Define a region before looking at data:
  - If in this region, assumption is unlikely/rejected



Alternative hypothesis comes into play for further understanding

- What choice of t is the most powerful?
- How much data I need?

IMPORTANT! We only **reject the null** when in critical region

No action otherwise!

- **Never accept the alt!** Could be other unconsidered alt hypothesis
- If not in critical region **null is acceptable**
- Rejecting also doesn't 100% guarantee the null is false, just was unlikely
- These are frequentist statements based on long-term repeated experiment probability
  - $\alpha = 0.05$ means 1/20 experiments will make a false rejection



σ	Frequency	False positives if 300 papers/year
1σ	16%	~weekly
2σ	2.2%	~bi-monthly
3σ	0.13%	~yearly
4σ	0.0032%	~10 years
5σ	0.00003%	~100 years

In HEP we often convert the p-value to a "significance" which is the x-value for the equivalent p-value of the Gaussian

Neyman-Pearson Lemma tells us that for simple $\boldsymbol{\theta_0}$ vs $\boldsymbol{\theta_1}$, the test-statistic with the lowest β at chosen α is from the likelihood ratio

$$t = \frac{L(\boldsymbol{\theta_0}|\boldsymbol{x})}{L(\boldsymbol{\theta_1}|\boldsymbol{x})}$$

But not true for our likelihoods $L(\mu, \boldsymbol{\theta}|x) \rightarrow$ Approximate by profiling

$$t_\mu = -2\ln\left(\frac{L(\mu, \hat{\hat{\boldsymbol{\theta}}}|\boldsymbol{x})}{L(\hat{\mu}, \hat{\boldsymbol{\theta}}|\boldsymbol{x})}\right)$$



"Easy" interpretation: higher t_μ means more incompatible with μ

In ATLAS we use slightly adjusted test-statistics based on this

Before we were asking: "What is the chance my signal is real for a specific model parameter?"

If don't see signal want to also ask: "What model parameters can I exclude?"



Calculate the "limits":

- What signal-strength/cross-section can I **not** exclude at $\alpha = 0.05$
- Here your SM+BSM model is your null-hypothesis
- You scan over different μ assumptions
  - Calculate the p-value
  - The excluded range is when $p < 0.05$

For searches often consider multiple hypothesis (e.x. masses)

- Re-run the limit procedure above for each mass/parameter then plot
- Each $\alpha = 0.05$ horizontal-slice (top) is one x-axis point's vertical-slice (bottom)



Example: Typical SM measurement



Hypothesis testing



- Design analysis
- Hypothesis testing: background (null) vs signal+background (alt)
  - Possibly reject background-only model
  - Background only p-value and significance
- Point and interval estimate of signal strength

Example: Typical SM measurement



Hypothesis testing



- Design analysis
- Hypothesis testing: SM (null) vs SM+BSM (alt)
  - Possibly reject SM hypothesis
  - SM p-value and significance
- Hypothesis testing: SM+BSM (null) vs BSM (alt)
  - Invert test to derive upper limit on BSM

Summary

Crash-course in basic statistics related to ATLAS

- Estimators, Hypothesis Testing, ATLAS specific topics
- Try to clear-up frequent misunderstandings of p-values, hypothesis testing, etc
- May not digest it all the first time through
  - Exposure is the best way to eventually understand remember!
  - Get involved in the statistical analysis of your analysis

Two notebooks to work through

- Background fitting, model-building, and hypothesis testing with $H \to \gamma\gamma$ open-data
- Upload a public likelihood model and run limit calculations

Separate github repository currently

- Use this docker link

BACKUP

Probability and definitions

Kolmogorov axioms define probably as a set X and a function P

1) $P(X) = 1$
2) For every subset S in X, $P(X) > 0$
3) For every countable disjoint subset $S_1, S_2, \ldots$, $P(\bigcup S_i) = P(\sum S_i)$

* Note X can be multi-dimensional (x, y) etc



The above is clear for discrete variables, but can extend to continuous probability distribution function (pdf) $p(x)$

- Only a definite integral of a pdf is a probability: $P(x_1 < x < x_2) = \int_{x_1}^{x_2} p(x)dx$
- Can usually change $\sum \rightarrow \int$ and $P(X) \rightarrow p(x)$ safely

Definitions continued

The conditional probability of X given Y is defined as:

$$P(X|Y) = \frac{P(X \cap Y)}{P(Y)}$$

A little bit of math gives you the famous Bayes Theorem:

$$P(Y|X) = \frac{P(X|Y)P(Y)}{P(X)}$$

So what? $\implies$ Allows you to relate the probability of a cause given it's effect!

Let's also define events as independent if $P(X \cap Y) = P(X)P(Y)$



Image from wikipedia

Expectation values

Any element of X or a function of it is called a random variable
For a random variable a and pdf p define the expectation value:

$$E[a] = \int_{-\infty}^{+\infty} a\, p(x) dx$$

The famous expectation values are:

- mean $E[x] = \mu_x$
- variance $E[(x-\mu)^2] = \sigma_x^2$

σ_x is usual measure of spread in a parameter $\rightarrow$ if Gaussian, my parameter should be within $[\mu - \sigma, \mu + \sigma]$ 68% of the time



The covariance between two random values $E[(a-\mu_a)(b-\mu_b)] = \sigma_{ab}$ is also an important measure of linear correlations

But you don't deal with $p(x)$ directly in experiments but some sample of data $x_1, x_2, ..., x_N$

An estimator $\hat{\theta}$ is a function of data used to approximate a model property θ

Example estimators for μ and σ

- Sample mean $\hat{\mu} = \frac{\sum_i^N x_i}{N}$
- Sample variance $\hat{\sigma}^2 = \frac{\sum_i^N (x_i - \hat{\mu})^2}{N-1}$

Some good properties for an estimator

- Consistent: $\lim_{N \to \infty} \hat{\theta} \to \theta$
- Unbiased: $b = E[\hat{\theta}] - \theta = 0$
- Efficient: lowest $MSE = \sigma^2_{\hat{\theta}} + b^2$
- Robust: single x_i can not make $\hat{\mu}$ an arbitrary value



* Note that estimators are a function of data and so are random variables too

We call the process of getting estimators from data as "fitting" $\to$ In almost all cases it is done numerically

An example of an estimator: Functional form χ^2 fitting
If you have some measured values y_i at point x_i with some gaussian uncertainty σ_i

Assume your data follows a function of parameters:

$$\lambda(x, \boldsymbol{\theta}) = \sum_j^N f_j(x)\theta_j$$

Define the best estimator $\hat{\theta}$ as the one which minimizes:

$$\chi^2 = \sum_i \frac{y_i - \lambda(x_i, \boldsymbol{\theta})}{\sigma_i}$$



If your λ is linear in θ_j can solve analytically, if non-linear can use software packages to numerically calculate the values

Likelihood Fits: Point estimates

Since we only care about maxima, we will often instead minimize the the negative log-likelihood for simplicity

Assume we have one sample and a likelihood model:

$$L(\mu|x_0) = N \exp\left(\frac{(x_0-\mu)^2}{2\sigma^2}\right)$$

$$-\ln L(\mu|x_0) = -\frac{(x_0-\mu)^2}{2\sigma^2} - \ln(N)$$

and the estimator $\hat{\mu}$

$$\frac{\partial(-\ln L)}{\partial\mu} = -2\frac{(x_0-\mu)}{2\sigma^2} = 0$$

$$\implies \hat{\mu} = x_0$$



What about the uncertainty on our estimator?

The Crámer-Rao bound gives:

$$\sigma_{\hat{\theta}} \geq \left(1 + \frac{\partial b}{\partial \theta}\right)^2 / E\left[\frac{\partial^2 \ln L}{\partial \theta^2}\right]$$



For efficient operators (like MLE) the equality holds, and often look at unbiased estimators

$\rightarrow$ The uncertainty on the estimator can be estimated from the curvature of the likelihood at the minimum

The term $I_{ij} = E\left[\frac{\partial^2 \ln L}{\partial \theta_i \theta_j}\right]_{\theta=\hat{\theta}}$ is called the Fisher information matrix

*This method is also known as HESS in HEP jargon

Another method can be derived for interval estimate

Expanding the likelihood around the extrema at leading order and using the CRB:

$$\ln L(\theta) \approx \ln L(\hat{\theta}) + \frac{1}{2!}\frac{\partial^2 \ln L}{\partial \theta^2}\bigg|_{\theta=\hat{\theta}} (\theta - \hat{\theta})^2$$
$$= \ln L_{max} + \frac{(\theta - \hat{\theta})^2}{2\sigma^2_{\hat{\theta}}}$$



which can be rephrased as

$$-\ln L(\hat{\theta} \pm N\sigma) = \ln L_{min} + \frac{N^2}{2}$$

$\rightarrow$ So can also estimate the intervals graphically with y-axis crossing points

*This method is also known as MINOS in HEP jargon

In frequentist setting, make statements on a hypothesis by defining a quantity of the data "test-statistic" and a critical region

1) Define null/alt hypotheses $\boldsymbol{\theta_0}/\boldsymbol{\theta_1}$
   - For a BSM search: null=SM, alternative=SM+BSM
2) Define a test-statistic t
   - E.x. the number of observed events
3) Derive the distribution of the test-static under the hypothesis $p(t|\boldsymbol{\theta_i})$
4) Defines a false negative/postive rate $\alpha = \int_t^\infty p(t|\boldsymbol{\theta_0})dt$, $\beta = \int_{-\infty}^t p(t|\boldsymbol{\theta_1})dt$
5) Define a threshold t_{thres} for a specific α/β threshold
   - Often $\alpha = 0.05$
6) Only now look at data t_{obs} and compare to t_{cut}
   - Or equivalently the p-value$= \int_{t_{obs}}^\infty p(t|\boldsymbol{\theta_0})dt$ to α



Hypothesis Testing

In frequentist setting, make statements on a hypothesis by defining a quantity of the data "test-statistic" and a critical region

To interpret we only **reject the null** when in critical region

- **Never accept the alt!** Could be other reasonable alts
- If not in critical region **null is acceptable**
- Rejecting null does not mean $p(\boldsymbol{\theta_0}) = 0$
- These are frequentist statements based on long-term repeated experiment probability
  - $\alpha = 0.05$ means 1/20 false null-rejections



Quantify null agreement with p-value$= \int_{t_{obs}}^{\infty} p(t|\boldsymbol{\theta_0})dt$

- How often you would observe **this value or more extreme** if the **null hypothesis was true**

In HEP we often convert the p-value to a "significance" which is the x-value for the equivalent p-value if the $p(t|\boldsymbol{\theta_0}) = Gauss(t|\mu = 0, \sigma = 1)$

- $3\,(5)\sigma$ is approximately $\alpha = 0.003\,(5 \times 10^{-7})$

Neyman Pearson Lemma

Neyman-Pearson Lemma tells us that for simple $\boldsymbol{\theta_0}$ vs $\boldsymbol{\theta_1}$, the test-statistic with the lowest β at chosen α is from the likelihood ratio

$$t = \frac{L(\boldsymbol{\theta_0}|\boldsymbol{x})}{L(\boldsymbol{\theta_1}|\boldsymbol{x})}$$

But not true for our likelihoods $L(\mu, \boldsymbol{\theta}|x) \rightarrow$ Approximate by profiling

$$t_\mu = -2\ln\left(\frac{L(\mu, \hat{\hat{\boldsymbol{\theta}}}|\boldsymbol{x})}{L(\hat{\mu}, \hat{\boldsymbol{\theta}}|\boldsymbol{x})}\right)$$



"Easy" interpretation: higher t_μ means more incompatible with μ

In ATLAS we use slightly adjusted test-statistics based on this

For searches often care about providing "limits":

- What signal-strength/cross-section can I **not** exclude at $\alpha = 0.05$
- Here your SM+BSM model is your null-hypothesis
  - Not the same as background-only p-value!
- "Invert" the hypothesis test:
  - For multiple μ assumptions calculate the p-value
  - Exclude values with $p < 0.05$

For searches often consider multiple hypothesis (e.x. masses)

- Separate model, test-statistic, and inversion for each
- In limit plot, each $\alpha = 0.05$ slice is one y-axis point





Asymptotic approximations

If assume θ_{true} true, Wald approximation says that log-likelihood test-statistic as $n \to \infty$ will follow a non-central χ^2 distribution with

- degrees of freedom=dim($\boldsymbol{\theta}$)   centrality=$\frac{\mu - \mu_{true}^2}{\sigma}$

This asymptotic approximation makes it very computationally easy to derive $p(t_\mu|\mu') \to$ quick analytic expression!

If not valid ($n \sim O(1)$) then need to do full frequentist computation using pseudo-experiments (aka "toys")

- Sample the likelihood $x' \sim L(\mu'|\hat{\hat{\boldsymbol{\theta}}})$ multiple times
- Calculate t_μ for each pseudo-experiment
- Histogram to approximate $p(t_\mu|\mu')$



ATLAS Test Statistic

In ATLAS we use NP-inspired test-statistics profiling over nuisance parameters

For evaluating signal+background model agreement (limit setting) use

$$\tilde{q}_\mu = \begin{cases} -2\ln\frac{L(\mu,\hat{\hat{\theta}}(\mu))}{L(0,\hat{\hat{\theta}}(0))} & \hat{\mu} < 0 \\ -2\ln\frac{L(\mu,\hat{\hat{\theta}}(\mu))}{L(\hat{\mu},\hat{\theta})} & 0 < \hat{\mu} < \mu \\ 0 & \mu < \hat{\mu} \end{cases}$$

For background-only model (discovery) agreement

$$q_0 = \begin{cases} -2\ln\frac{L(\mu,\hat{\hat{\theta}}(\mu))}{L(0,\hat{\hat{\theta}}(0))} & \hat{\mu} > 0 \\ 0 & \hat{\mu} < 0 \end{cases}$$

The ranges are defined so that:

- Only look for upward fluctuations
- For searches, fluctuations larger then a specific signal strength are also consistent with it

Confidence intervals and Neyman Construction

Let's be more precise when we say we measure/exclude something

An X% **confidence interval** on a theory parameter θ means if we repeat the experiment our **interval** would cover the true parameter X% of the time

- Theory parameter is fixed constant of nature
- The bound edges change across repeated experiments
- **Does not say** that X% prob your derived interval covers true value

Can define the procedure from assumed pdf beforehand:

- Once observe x_{obs} can derive interval $[\theta_-, \theta_+]$
- Interval can be also one-sided e.x. $[-\infty, \theta_+]$
- For Gaussian pdf 68% CI same as $[\mu - \sigma, \mu + \sigma]$



From Cranmer



More likelihood models than binned-Poisson or Gaussain can be used:

$$L(N_S, N_B, \boldsymbol{\theta}|\boldsymbol{x}) = \frac{(N_S+N_B)^N e^{-N_S+N_B}}{N!} \prod_i^N \frac{N_S}{N_S+N_B} p_S(x_i|\boldsymbol{\theta}) + \frac{N_B}{N_S+N_B} p_B(x_i|\boldsymbol{\theta})$$

- $p_{S/B}(x_i|\boldsymbol{\theta})$ are pdf choices for events from the processes
  - E.x. $p_S(x_i|\boldsymbol{\theta}) = Gauss(x_i)$
    $p_B(x_i|\boldsymbol{\theta})$ exponential
- "Extended" term at front since event count N a Poisson variable
  - $N_{S/B}$ are the expected yields



Exploits all structure within your data!
- Better variances!
- Minimization difficult as N large
- Need good assumptions on pdf

Multiple dimensions

Mainly discusses situation where one theory parameter, but most results scale for case with several "parameters-of-interest" $\boldsymbol{\theta}$

Main thing to remember is that asymptotically follow a χ^2 with higher d.o.f. parameter $k > 1$

Find intervals by drawing point where $\log L(\boldsymbol{\theta}) = \log L_{max} + \frac{Q_k}{2}$

- Look-up tables for Q_k threshold at chosen CI



Error propagation

First useful example of statistics is how to propagate uncertainty

If I measure some observables $\boldsymbol{x}$ (say object 4-vectors) and it's related to another quantify $y(\boldsymbol{x})$ (say the invariant mass), at first order

$$y(\boldsymbol{x}) = y(\boldsymbol{\mu}) + \sum_i \frac{\partial y}{\partial x_i}\Big|_{\boldsymbol{x}=\boldsymbol{\mu}} + \ldots$$

The mean and variation of the new quantity is:

$$\mu_y = y(\boldsymbol{\mu})$$

$$\sigma_y^2 = \sum_i \sum_j \frac{\partial y}{\partial x_i}\frac{\partial y}{\partial x_j}\Big|_{\boldsymbol{x}=\boldsymbol{\mu}} \sigma_{x_i x_j}$$

You will often see we transform to new basis so that the covariances $\sigma_{y_i y_j}$ are zero to simply calculations
$\rightarrow$ eiegenvector decomposition of jet and b-jet experimental systematics

Toy Results

Proper limit calculations are done by toy calculations (very CPU expensive!)



Don't assume toy calculations perfect

- Make sure distributions are smooth with enough statistics
- Check search range is reasonable
  - 0.05 crossing visible even for $\pm 2\sigma$ band
- Make sure no odd features
  - Toy fits can fail
  - Nuisance parameters can act odd: MC stats, spurious signal

CLs

For limits as mentioned calculate signal+background p-values

But remember by construction will exclude 5% of the time

- Even for cases where we have no signal sensitivity!



Combat this with CLs approach, replace normal p-value with $CLs = \frac{p_{s+b}}{p_b} = \frac{\int_{t_{obs}}^{\infty} p(t|s+b)dt}{\int_{t_{obs}}^{\infty} p(t|b)dt}$

- Penalty if s+b not very separable from b
- Conservative: Doesn't match 95% CI anymore



But what about "expected results" $\rightarrow$ define as the median result given a (background) hypothesis

In the asympotic approximation can use the Asimov dataset

- Artificial dataset where all all estimators are the true values
- Easy estimators and also their uncertainty
- Logic: median(toy result)= result(median dataset)

Can be used as a shortcut to derive the expected result and it's uncertainty





Analyzers may know full toy calculations is a very CPU/time intensive task
$\rightarrow$ but **when asymptotics breakdown they are overly aggressive!**

Better-then-zero effect : The 95% upper-limit for a Poisson experiment with $n_{obs} = 0$ is 3

- Not true for asymptotics with systematics!

Quick rule-of-thumb **asymptotic don't apply if you see**:
$N_{excl} = \sigma_{excl} \times L \times \mathrm{acc} \lesssim 5$

Where do I need to run toys

- When have sensitive bins near this threshold
- Typically impact limits ~20-30%

Look elsewhere effect

Our hypothesis testing definitions was framed interpretating data **only between two hypothesis**

- But we are often doing many hypothesis tests at once!
- Each "local" hypothesis p-value misleading as not independent!
- Need to define a "global" p-value to see deviations anywhere in data

a.k.a. Look-elsewhere-effect (a.k.a. multiple comparisons problem)

Gross-Vitells estimation formula on the number of up-crossings N_{up} at arbitrary threshold Z_{ref}:

$$p_{global} = p_{local} + N_{up} e^{\frac{1}{2}(Z_{local}^2 - Z_{ref}^2)}$$

Can also measure with fully pseudo-experiment procedure

Functional Form Fits

Sometimes can't rely on simulations for deriving a template and need to fit some functional form to the data

Use simplest functional form allowed by data → provided by F-test

Generally three potential kinds of uncertainty on a background fit:

- Fit quality uncertainty:
  - Evaluate from hessian of fit or from toys
- Function-choice uncertainty (alternate pdf)
- "Spurious Signal" uncertainty
  - Bias of fitting signal+background model on background-only data





Spurious Signal uncertainty

- Bias of fitting signal+background model on background-only template
- $S_{spur} = N_{fit} - N_{template}$
- Add as additive nuisance parameter to model $Pois(x|\mu s + S_{spur}\theta_{SS} + b)$
- When possible evaluate fit from high-MC template or high-stats region

To keep impact of this bias on sensitivity $< 5\,(12)\%$ need $S_{spur}/\sigma_{S_{spur}} < 0.3\,(0.5)$

- Background model may be too flexible and can pick-up fake signal
  - Try less flexible fit model
- Understand if signal model too wide or background like





Combinations

What if we want to combine several measurements for optimal sensitivity

Often assume a multi-variate Gaussian likelihood (aka BLUE method):

$$\ln L \propto (\boldsymbol{x} - \mathbf{1})^{\intercal} \boldsymbol{V}^{-\mathbf{1}} (\boldsymbol{x} - \mathbf{1})$$

- $\boldsymbol{x}$ is the vector of all measurments
- $\boldsymbol{V}$ is their correlation matrix
  - Often will decompose into $\boldsymbol{V} = \boldsymbol{V}_{stat} + \boldsymbol{V}_{syst} + \ldots$

Assumptions:

- Data x is statistically independent, otherwise need to evaluate with bootstrapping
- Gaussian likelihood valid, otherwise need to do full likelihood level combination
  - Often the case

Reminder of our goal

Typical Likelihood looks like:

$$L(\mu, \boldsymbol{\theta}) = \prod_r \prod_i \mathrm{Pois}\Big(n_{i_r}^{obs} | n_{i_r}^{sig}(\mu, \theta_k) + n_{i_r}^{bkg}(\theta_k)\Big) \prod_k f_k(\theta_k)$$

i=bin, r=region, k=systematic

To construct our likelihood need to specify few things:

- The r's and i's aka the regions and the bins
- The θ_k aka the systematic (as well as constraint $f_k(\theta_k)$)
- The actual parameter we care about μ

Then we connect these to the actual data we have

- Specify the n's aka the bin contents from our histograms
- Specify how the systematics effect these bins $n(\theta_k)$, aka variation histograms

The output ROOT/pyhf "workspace" holds the the likelihood in a common format and also the n's

- You can do all the "interpretation" from this common format
  - Limits, plots, etc

Code workflow



The by-far most common mistake I see is analyzers assuming workspace built correctly and jumping right to limits

- p-values, limits, etc tell very summarized info so almost impossible to debug from there
- Many small simple things can go wrong: Forget samples/systematics
- Many complex things can go wrong: Systematics have weird shapes, large pulls
  - Some of these can be "physics" and not bugs

Statistics in 1 minute

For profile-likelihood fits everything we do is related to the likelihood $L(\mu, \boldsymbol{\theta})$

- A fit finds the minimum of $L(\mu, \theta)$, labeled by $\hat{\mu}, \hat{\boldsymbol{\theta}}$
- You can estimate the uncertainties on the parameters by the curvature of $L(\mu, \theta)$ near the minimum
- Often care about fits at specific hypothesis; e.g $\mu = 0$ background-only
- Asimov data$=$ artificial data where $\hat{\mu}, \hat{\boldsymbol{\theta}}$ are the “true” generated values

Limits are related to p-values

- Probability that data is consistent with some (alternate) hypothesis
- Background-only p_0: Consistency of background-only with data
- Limit: Value of μ (aka cross-section) for which p-value is 0.05

Difficulties with profile-likelihood usually related to understanding your systematic $\boldsymbol{\theta}$

- Always understand your likelihood before running limits

Typical nomenclature/jargon

- Observable: The variable you are fitting (aka the x_i)
- Sample: A single contribution to your total s_i/b_i, which is usually divided into different physics processes: aka V+jets, $t\bar{t}$,...
- A category/region: A signal/control/validation region of your analysis. Extends the i in your $\prod_i$
- Template: the histogram of your nominal s_i/b_i, or the histogram after a $\pm 1\sigma$ uncertainty shift
- Normalization uncertainties: Systematic which uniformly shifts a sample rate up/down in a specific region
- Shape uncertainties: Systematic which isn't uniform across the sample+region
- Correlated systematic: The $\pm 1\sigma$ shift of the systematic is simultaneous across the region/sample, aka one parameter encompasses effect across samples/regions