

Particle Detectors 2/2

Werner Riegler, CERN, werner.riegler@cern.ch

The ‘Standard Model’


matter particles

	1st gen.	2nd gen.	3rd gen.
QUARK	 up	 charm	 top
LEPTON	 down	 strange	 bottom

guage particles

Strong Force  Gluon
Electro-Magnetic Force  photon
Weak Force    W^+ W^- Z W bosons Z boson

scalar particle(s)



The ‘Standard Model’

$$\begin{aligned} L_{GSW} &= L_0 + L_H + \sum_l \left\{ \frac{g}{2} \bar{L}_l \gamma_\mu \vec{\tau} L_l \vec{A}^\mu + g' \left[\bar{R}_l \gamma_\mu R_l + \frac{1}{2} \bar{L}_l \gamma_\mu L_l \right] B^\mu \right\} + \\ &+ \frac{g}{2} \sum_q \bar{L}_q \gamma_\mu \vec{\tau} L_q \vec{A}^\mu + \\ &+ g' \left\{ \frac{1}{6} \sum_q [\bar{L}_q \gamma_\mu L_q + 4 \bar{R}_q \gamma_\mu R_q] + \frac{1}{3} \sum_{q'} \bar{R}_{q'} \gamma_\mu R_{q'} \right\} B^\mu \end{aligned}$$

$$\begin{aligned} L_H &= \frac{1}{2} (\partial_\mu H)^2 - m_H^2 H^2 - h \lambda H^3 - \frac{h}{4} H^4 + \\ &+ \frac{g^2}{4} (W_\mu^+ W^\mu + \frac{1}{2 \cos^2 \theta_W} Z_\mu Z^\mu) (\lambda^2 + 2 \lambda H + H^2) + \\ &+ \sum_{l,q,q'} (\frac{m_l}{\lambda} \bar{l} l + \frac{m_q}{\lambda} \bar{q} q + \frac{m_{q'}}{\lambda} \bar{q}' q') H \end{aligned}$$

Over the last century
this 'Standard Model' of
Fundamental Physics was discovered
by studying

Radioactivity

Cosmic Rays

Particle Collisions (Accelerators)

A large variety of Detectors and
experimental techniques have been
developed during this time.

Scales

$$E = m a^2$$

$$E = m b^2$$

$$E = m c^2 \quad \leftarrow \text{Energy} \cong \text{Mass}$$

⋮


$$m(\text{electron}) = 9.1 \cdot 10^{-31} \text{ kg}$$

$$m_e c^2 = 8.19 \cdot 10^{-14} \text{ J}$$

$$= 510\,999 \text{ Electron Volt (eV)}$$


$$= 0.511 \text{ MeV}$$

$$1 \text{ Electron Volt} = e_0 \cdot 1V = 1.603 \cdot 10^{-19} \text{ J}$$



1 Electron Volt = Energy an Electron gains as
it traverses a Potential Difference of 1V

Build your own Accelerator



$$E_{\text{kin}} = 1.5 \text{ eV} =$$

2 615 596 km/h

Scales

Visible Light: $\lambda = 500 \text{ nm}$, $h\nu \approx 2.5 \text{ eV}$

Excited States in Atoms: 1-100 keV "X-Rays"

Nuclear Physics: 1-50 MeV

Particle Physics: 1-1000 GeV (LHC 14 TeV)

Higher Measured Energy: 10^{20} eV (Cosmic Rays)

Basics

Lorentz Boost:

$$\mu^- \rightarrow e^- + \bar{\nu}_e + \gamma_\mu \quad \gamma = 2.2 \cdot 10^{-6} s$$

E.g. Produced by Cosmic Rays ($p, He, Li \dots$)
 colliding with air in the upper atmosphere $\sim 10 \text{ km}$

$$s = v \cdot \gamma \sim c \cdot \gamma = 660 \text{ m}$$

But we see Muons here on Earth

$$E_\mu \sim 2 \text{ GeV}, m_\mu c^2 = 105 \text{ MeV} \rightarrow \gamma \sim 19$$

$$\text{Relativity: } \bar{\gamma} = \gamma \cdot \gamma$$

$$s = c \cdot \bar{\gamma} = 12.5 \text{ km} \rightarrow \text{Earth}$$

$$\text{Pions: } \pi^+, \pi^- \quad \gamma \sim 2.6 \cdot 10^{-8} s, m_\pi c^2 = 135 \text{ MeV}$$

$$2 \text{ GeV} \rightarrow s = 115 \text{ m}$$

Pions were discovered in Emulsions exposed
 to Cosmic Rays on high Mountains.

<http://pdg.lbl.gov>

~ 180 Selected Particles

$\pi, W^\pm, Z^0, g, e, \mu, \tau, \nu_e, \nu_\mu, \nu_\tau, \pi^\pm, \pi^0, \eta, f_0(600), g(870),$
 $w(782), \eta'(858), f_0(980), a_0(980), \phi(1020), h_1(1170), b_1(1235),$
 $a_1(1260), f_2(1270), f_1(1285), \eta(1395), \pi(1300), a_2(1320),$
 $f_0(1370), f_1(1420), w(1420), \eta(1440), a_0(1450), g(1450),$
 $f_0(1500), f_2'(1525), w(1650), w_3(1670), \pi_2(1670), \phi(1680),$
 $g_3(1690), g(1700), f_0(1710), \pi(1800), \phi_3(1850), f_2(2010),$
 $a_4(2040), f_4(2050), f_2(2300), f_2(2340), K^\pm, K^0, K_s^0, K_L^0, K^*(892),$
 $K_1(1270), K_1(1400), K^*(1410), K_2^*(1430), K_2^*(1430), K^*(1680),$
 $K_2(1770), K_3^*(1780), K_2(1820), K_4^*(2045), D^\pm, D^0, D^*(2007)^0,$
 $D^*(2010)^\pm, D_1(2420)^0, D_2^*(2460)^0, D_2^*(2460)^\pm, D_s^\pm, D_s^{*\pm},$
 $D_{s1}(2536)^\pm, D_{s1}(2573)^\pm, B^\pm, B^0, B_s^0, B_c^\pm, \eta_c(1S), J/\psi(1S),$
 $\chi_{c0}(1P), \chi_{c1}(1P), \chi_{c2}(1P), \psi(2S), \psi(3770), \psi(4040), \psi(4160),$
 $\psi(4415), \tau(1S), \chi_{b0}(1P), \chi_{b1}(1P), \chi_{b2}(1P), \tau(2S), \chi_{b3}(2P),$
 $\chi_{b2}(2P), \tau(3S), \tau(4S), \tau(10860), \tau(11020), p, n, N(1440),$
 $N(1520), N(1535), N(1650), N(1675), N(1680), N(1700), N(1710),$
 $N(1720), N(2190), N(2220), N(2250), N(2600), \Delta(1232), \Delta(1600),$
 $\Delta(1620), \Delta(1700), \Delta(1905), \Delta(1910), \Delta(1920), \Delta(1930), \Delta(1950),$
 $\Delta(2420), \Lambda, \Lambda(1405), \Lambda(1520), \Lambda(1600), \Lambda(1670), \Lambda(1690),$
 $\Lambda(1800), \Lambda(1810), \Lambda(1820), \Lambda(1830), \Lambda(1890), \Lambda(2100),$
 $\Lambda(2110), \Lambda(2350), \Sigma^+, \Sigma^0, \Sigma^-, \Sigma(1385), \Sigma(1660), \Sigma(1670),$
 $\Sigma(1750), \Sigma(1775), \Sigma(1915), \Sigma(1940), \Sigma(2030), \Sigma(2250), \Xi^0, \Xi^-,$
 $\Xi(1530), \Xi(1690), \Xi(1820), \Xi(1950), \Xi(2030), \Omega^-, \Omega(2250)^-,$
 $\Lambda_c^+, \Lambda_c^+, \Sigma_c(2455), \Sigma_c(2520), \Xi_c^+, \Xi_c^0, \Xi_c^+, \Xi_c^0, \Xi(2645)$
 $\Xi_c(2780), \Xi_c(2815), \Omega_c^0, \Lambda_b^0, \Xi_b^0, \Xi_b^-, t\bar{t}$

There are Many more

Particle	Mass (MeV)	Lifetime τ (s)	$c\tau$
γ	0	∞	\sim
$\pi^\pm (\bar{u}\bar{d}, \bar{d}\bar{u})$	140	$2.6 \cdot 10^{-8}$	7.8 m
$K^\pm (\bar{u}\bar{s}, \bar{s}\bar{u})$	494	$1.2 \cdot 10^{-8}$	3.7 m
$K^0 (\bar{d}\bar{s}, \bar{s}\bar{d})$	497	$5.1 \cdot 10^{-8}$ $8.9 \cdot 10^{-11}$	15.5 m 2.7 cm
$D^\pm (\bar{c}\bar{d}, \bar{c}\bar{d})$	1869	$1.0 \cdot 10^{-12}$	375 μm
$D^0 (\bar{c}\bar{u}, \bar{u}\bar{c})$	1864	$4.1 \cdot 10^{-13}$	123 μm
$D_s^+ (\bar{c}\bar{s}, \bar{s}\bar{c})$	1969	$4.9 \cdot 10^{-13}$	147 μm
$B^\pm (\bar{u}\bar{s}, \bar{s}\bar{u})$	5279	$1.7 \cdot 10^{-12}$	502 μm
$B^0 (\bar{b}\bar{d}, \bar{d}\bar{b})$	5279	$1.5 \cdot 10^{-12}$	462 μm
$B_s^0 (\bar{s}\bar{b}, \bar{b}\bar{s})$	5370	$1.5 \cdot 10^{-12}$	438 μm
$B_c^+ (\bar{c}\bar{s}, \bar{s}\bar{c})$	~6400	$\sim 5 \cdot 10^{-13}$	150 μm
$\rho (\bar{u}u\bar{d})$	938.3	$> 10^{53} \text{ s}$	∞
$n (\bar{u}u\bar{d})$	939.6	885.7 s	$2.655 \cdot 10^8 \text{ km}$
$\Lambda^0 (\bar{u}\bar{d}s)$	1115.7	$2.6 \cdot 10^{-10}$	7.89 cm
$\Sigma^+ (\bar{u}\bar{u}s)$	1189.4	$8.0 \cdot 10^{-11}$	2.404 cm
$\Sigma^- (\bar{d}\bar{d}s)$	1197.4	$1.5 \cdot 10^{-10}$	4.434 cm
$\Xi^0 (\bar{u}\bar{s}s)$	1315	$2.9 \cdot 10^{-10}$	8.71 cm
$\Xi^- (\bar{d}\bar{s}s)$	1321	$1.6 \cdot 10^{-10}$	4.91 cm
$\Omega^- (\bar{s}\bar{s}s)$	1672	$8.2 \cdot 10^{-11}$	2.461 cm
$\Lambda_c^+ (\bar{u}\bar{d}c)$	2285	$\sim 2 \cdot 10^{-13}$	60 μm
$\Xi_c^+ (\bar{u}\bar{s}c)$	2466	$4.4 \cdot 10^{-13}$	132 μm
$\Xi_c^0 (\bar{d}\bar{s}c)$	2472	$\sim 1 \cdot 10^{-13}$	29 μm
$\Xi_c^0 (\bar{s}\bar{s}c)$	2698	$6.0 \cdot 10^{-14}$	19 μm
$\Lambda_b (\bar{u}\bar{d}b)$	5620	$1.2 \cdot 10^{-12}$	368 μm

"Secondary
Vertices"

From the 'hundreds' of Particles listed by the PDG there are only ~ 27 with a life time $c\tau > \sim 1\mu\text{m}$ i.e. they can be seen as 'tracks' in a Detector.

~ 13 of the 27 have $c\tau < 500\mu\text{m}$ i.e. only $\sim \text{mm}$ range at GeV Energies.
→ "short" tracks measured with Emulsions or Vertex Detectors.

From the ~ 14 remaining particles

$e^\pm, \mu^\pm, \gamma, \pi^\pm, K^\pm, K^0, p^\pm, n$


are by far the most frequent ones

A particle Detector must be able to identify and measure Energy and Momenta of these 8 particles.

Basics

Invariant Mass:

LAB:



$$\text{Relativity: } \tilde{a} = \begin{pmatrix} a_0 \\ \vec{a} \end{pmatrix}, \tilde{b} = \begin{pmatrix} b_0 \\ \vec{b} \end{pmatrix}, \tilde{a}\tilde{b} = a_0 b_0 - \vec{a}\cdot\vec{b}$$

$$E = mc^2\gamma, \vec{p} = m\vec{v}\gamma$$

$$\tilde{p} = \begin{pmatrix} E \\ \vec{p} \end{pmatrix}, \tilde{p}_1 = \begin{pmatrix} E_1 \\ \vec{p}_1 \end{pmatrix}, \tilde{p}_2 = \begin{pmatrix} E_2 \\ \vec{p}_2 \end{pmatrix}$$

$$\tilde{p} = \tilde{p}_1 + \tilde{p}_2 \quad \text{Energy + Momentum Conservation}$$

$$\tilde{p}^2 = (\tilde{p}_1 + \tilde{p}_2)^2 \rightarrow \tilde{p}\tilde{p} = \tilde{p}_1\tilde{p}_1 + \tilde{p}_2\tilde{p}_2 + 2\tilde{p}_1\tilde{p}_2$$


$$\underline{M^2c^2 = m_1^2c^2 + m_2^2c^2 + 2\left(\frac{E_1E_2}{c^2} - p_1p_2 \cos\theta\right)}$$

- Measuring Momenta and Energies OR
- Measuring Momenta and identifying Particles

gives the Mass of the original Particle

Momentum Measurement

Magnetic Spectrometer: A charged particle describes a circle in a magnetic field:



$$L = R \cdot \theta$$

$$S = R(1 - \cos \frac{\theta}{2}) \sim R \frac{\theta^2}{8} = \frac{L^2}{8R} \rightarrow R = \frac{L^2}{8S}$$

$$\Delta P = 0.3 B \Delta R = 0.3 B \frac{L^2}{8S^2} \Delta S$$

$$\Delta S = \frac{\sigma_x}{\sqrt{N}} \quad \sigma_x \dots \text{point resolution}, N \dots \text{Measurement Points}$$


$$\frac{\Delta P}{P} \cdot \frac{\Delta S}{S} = \frac{\sigma_x [\text{n}]}{\sqrt{N}} \cdot \frac{3.3 \cdot 8 P \left[\frac{\text{GeV}}{c} \right]}{B [\text{T}] \cdot L^2 [\text{m}^2]}$$

E.g.: $P = 10 \frac{\text{GeV}}{c}$, $B = 1 \text{T}$, $L = 1 \text{m}$, $\sigma_x = 200 \mu\text{m}$, $N = 25$

$$\frac{\Delta P}{P} = 0.01 \rightarrow 1\%$$

Limit → Multiple Scattering

Multiple Scattering




$$P \left[\frac{e_0 v}{c} \right] = 0.3 R[n] B[T]$$

$$\theta = \frac{L}{R} = \frac{L}{P} \cdot 0.3 B$$

$$\frac{\Delta P}{P} = \frac{\Delta \theta}{\theta} = \frac{\theta_0}{\theta} \sim \frac{0.05}{B B[T] L[m]} \cdot \sqrt{\frac{L}{x_0}}$$

→ Independent of P



$$\left| \frac{\Delta P}{P} \right|_{tot} = \sqrt{\left(\frac{\Delta \phi}{P} \right)_{Sog}^2 + \left(\frac{\Delta \phi}{P} \right)_{ns}^2}$$





Multiple Scattering

ATLAS Muon Spectrometer:
 $N=3$, $\sigma=50\text{um}$, $P=1\text{TeV}$,
 $L=5\text{m}$, $B=0.4\text{T}$

$\Delta p/p \sim 8\%$ for the most energetic muons at LHC





$$\left. \begin{array}{l} e^\pm \quad m_e = 0.511 \text{ MeV} \\ \mu^\pm \quad m_\mu = 105.7 \text{ MeV} \sim 200 m_e \\ \gamma \quad m_\gamma = 0, \quad Q = 0 \end{array} \right\} EM$$

$$\left. \begin{array}{l} \pi^\pm \quad m_\pi = 139.6 \text{ MeV} \sim 270 m_e \\ K^\pm \quad m_K = 493.7 \text{ MeV} \sim 1000 m_e \\ p^\pm \quad m_p = 938.3 \text{ MeV} \sim 2000 m_e \\ K^0 \quad m_{K^0} = 497.7 \text{ MeV} \quad Q=0 \\ n \quad m_n = 939.6 \text{ MeV} \quad Q=0 \end{array} \right\} \begin{array}{l} EM, Strong \\ \sim 3.5 m_\pi \\ Strong \end{array}$$

The Difference in
 Mass, Charge, Interaction
 is the key to the Identification

Tracking:


Momentum by bending in the B-field
Secondary vertices

Calorimeter:





Energy by absorption

Muons:

Only particles passing through calorimeters



- Electrons ionize and show Bremsstrahlung due to the small mass
- Photons don't ionize but show Pair Production in high τ Material. From here on equal to e^\pm
- Charged Hadrons ionize and show Hadron Shower in dense Material.
- Neutral Hadrons don't ionize and show Hadron Shower in dense Material
- Muons ionize and don't shower



Vertex Detector


Inner Tracking Chamber

Time Projection Chamber

Electromagnetic Calorimeter

Hadron Calorimeter

Muon Detectors







Fig. 1 - The ALEPH Detector


- Vertex Detector
- Inner Track Chamber
- Time Projection Chamber
- Electromagnetic Calorimeter
- Superconducting Magnet Coil
- Hadron Calorimeter
- Muon Detection Chambers
- Luminosity Monitors

$\gamma, e^\pm, \pi^\pm, k^\pm$

K^0, p, n, μ^\pm




CMS
A Compact Solenoidal Detector for LHC




$$Z \rightarrow e^+ e^-$$

Two high momentum charged particles depositing energy in the Electro Magnetic Calorimeter



$$Z \rightarrow \mu^+ \mu^-$$

Two high momentum charged particles traversing all calorimeters and leaving a signal in the muon chambers.



Interaction of Particles with Matter

Any device that is to detect a particle must interact with it in some way → almost ...

In many experiments neutrinos are measured by missing transverse momentum.

E.g. e^+e^- collider. $P_{\text{tot}}=0$,


If the $\sum p_i$ of all collision products is $\neq 0 \rightarrow$ neutrino escaped.



“Did you see it?”
“No nothing.”
“Then it was a neutrino!”


$$W^+W^- \rightarrow e^+ \square + \square_e + \square_{\mu}$$

Single electron, single Muon, Missing Momentum




$$Z \rightarrow q \bar{q}$$

Two jets of particles




$$Z \rightarrow q \bar{q} g$$

Three jets of particles




Two secondary vertices with characteristic decay particles giving invariant masses of known particles.

Bubble chamber like – a single event tells what is happening. Negligible background.





ALEPH Higgs Candidate





Undistinguishable background exists. Only statistical excess gives signature.



2010 ATLAS W, Z candidates !



2010 ATLAS W, Z candidates !



Higgs Boson at CMS



Particle seen as an excess of two photon events above the irreducible background.

Principles:

Only a few of the numerous known particles have lifetimes that are long enough to leave tracks in a detector.

Most of the particles are measured through the decay products and their kinematic relations (invariant mass). Most particles are only seen as an excess over an irreducible background.

Some short lived particles (b,c -particles) reach lifetimes in the laboratory system that are sufficient to leave short tracks before decaying → identification by measurement of short tracks.

In addition to this, detectors are built to measure the 8 particles

$$e^\pm, \mu^\pm, \gamma, \pi^\pm, K^\pm, K^0, p^\pm, n$$

Their difference in mass, charge and interaction is the key to their identification.


Detector Technologies

Solid state detectors close to the collision point for excellent position resolution to find primary and secondary vertices → **silicon pixel detectors**.

Solid state detectors (**silicon strip detectors**) or gas detectors at larger distances for tracking and momentum measurement.



Massive **calorimeters** with alternating layers of passive absorber material and active detector material for measurement of particle energies.

Silicon Pixel Detectors




ATLAS: 1.4×10^8 pixels

40 000 000 ‘images’ per second.




ATLAS Silicon Pixel Detector




Silicon Strip Detectors

Every electrode is connected to an amplifier →
Highly integrated readout electronics.


Two dimensional readout is possible.



CMS Outer Barrel Module



Silicon Strip Detectors



Time Projection Chamber (TPC):



Gas volume with parallel E and B Field.

B for momentum measurement. Positive effect:

Diffusion is strongly reduced by E//B (up to a factor 5).


Drift Fields 100-400V/cm. Drift times 10-100 μ s.

Distance up to 2.5m !



ALICE TPC: Construction Parameters

- Largest TPC:
 - Length 5m
 - Diameter 5m
 - Volume 88m³
 - Detector area 32m²
 - Channels ~570 000
- High Voltage:
 - Cathode -100kV
- Material X₀
 - Cylinder from composite materials from airplane industry ($X_0 = \sim 3\%$)



ALICE TPC: Pictures of the Construction

Precision in z: 250 μm



End plates 250 μm



Wire chamber: 40 μm




ALICE TPC Construction

My personal contribution:



A visit inside the TPC.



TPC installed in the ALICE Experiment




First 7 TeV p-p Collisions in the ALICE TPC in March 2010 !




11/16/2011

First Pb Pb Collisions in the ALICE TPC in Nov 2010 !




The Geiger Counter reloaded: Drift Tube

ATLAS MDT R(tube) =15mm



Calibrated Radius-Time correlation






Primary electrons are drifting to the wire.

Electron avalanche at the wire.

The measured drift time is converted to a radius by a (calibrated) radius-time correlation.

Many of these circles define the particle track.

ATLAS Muon Chambers



ATLAS MDTs, 80 μ m per tube

The Geiger counter reloaded: Drift Tube

Atlas Muon Spectrometer, 44m long, from r=5 to 11m.

1200 Chambers




6 layers of 3cm tubes per chamber.

Length of the chambers 1-6m !

Position resolution: $80\mu\text{m}/\text{tube}$, $<50\mu\text{m}/\text{chamber}$ (3 bar)

Maximum drift time $\approx 700\text{ns}$

Gas Ar/CO₂ 93/7




Detector Systems

CERN Neutrino Gran Sasso


(CNGS)

CNGS

If neutrinos have mass:



Muon neutrinos produced at CERN.
See if tau neutrinos arrive in Italy.



CNGS Project


CNGS (CERN Neutrino Gran Sasso)

- A long base-line neutrino beam facility (732km)
- send ν_μ beam produced at CERN
- detect ν_τ appearance in OPERA experiment at Gran Sasso




direct proof of ν_μ - ν_τ oscillation (appearance experiment)


CNGS




CNGS



CNGS



Radial Distribution of the ν_μ -Beam at GS



5 years CNGS operation, 1800 tons target:

- 30000 neutrino interactions
- ~150 ν_τ interactions
- ~15 ν_τ identified
- < 1 event of background

Neutrinos at CNGS: Some Numbers

For 1 year of CNGS operation, we expect:

protons on target

2×10^{19}

pions / kaons at entrance to decay tunnel

3×10^{19}

ν_μ in direction of Gran Sasso

10^{19}

ν_μ in 100 m^2 at Gran Sasso

3×10^{14}

ν_μ events per day in OPERA

≈ 2500

ν_τ events (from oscillation)

≈ 2




Opera Experiment at Gran Sasso

Basic unit: brick

56 Pb sheets + 56 photographic films (emulsion sheets)

Lead plates: massive target


Emulsions: micrometric precision



Opera Experiment at Gran Sasso



31 target planes / supermodule In total: 206336 bricks, 1766 ton




First observation of CNGS beam neutrinos : August 18th, 2006

Opera Experiment at Gran Sasso




Second Super-module



Scintillator planes 5900 m^2
8064 7m long drift tubes


Details of the first spectrometer




3050 m^2 Resistive Plate Counters
2000 tons of iron for the two magnets

Opera Experiment at Gran Sasso

The Brick Manipulator System (BMS) prototype:
a lot of fun for children and adults !



Tests with the prototype wall





The robotised "Ferrari" for
insertion/extraction of bricks with
vacuum grip by Venturi valve




"Carousel" brick dispensing
and storage system

First Tau Candidate seen a few weeks ago !





AMANDA



AMANDA

Antarctic Muon And Neutrino Detector Array

AMANDA





South Pole



AMANDA


AMANDA-II

Depth



top view

200 m



**Photomultipliers in the Ice,
looking downwards.
Ice is the detecting medium.**





AMANDA

Look for upwards going Muons from Neutrino Interactions.



Cherenkov Light propagating through the ice.

→ Find neutrino point sources in the universe !




AMANDA

Event Display



Up to now: No significant point sources but just neutrinos from cosmic ray interactions in the atmosphere were found .

→ Ice Cube for more statistics !






AMS

Alpha Magnetic Spectrometer


Try to find Antimatter in the primary cosmic rays.
Study cosmic ray composition etc. etc.

AMS

Will be installed on the space station.



AMS



AMS

