

First ever α -scattering with radioactive ion beams:

a novel technique to determine α -nuclear
potentials away from stability



Ciências
ULisboa

ISOLDE Workshop and Users meeting


Wednesday, 3rd December 2025

Francisco G. Barba



Outline

- Why study α -nuclear potentials?
- Challenges of the measurement
- Setup and benchmark of technique
- Status of ${}^4\text{He}({}^A\text{Sn}, {}^4\text{He}){}^A\text{Sn}$ analysis with exotic Sn



Motivation


Solar Abundance Distribution (SAD)

Bulk of elements:

- BBN
- Fusion phase


Heavy elements:

- **Neutron capture** processes




Motivation

p-nuclei





Motivation

Sensitivity studies of gamma Process



Motivation


α -nuclear potentials: elastic scattering



Motivation

α -nuclear potentials: elastic scattering



Mass dependence in stable Sn isotopes



D. Galaviz *et al.*,
PRC71, 0650802 (2005)

Motivation

α -nuclear potentials: elastic scattering




Si/He Targets



Magnetron Sputtering Si/He thin films


Element	Si	He	C	N	O	H
atoms/cm ² (x10 ¹⁵)	2820	1280	100	390	380	475




Exotic
beam



Self-supported Si:He target



RBS spectrum of Si/He target using 2.0 MeV protons and 165° scattering angle




SEM cross section of the Si/He target

Si/He Targets




Magnetron Sputtering Si/He thin films


Element	Si	He	C	N	O	H
atoms/cm ² (x10 ¹⁵)	2820	1280	100	390	380	475



Self-supported Si:He target



RBS spectrum of Si/He target using 2.0 MeV protons and 165° scattering angle




SEM cross section of the Si/He target

Si/He Targets



Magnetron Sputtering
Si/He thin films



Elastic Scattering in
inverse kinematics


Element	Si	He	C	N	O	H
atoms/cm ² (x10 ¹⁵)	2820	1280	100	390	380	475




Exotic
beam



Self-supported Si:He target




RBS spectrum of Si/He target using 2.0 MeV protons and 165° scattering angle




SEM cross section of the Si/He target


Setup



Setup




Setup




Beam

Setup

Telescope A
19° to 51°




Setup



Telescope A
19° to 51°

Detector B
46° to 65°

Setup



Telescope A
19° to 51°

DSSD 300
μm

DSSD 300 μm +
PAD 1500 μm

DSSD 1000 μm

A

B

C


Detector C
60° to 78°

Detector B
46° to 65°

Beam

IS698 Experiment

Measurement of the ${}^4\text{He}({}^A\text{Sn}, {}^4\text{He}){}^A\text{Sn}$ in inverse kinematics at the same $E_{\text{c.m.}}$
mass dependence of α -nuclear potentials along the Sn isotopic chain



Beam and targets

Pilot Beam

$$E_{\text{Beam}} = 4.9 \text{ MeV/u}$$

Cocktail of ^{12}C , ^{16}O , ^{20}Ne and ^{32}S

Sn Beam


$$E_{\text{Beam}} = 4.9 \text{ MeV/u}$$

^{112}Sn ($I \approx 30 \text{ pA}$)

^{108}Sn ($I \approx 50 \text{ pA}$)

^{110}Sn ($I \approx 80 \text{ pA}$)

^{109}Sn ($I \approx 90 \text{ pA}$)



Targets

$^{197}\text{Au} \approx 300 \mu\text{g}/\text{cm}^2$


$^4\text{He} \approx 2 \times 10^{18} \text{ atoms}/\text{cm}^2$




Experimental Data

${}^4\text{He}({}^A\text{Sn}, {}^4\text{He}){}^A\text{Sn}$ @ 4.9 MeV/u

${}^{112}\text{Sn}$ beam scattering on **Si/He-target**



${}^{108}\text{Sn}$ beam scattering on **Si/He-target**




Before counting...!

Amount of ^4He over time

^4He amount decreases

Need to account the different amount of ^4He on each run

- ★ Normalize the integral of ^4He to the ratio He/Si at 60°



Before counting...!


Amount of ^4He over time

^4He amount decreases

Need to account the different amount of ^4He on each run



- ★ Normalize the integral of ^4He to the ratio He/Si at 60°

$$\alpha_i = \frac{N_{He}^i(60^\circ)}{N_{Si}^i(60^\circ)} \frac{N_{Si}^0(60^\circ)}{N_{He}^0(60^\circ)}$$




Before counting...!

Amount of ^4He over time



p-RBS spectrum of the F3 Si:He in different target points after IS698

He/Si ratio obtained from p-EBS analysis of the F3 Si:He after IS698



$$\alpha_i = \frac{N_{\text{He}}^i(60^\circ)}{N_{\text{Si}}^i(60^\circ)} \frac{N_{\text{Si}}^0(60^\circ)}{N_{\text{He}}^0(60^\circ)}$$

Preliminary results


${}^4\text{He}({}^{112}\text{Sn}, {}^{112}\text{Sn}){}^4\text{He}$ @ 18.9 MeV (CM)

$$\frac{d\sigma_{He}}{d\Omega}(\theta) = \sum_i \left(\frac{N_{He}^i(\theta) / \alpha_i}{N_{Si}^i(60^\circ)} \right) \left(\frac{d\sigma_{Si}}{d\Omega}(60^\circ) \right)_{\text{Ruth}}$$




Preliminary results

${}^4\text{He}({}^{112}\text{Sn}, {}^{112}\text{Sn}){}^4\text{He}$ @ 18.9 MeV (CM)

$$\frac{d\sigma_{He}}{d\Omega}(\theta) = \sum_i \left(\frac{N_{He}^i(\theta) / \alpha_i}{N_{Si}^i(60^\circ)} \right) \left(\frac{d\sigma_{Si}}{d\Omega}(60^\circ) \right)_{\text{Ruth}}$$



Preliminary results



Conclusions

- **Four angular distributions** measured covering a large angular range.
- **Good performance** of high concentration ^4He targets
- **Analysis defined** for all the angular regions.
- **Technique benchmarked** with previously measured data in direct kinematics
- **Preliminary data on radioactive** nuclei showing consistency of the method

Acknowledgments



D. Galaviz, J. Ferrer, A. M. Sánchez-Benítez, L. Acosta, B. Amorim, F. G. Barba, M. L. Benito, M. J. Borge, J. Cederkall, M. M. R. Chisti, M. La Cognata, J. G. Correia, Y. Demane, A. Di Pietro, C. Ducoin, E. Fadakis, A. Fernández, B. Fernandez, J. P. Fernández, P. Figuera, L. M. Fonseca, L. M. Fraile, S. Freeman, V. García, V. Godinho, J. Gómez Camacho, D. Hufschmidt, K. Johnston, U. Koester, L. Lamia, B. Olaizola, A. Perea, R. Pires, A. Rodriguez, S. Rothe, E. Siesling, O. Tengblad, P. Teubig, A. Tumino, M. Xarepe



TECHNISCHE
UNIVERSITÄT
DARMSTADT



LUNDS
UNIVERSITET



Fundação
para a Ciência
e a Tecnologia



RENASCER Project: CERN/FIS-PAR/0009/2021

SCORE Project: EXPL/FIS-NUC/0364/2021



FCT PhD Fellowship: UI/BD/154300/2022

Spanish Project: PID2023-147569NB-C22

BACKUP SLIDES



Detector A

- Separate what is punchthrough (consider Elastic Peak higher energy)
- Projection on Total Energy Axis



Detector A



- Select regions for elastically scattered alpha-particles



Kinematics!

Detector A

- Select regions for elastically scattered alpha-particles



Kinematics!


Detector B

I wish they all could be detectors B....!

Fit Gaus + pol1

Integrate in centroid +/- sigma

Apply counter-bg/counter factor to integral
in **each pixel**





Detector C

Separate Linear bg from quadratic bg

Fit gaus+gaus+pol

Integrate in centroid1-sigma1 and centroid2+sigma2


Apply counter-bg/counter to the amounts in **each pixel**



Backup: Pilot Beam calculated xs




20Ne + 197Au elastic scattering Elab=98 MeV



Detector A – inelastic channels

E(level) (keV)	XREF	J ^π (level)
0.0	ABCDEFGHIJKLMNO PQ	0+
1256.69 4	BCDEFGHIJKLMN PQ	2+
2150.87 5	BC E LMN	2+
2190.81 6	C E H LMNO Q	0+
2247.39 6	BC EF HIJKLMN P	4+
2354.21 6	BC EF H LMN PQ	3-
2476.16 11	BC E H MN Q	2+
2520.70 7	BC EFGHI MN P	4+
2549.22 14	EF HIJK MN	6+
2556.6 3	B	(2+)
2617.62 18	E MN	0+
2721.06 14	B E MN	2+
2756.02 9	B E M	3+


Energy deposited by elastic scattered ⁴He in different Si thicknesses




Detector A

E(level) (keV)	XREF	J ^π (level)
0.0	ABCDEFGHIJKLMNO P Q	0+
1256.69 4	BCDEFGHIJKLMN PQ	2+
2150.87 5	BC E LMN	2+
2190.81 6	C E H LMNO Q	0+
2247.39 6	BC EF HIJKLMN P	4+
2354.21 6	BC EF H LMN PQ	3-
2476.16 11	BC E H MN Q	2+
2520.70 7	BC EFGHI MN P	4+
2549.22 14	EF HIJK MN	6+
2556.6 3	B	(2+)
2617.62 18	E MN	0+
2721.06 14	B E MN	2+
2756.02 9	B E M	3+


Energy deposited by scattered ⁴He in 1050 μm Si for different Sn excited states




- Energy deposited by alphas from elastic scattering at different thicknesses




- Adjusted gain **in pad** and angular shifted to meet the kinematic curves





- Assuming thickness of **1050 um** in the telescope





- Energy deposited by elastic alphas in deltaE detector (300 um)



- Adjusted deltaE thickness to 315 μm and removed angular shift





Angular Calibration

$^{197}\text{Au}(^{20}\text{Ne},^{20}\text{Ne})^{197}\text{Au}$ @ 4.9 MeV/u

Pilot Beam Spectra

