



www.ippp.dur.ac.uk



On-Shell Methods for QCD Matrix Elements

Simon Badger, University of Durham IPPP

IOP HEPP06, Warwick, 11th April 2006

Work in collaboration with Nigel Glover and Valya Khoze.

Talk Outline

- Off-Shell Methods - Feynman Diagrams.
- On-Shell Methods: Recursion Relations.
- On-Shell Methods: MHV Rules.
- Conclusions and Outlook.

Off-Shell Methods: Feynman Diagrams

- Advantages:

- Direct link to Lagrangian.
- Easy to adapt to amplitudes with mass/spin.
- Easy to automate.

[MADEVENT,GRACE]

- Disadvantages:

- Large number of diagrams.

# Number of Particles	4	5	6	7	8	9	10
# Feynman Diagrams	4	25	120	2485	34300	559405	10525900

- High multiplicity final states will be significant at the LHC.
- Large cancellations between diagrams.
- Higher order corrections extremely difficult to evaluate.

Colour Ordering and Helicity Amplitudes

- Extracting group factors from Feynman rules allows exploitation and group symmetries.
- Reduces the number of diagrams that need to be calculated.

# Number of Particles	4	5	6	7	8	9	10
# Feynman Diagrams	4	25	120	2485	34300	559405	10525900
# Colour Structures	3	10	36	133	501	1991	7335


- **Helicity Amplitudes:** Computing matrix elements as complex numbers for fixed spins reduces computation of squared matrix elements.
- Write massless vectors in terms of 2-component spinors:

$$|p]_{\dot{\alpha}} |p\rangle_{\alpha} = \sigma_{\alpha\dot{\alpha}}^{\mu} p_{\mu} \quad \langle pq\rangle [qp] = (p + q)^2$$

Off-Shell Recursion

Amplitudes can be built out of amplitudes with fewer external legs by joining off-shell currents

[Berends, Giele '86-'88]



Off-shell gluons shown in red.


Well suited to numerical computations.

[ALPGEN, HELAC/PHEGAS]

On-Shell Recursion: Tree Level Relation

New recursion relations for tree level gluon amplitudes.

[Britto,Cachazo,Feng hep-th/0412308]



All particles **on-shell**, all propagators $\sim 1/P^2$.

Proved using only simple complex analysis.

[Britto,Cachazo,Feng,Witten hep-th/0501052]


On-Shell Recursion: Proof

Consider a generic scattering amplitude:

$$\mathcal{A}(p_1, \dots, p_n) \rightarrow \mathcal{A}(p_1(z), p_2(z), p_3, \dots, p_n) = \mathcal{A}(z)$$

where $p_1(z) + p_2(z) = p_1 + p_2$ and $p_1(z)^2 = p_2(z)^2 = 0$.

$$0 = \frac{1}{2\pi i} \int_{\gamma} dz \frac{\mathcal{A}(z)}{z} = \mathcal{A}(0) + \sum_{i=3}^{n-1} \sum_{h=\pm} \mathcal{A}_L(z_i) \frac{1}{P_{1,i}^2} \mathcal{A}_R(z_i)$$



- Condition for complex momenta can be solved for massless particles:
 $|1\rangle \rightarrow |1\rangle + z|2\rangle, \quad |2\rangle \rightarrow |2\rangle - z|1\rangle.$
- Holds for more general shifts into complex momenta.
- Relies on vanishing of $\mathcal{A}(z)$ at the $z \rightarrow \infty$ boundary.

On-Shell Recursion: Applications

Generates extremely compact expressions for many amplitudes:

- Works for amplitudes of gluons and quarks. [Luo,Wen hep-th/0501121,hep-th/0502009]
- Applies to massive propagating particles: [SB,Glover,Khoze,Svrček hep-th/0504159]
[Forde,Kosower hep-ph/0507292]
 - Propagator simply changes from $P_{1,i}^2 \rightarrow P_{1,i}^2 - m_P^2$
 - Solutions to constraints for shifting one external massive particle.
- Multi-Vector Bosons Currents and Massive Fermions [SB,Glover,Khoze hep-th/0507173]
[Ozeren,Stirling hep-ph/0603071],[Rodrigo hep-ph/0508138],[Schwinn,Weinzierl hep-th/0601012]


MHV Rules

Maximal Helicity Violating (MHV) amplitudes take a remarkably simple form:


$$\mathcal{A}_n^{(0)}(\dots, i^-, \dots, j^-, \dots) = \frac{\langle ij \rangle^4}{\langle 12 \rangle \langle 23 \rangle \dots \langle n1 \rangle}$$

Following from Witten's twistor string/ $\mathcal{N} = 4$ duality, Cachazo, Svrček and Witten proposed a scalar perturbation theory with the MHV amplitudes as vertices.

[Cachazo, Svrček, Witten hep-th/0403047]



NMHV




N^2 MHV

Lines in **red**(black) indicate **negative**(**positive**) helicity gluons.

MHV Rules

Rules generate simple n -point amplitudes for specific helicity configurations.



- Has been applied to tree level amplitudes with gluons, quarks, Higgs and Vector Bosons

[Georgiou,Khoze hep-th/0404072], [Georgiou,Glover,Khoze hep-th/0407027]

[Dixon,Glover,Khoze hep-th/0411092], [SB,Glover,Khoze hep-th/0412275], [Bern,Forde,Kosower,Mastrolia hep-th/0412275]

- Proved by showing equivalence between MHV method and recursion relation with special shift to complex momenta:

Shift all negative helicity particles: $|\widehat{m}_i] \rightarrow |\widehat{m}_i] + z r_i |\eta]$


This ensures no $\overline{\text{MHV}}$ 3-vertices appear.

[Risager hep-th/0508206]

Outlook

- On-shell methods have provided efficient new ways to calculate tree level matrix elements.
- Methods extremely general and easy to apply to many different gauge theories at tree level.

Progress at NLO - Combining on-shell recursion and unitarity methods to find QCD matrix elements:



- Gluon and Quark “finite” amplitudes. [Bern,Dixon,Kosower hep-ph/0501240+hep-ph/0505055]
- Gluon MHV amplitudes. [Bern,Dixon,Kosower hep-ph/0507005, Forde,Kosower hep-th/0509358]
- MHV rules at 1-loop. [Bedford,Brandhuber,Spence,Travaglini hep-th/0412108]
- Target to find the first six gluon NMHV amplitude.