

Chiral perturbation theory tests

Wednesday 13 June 2012 10:00 (25 minutes)

The NA62 experiment at CERN collected a large sample of charged kaon decays with a low intensity beam and minimum bias trigger conditions in 2007. This allowed measurements of a number of rare decays that are difficult to address in conventional high intensity experiments with highly selective trigger conditions. In particular, large samples of $K^{+-} \rightarrow \pi \gamma \gamma$ and $K^{+-} \rightarrow e \nu \gamma$ decays have been collected, allowing precision tests of the Chiral Perturbation Theory. The status and first results of these analyses are presented.

Summary

The NA62 experiment at CERN collected a large sample of charged kaon decays with a low intensity beam and minimum bias trigger conditions in 2007. This allowed measurements of a number of rare decays that are difficult to address in conventional high intensity experiments with highly selective trigger conditions. In particular, large samples of $K^{+-} \rightarrow \pi \gamma \gamma$ and $K^{+-} \rightarrow e \nu \gamma$ decays have been collected, allowing precision tests of the Chiral Perturbation Theory. The status and first results of these analyses are presented.

Author: FANTECHI, Riccardo (Sezione di Pisa (IT))

Presenter: FANTECHI, Riccardo (Sezione di Pisa (IT))

Session Classification: Kaon Physics

Track Classification: Kaon Physics