

# *Implications of LHC results on LC physics*

***G. Moortgat-Pick, Jenny List, Sabine Riemann***

- **Status LHC results**
- **Impact from LHC findings**
- **Impact from (still) LHC non-findings**
- **Conclusions: possible steps and Outlook**

# *Status LHC results -- in short*

- **Excess compatible with a SM-like Higgs around  $m_H \sim 125$  GeV**
  - Should become clear this year.....
  - Would be an absolute revolution!
- **Limits in SUSY coloured sector (approx.):**
  - $m_g > 700$  GeV,  $m_q > 800$  GeV
  - 3<sup>rd</sup> generation: much weaker
  - Results in  $\Lambda > 31$  TeV in GMSB
- **Limits on  $Z'$ ,  $W'$ :  $\sim 2$  TeV**
- **And more limits on ED, exotics, 4<sup>th</sup> generation etc.**

**Room left for a Linear Collider? Which energy steps?**

# *Status LHC results -- in short*

- **Excess compatible with a SM-like Higgs around**

**$m_H \approx 125$  GeV**

- *Currently many new LC reports under work: CLIC CDR just finished, ILC TDR to be finished in end 2012 ....*
- *Please have a look for collider specific details.*
- *This talk is a personal approach to outline possible first LC steps as reaction on the encouraging LHC results.*
- *Therefore mainly talks of the recent LC Forum (only physics addressed) have been used*
- **And more limits in ED, exotics, 4<sup>th</sup> generation etc.**

**Room left for a Linear Collider? Which energy steps?**

# *Impact from LHC findings*

- If a Higgs at  $m_H=125$  GeV :

The Higgs boson is **a completely new type** of particle that we have never encountered. The properties are **experimentally totally unknown**.

- Rich phenomenology at LHC and a LC


- Distinction SM-Higgs vs. SUSY Higgs or something else
- Precise measurement of couplings and BRs needed
- Trilinear couplings measurement

- *'The properties of the Higgs boson, to be discovered at the LHC, must be thoroughly investigated in a good condition at the ILC'* (K. Kawagoe)

# Which energy?

- $E_{\text{CM}} \sim M_{\text{H}} + 120 \text{ GeV} (\sim 250 \text{ GeV})$ 
  - Precise measurement of the mass and production cross section
  - Determination of the spin
  - Precise measurement of the decay branching fractions
- $E_{\text{CM}} \sim 500 \text{ GeV}$  or higher
  - Measurement of the Higgs self-coupling
    - Essential for the Higgs potential and EWSB mechanism
  - Measurement of Top Yukawa coupling


See talk K. Kawagoe at  
LCForum, Hamburg 2/2012



# Task list of the LC -- some 'new' studies


- Couplings measurement:  $\sqrt{s}=250$  GeV (LOI)  $\sqrt{s}=250$  GeV

Muon Channel



Very Precise Measurement  
S/B = 8 in Peak Region


Electron Channel



Less Precise  
Bremsstrahlung in detector material

Combined:  $\Delta M_H = 0.035$  GeV,  $\Delta \sigma_{HZ} / \sigma_{HZ} = 0.027$

See talk R. Poeschl at LCForum



$$\sigma_{HZ} \sim g_{HZZ}^2$$


=> Precision in  $g_{HZZ}$  coupling 1-2%

Sensitivity to 15% deviations  
SM prediction of cross section

# Task list of the LC -- some 'new' studies


- Couplings measurement:  $\sqrt{s}=250$  GeV (LOI)  $\sqrt{s}=250$  GeV

Muon Channel




Very Precise Measurement  
S/B = 8 in Peak Region

Electron Channel



Less Precise  
Bremsstrahlung in detector material



## Further results:

See talk R. Poeschl  
at LCForum


### Higgs BR results

- $\Delta BR/BR(bb) \sim 3\%$
- $\Delta BR/BR(cc) \sim 9\%$
- $\Delta BR/BR(gg) \sim 10\%$
- Error includes  $\Delta\sigma_{HZ}$

# Exciting Higgs story not yet finished

- Since  $m_H \sim$  free parameter in SM at tree level
  - Crucial relations exist, however, between  $m_{\text{top}}$ ,  $m_W$  and  $\sin^2\theta_{\text{eff}}$
  - If nothing else appears in the electroweak sector, these relations have to be urgently checked in order to
    - a) distinguish between SM and Higgs in BSM models (remember Abdels talk,  $\Delta m_H \sim m_{\text{top}}^4$  !)
    - b) Close the SM picture
- Which strategy should one aim?
  - exploit **precision observables** and check, whether the measured values fit together at quantum level
  - $m_Z, m_W, \alpha_{\text{had}}, \sin^2\theta_{\text{eff}}$  und  $m_{\text{top}}$

# Higgs story has just started ...



**LEP:**

$$\sin^2\theta_{\text{eff}}(A_{\text{FB}}^b) = 0.23221 \pm 0.00029$$

**SLC:**

$$\sin^2\theta_{\text{eff}}(A_{\text{LR}}) = 0.23098 \pm 0.00026$$

**World average:**

$$\sin^2\theta_{\text{eff}} = 0.23153 \pm 0.00016$$

- **Uncertainties from input parameters:  $\Delta m_Z$ ,  $\Delta\alpha_{\text{had}}$ ,  $m_{\text{top}}$**

*Heinemeyer, Weiglein*


- $\Delta m_Z = 2.1 \text{ MeV}$ :  $\Delta \sin^2\theta_{\text{eff}}^{\text{para}} \sim 1.4 \times 10^{-5}$
- $\Delta\alpha_{\text{had}} \sim 10$  (5 future)  $\times 10^{-5}$ :  $\Delta \sin^2\theta_{\text{eff}}^{\text{para}} \sim 3.6$  (1.8 future)  $\times 10^{-5}$
- $\Delta m_{\text{top}} \sim 1 \text{ GeV}$  (Tevatron/LHC):  $\Delta \sin^2\theta_{\text{eff}}^{\text{para}} \sim 3 \times 10^{-5}$
- $\Delta m_{\text{top}} \sim 0.1 \text{ GeV}$  (ILC):  $\Delta \sin^2\theta_{\text{eff}}^{\text{para}} \sim 0.3 \times 10^{-5}$

# To close the story...


$\sqrt{s}=92 \text{ GeV}$

## 1. Measure $\sin^2\theta_{\text{eff}}$ via $A_{\text{LR}}$ , $A_{\text{FB}}$ with high precision


See talk S. Heinemeyer at LCForum



← **LEP value**  
**disfavours both,**  
**SM+MSSM**



↑ **SLD value**  
**disfavours SM**



**World average** →  
**happy with both!**

**Central value has**  
**large impact !!!**

and...

$\sqrt{s}=350 \text{ GeV}$

## 2. measure the top mass

See talk P. Uwer at LCFForum


- LHC: Difficult (if not impossible) to go below  $\Delta m_{\text{top}} < 1 \text{ GeV}$ 
  - Because of non perturbative corrections, intrinsic uncertainty generator mass  $\leftrightarrow$  pole mass, higher order corrections
- Since top quark does not appear as asymptotic state:  $m_{\text{top}}$  is just a parameter, i.e. **renormalisation scheme dependent**
- **only indirect determination possible**
- **Observable should show good sensitivity to  $m$ :**  $\frac{\Delta O}{O} \leftrightarrow \frac{\Delta m_t}{m_t}$
- **LC: Peak position remains stable:  $m_t=100 \text{ MeV}$  estimate still valid!**
- **However: dedicated threshold scan required!**
- **Threshold scan meets all the requirements, theory is still improving**
- **Alternative approaches are in principle possible, however reachable accuracy not yet clear**

and...

$\sqrt{s}=350 \text{ GeV}$

## 2. measure the top mass

See talk P. Uwer at LCForum



Important shift due to non-logarithmic NNNLO terms

- LC: Peak position remains stable:  $m_t=100 \text{ MeV}$  estimate still valid!
- However: dedicated threshold scan required!
- **Threshold scan meets all the requirements, theory is still improving**
- Alternative approaches are in principle possible, however reachable accuracy not yet clear

# Why Higgs self coupling?

- Needed to determine the Higgs potential
- May also be important to distinguish nature of Higgs sector, THDM or SUSY!

$\sqrt{s} \geq 500 \text{ GeV}$

See talk J. Tian at LCForum

Higgs Potential:  $V(\eta_H) = \frac{1}{2} m_H^2 \eta_H^2 + \lambda v \eta_H^3 + \frac{1}{4} \lambda \eta_H^4$


physical Higgs field      mass term      trilinear coupling      quartic Higgs coupling, which is difficult to measure at both LHC and ILC, even SLHC!

SM:  $\lambda = \lambda_{SM} = \frac{m_H^2}{2v^2}$        $v \sim 246 \text{ GeV}$



# Full simulation study under work

See talk J. Tian

$\sqrt{s}=500 \text{ GeV}$




$$\frac{\Delta\lambda}{\lambda} = 1.8 \frac{\Delta\sigma}{\sigma}$$



$$\sigma = a\lambda^2 + b\lambda + c$$

$$\frac{\Delta\lambda}{\lambda} = 0.85 \frac{\Delta\sigma}{\sigma}$$


$\sqrt{s}=1000 \text{ GeV}$



# Full simulation study under work



See talk J. Tian

$\sqrt{s}=500 \text{ GeV}$




$$\frac{\Delta\lambda}{\lambda} = 1.8 \frac{\Delta\sigma}{\sigma}$$

$$\sigma = a\lambda^2 + b\lambda + c$$





$$\frac{\Delta\lambda}{\lambda} = 0.85 \frac{\Delta\sigma}{\sigma}$$

$\sqrt{s}=1000 \text{ GeV}$



# Impact from (still) LHC non-findings

- SUSY: still strongly motivated and beautiful, but
  - so far, nothing, only rather heavy exclusion limits in the coloured sector
  - just **constrained models** (mSUGRA, CMSSM) under tension !



- What can a LC do? Which further input do we have?

# What could be the 1<sup>st</sup> step in such scenarios?

- Assume only Higgs@LHC but no hints for SUSY

- Really SM?

- Help from  $\sin^2\theta_{\text{eff}}$ ?

- If GigaZ precision:


- i.e.  $\Delta m_{\text{top}}=0.1$  GeV...

- Deviations measurable

- $\sin^2\theta_{\text{eff}}$  can be the crucial quantity to outline the scale!

$\sqrt{s}=92$  GeV

Heinemeyer, Weber, Weiglein



# Further SUSY facts

- Please remember (see talk, S. Heinemeyer, yesterday):
  - Low energy experiments,  $(g-2)_\mu$ , favour rather low SUSY masses in the electroweak sector:

$$\delta a_\mu(\text{N.P.}) = \mathcal{O}(C) \left( \frac{m_\mu}{M} \right)^2, \quad C = \frac{\delta m_\mu(\text{N.P.})}{m_\mu}$$

See talk D. Stoeckinger  
at LCForum

- C very model dependent, SUSY/ED  $\sim \mathcal{O}(\alpha/4\pi \dots)$
- LHC results prefer rather heavy coloured sector in 1<sup>st</sup> +2<sup>nd</sup> generation ( $m_g > 800$  GeV)
- Way out: rather simple
  - Either just decouple uncoloured and coloured sector and take hybrid models of SUSY breaking
  - Just leave out the constrained minimal models, that's all

*Personal remark: I am not worried at all, just enjoy LHC results*



# Further play: new TDR benchmark

$\sqrt{s}=500 \text{ GeV}$

Berggren, List, Rolbiecki


$m_H=125 \text{ GeV}$ :

$\tan\beta=10$

$M_2=225 \text{ GeV}$

$\mu=200 \text{ GeV}, \dots$

- Wonderful spectrum with  
rich phenomenology!



***There is not really a problem with current LHC bounds vs SUSY !***

# One gravitino DM induced example

- See Sergei's talk later today: nightmare (LHC) scenario


- Gravitino LSP (DM)
- Neutralino is the NLSP and Higgsino-like
- Charginos are mass-degenerate with Neutralinos
- Higgs mass around 125 GeV
- All colored particles are inaccessible at the LHC

Light chargino and neutralinos  $\sim 130$  GeV,  
 Further scalars  $> 600$  GeV  
 Gluino etc,  $> 1$  TeV

- Apply in future LC features to such a scenario:

- Precise fundamental SUSY parameter determination  $M_1, M_2, \mu, \tan\beta$  possible, if only  $m_{\chi_{01}}, m_{\chi_{02}}$  and  $m_{\chi_{\pm 1}}$  is accessible
- Required features: polarized beams, different beam energies, threshold scans

Choi, SY ea



# Assume hints for SUSY at LHC: task List LC

- Precise determination of fundamental MSSM parameters

- Independent from the SUSY breaking scheme
- Since masses and cross section O(% - ‰) level achievable, parameter determination also O(%)

- Inclusion of **electroweak higher order corrections** required

See A. Bharucha talk at LCForum

- Already under work now:

- Why already now? Study, i.e. how strong the impact of mass measurements via **threshold scans** is etc.

- Provide **accurate predictions** for heavier states!

e+e- to  $\chi^+_1 \chi^-_1$

Parameter	NLO result $\pm 1\sigma (\pm 2\sigma)$	LO result $\pm 1\sigma$
$M_1 / \text{GeV}$	$125.0 \pm 0.6 (\pm 1.2)$	$122.0 \pm 0.5$
$M_2 / \text{GeV}$	$250.0 \pm 1.6 (\pm 3.0)$	$260.7 \pm 1.4$
$\mu / \text{GeV}$	$180.0 \pm 0.7 (\pm 1.3)$	$176.5 \pm 0.5$
$\tan \beta$	$10.0 \pm 1.3 (\pm 2.6)$	$27.0 \pm 9.0$
$m_{\tilde{\nu}} / \text{GeV}$	$1500 \pm 20 (\pm 40)$	$2230 \pm 50$
$m_{\tilde{t}_2} / \text{GeV}$	$800^{+220}_{-170} (+540_{-280})$	–

# Task List cont.

- **Goal: reveal the 'structure' of NP**
  - **Prove all predicted properties: couplings, mass relations, CP properties, spin, quantum numbers etc.**
  - **For some properties as e.g. spin verification, very good methods have been developed for the LHC** (*remember methods of Barr and of Rolbiecki, Tattersall etc.*)
- **However, 'model independence' and 'accuracy at quantum level' requires a LC**
  - *Remember the results of the LHCLC report, >400 pages,..... still valid, nothing has really changed since then*

# *Conclusions: possible steps*

- LHC results are strongly encouraging !
- If new particle at  $m_{\gamma}=125$  GeV confirmed:
  - rich phenomenology for both LHC and LC
- Reasonable steps:

## *‘elementary’*

- $\sqrt{s}=250$  GeV, 350 GeV and 92 GeV (do not forget, it might be crucial already now!)

## *‘graduate’*

- $\sqrt{s}=500$  GeV, 1 TeV (*if hints at LHC for NP found*)

## *‘post-graduate’*

- $\sqrt{s}=1.5$  TeV and more (*if eg Z', ED, SUSY at LHC found*)

# and Outlook

- **Physics prospects couldn't be better.....**
  - Maybe a Higgs at 125 GeV
  - LHC runs perfectly well
  - LC is ready to start
  - Encouraging activities in Japan towards a LC
- **A LCreview (*physics!*) from 92 GeV to multi-TeV under work**
  - Please have a look on [lcforum.desy.de](http://lcforum.desy.de) (more details on the web or ask S. Riemann, I. Fleck, F. Simon and myself)
  - Feel invited and encouraged to submit LCnotes (ILC,CLIC) on:  
[www-flc.desy.de/lcnotes](http://www-flc.desy.de/lcnotes)


***This year is crucial to get answers to our physics questions!***

# R-parity violation

- Much lower mass bounds in such models:
- RPV often leads to displaced vertices
- Dedicated simulations also at LC

## Conclusion

- R-parity violation is an interesting alternative to “normal” SUSY
- bRPV could explain neutrino parameters
- some RPV LHC searches ongoing
- especially for high squark masses: sensitivity of ILC expected to be better



See B. Vormwald talk at LCForum