

# Higgs and SFitter



- Introduction
- 7 TeV results
- 14 TeV results
- Conclusions

LHC2TSP  
CERN  
March 28, 2012  
Dirk Zerwas  
LAL Orsay



On behalf of:

SFitter: R.Lafaye, T.Plehn, M.Rauch, D.Zerwas  
and Higgs friends: C. Englert, M. Duehrssen, M.Klute,  
P.Zerwas



Measuring the Higgs Sector, R. Lafaye, T. Plehn, M. Rauch, D. Zerwas, M. Dührssen  
Published in JHEP 0908 (2009) 009.

Measuring Hidden Higgs and Strongly-  
Interacting Higgs Scenarios.  
S.Bock, R. Lafaye, T. Plehn, M. Rauch, D. Z.,  
P.M. Zerwas  
Phys.Lett. B694 (2010) 44-53

# The Higgs sector: 14TeV

**Gluon fusion and VBF in well defined final states** (many authors and papers)

**Example for  $M_H=120\text{GeV}$**

production	decay	$S + B$	$B$	$S$	$\Delta S^{(\text{exp})}$	$\Delta S^{(\text{theo})}$
$gg \rightarrow H$	$ZZ$	13.4	6.6 ( $\times 5$ )	6.8	3.9	0.8
$qqH$	$ZZ$	1.0	0.2 ( $\times 5$ )	0.8	1.0	0.1
$gg \rightarrow H$	$WW$	1019.5	882.8 ( $\times 1$ )	136.7	63.4	18.2
$qqH$	$WW$	59.4	37.5 ( $\times 1$ )	21.9	10.2	1.7
$t\bar{t}H$	$WW(3\ell)$	23.9	21.2 ( $\times 1$ )	2.7	6.8	0.4
$t\bar{t}H$	$WW(2\ell)$	24.0	19.6 ( $\times 1$ )	4.4	6.7	0.6
inclusive	$\gamma\gamma$	12205.0	11820.0 ( $\times 10$ )	385.0	164.9	44.5
$qqH$	$\gamma\gamma$	38.7	26.7 ( $\times 10$ )	12.0	6.5	0.9
$t\bar{t}H$	$\gamma\gamma$	2.1	0.4 ( $\times 10$ )	1.7	1.5	0.2
$WH$	$\gamma\gamma$	2.4	0.4 ( $\times 10$ )	2.0	1.6	0.1
$ZH$	$\gamma\gamma$	1.1	0.7 ( $\times 10$ )	0.4	1.1	0.1
$qqH$	$\tau\tau(2\ell)$	26.3	10.2 ( $\times 2$ )	16.1	5.8	1.2
$qqH$	$\tau\tau(1\ell)$	29.6	11.6 ( $\times 2$ )	18.0	6.6	1.3
$t\bar{t}H$	$b\bar{b}$	244.5	219.0 ( $\times 1$ )	25.5	31.2	3.6
$WH/ZH$	$b\bar{b}$	228.6	180.0 ( $\times 1$ )	48.6	20.7	4.0

**Duehrssen et al.: Phys.Rev.D70:113009,2004.  
hep-ph/0406323**

**$t\bar{t}H \rightarrow b\bar{b}$ : 50% signal reduction**

**Hbb: J. M. Butterworth, A. R. Davison, M. Rubin, G. P. Salam Phys.Rev.Lett.100:242001,2008  
ATLAS: ATL-PHYS-PUB-2009-088.**

## Experimental Errors

Measurement of luminosity	5 %
Detector efficiency	2 %
Lepton reconstruction efficiency	2 %
Photon reconstruction efficiency	2 %
WBF tag-jets / jet-veto efficiency	5 %
$b$ -tagging efficiency	3 %
$\tau$ -tagging efficiency (hadronic decay)	3 %
Lepton isolation efficiency (decay $H \rightarrow ZZ \rightarrow 4l$ )	3 %

## Theory Errors

$\sigma$ (gluon fusion)	13 %	$\text{BR}(H \rightarrow ZZ)$	1 %
$\sigma$ (weak boson fusion)	7 %	$\text{BR}(H \rightarrow WW)$	1 %
$\sigma$ ( $VH$ -associated)	7 %	$\text{BR}(H \rightarrow \tau\tau)$	1 %
$\sigma$ ( $t\bar{t}$ -associated)	13 %	$\text{BR}(H \rightarrow c\bar{c})$	4 %
		$\text{BR}(H \rightarrow b\bar{b})$	4 %
		$\text{BR}(H \rightarrow \gamma\gamma)$	1 %
		$\text{BR}(H \rightarrow Z\gamma)$	1 %
		$\text{BR}(H \rightarrow gg)$	2 %

# The Higgs sector: 7TeV


ATLAS	WW	0-jet
	WW	1-jet
CMS	$\gamma\gamma$	
	$ZZ \rightarrow 4\ell$	
	$\gamma\gamma$	di-jet
	$ZZ \rightarrow 4\ell$	
	WW	0-jet
	WW	1-jet
	WW	2-jet
	$\tau\tau$	0/1-jet
	$\tau\tau$	Boosted
	$\tau\tau$	VBF
	$b\bar{b}$	WH
	$b\bar{b}$	$Z(\rightarrow \ell\bar{\ell})H$
	$b\bar{b}$	$Z(\rightarrow \nu\bar{\nu})H$

Cross checks performed (A.Read's comment):

- exclusion
- signal strength

Correlated measurements and parameters: apply SUSY search techniques for parameter extraction

Difficulty: convolution of Gaussian+Poisson+Flat errors




# The coupling definitions

**Definition:**  $\Delta X$  deviation of  $XXH$  coupling from SM value:

$$g_X \rightarrow g_X^{\text{SM}} (1 + \Delta X)$$

**Loop induced coupling:**


$$g_X \rightarrow g_X^{\text{SM}} (1 + \Delta X^{\text{SM}} + \Delta X)$$



Add  $\Delta g$  and  $\Delta \gamma$ : sign preference power of  $H \rightarrow \gamma\gamma$  disappears

**Common assumptions:**

- fermion universality (charm)
- no invisible decays





- general positive correlation among non-Hbb couplings due to total width  $\approx$  Hbb
- frequentist approach better adapted (no real secondary minima)

**Measurements at LHC:**

$$\sigma \cdot \text{BR} \cdot L \cdot \sim g^2 \cdot g^2 / \Gamma$$

**blind to simultaneous coupling/ $\sqrt{\text{width}}$  changes:**



# Hidden sector: 14 TeV

Patt, Wilczek

## Higgs portal:


- hidden sector
- all widths modified by the same parameter
- set  $\Gamma_{\text{hid}}=0$

$$\sigma = \cos^2 \chi \sigma^{\text{SM}}$$


$$\Gamma_{\text{vis}} = \cos^2 \chi \Gamma_{\text{vis}}^{\text{SM}}$$

$$\Gamma_{\text{inv}} = \cos^2 \chi \Gamma_{\text{inv}}^{\text{SM}} + \Gamma_{\text{hid}} .$$


$\cos^2 \chi = 1$



$\cos^2 \chi = 0.8$



$\cos^2 \chi = 0.6$




well reconstructed (step=10GeV)

Typical precision:  
10-20%  
better than cpl

Non-discovery of Higgs (<140GeV) does not mean non-existence (Englert et al, Phys.Lett. B703 (2011) 298-305 ):

	# observable Higgs bosons	
$\sin^2 \chi \lesssim 0.2$	1	SM-type Higgs $H_1$ ( $\sigma_{H_1} \geq 3, \sigma_{H_2} \leq 1$ )
$0.3 \lesssim \sin^2 \chi \lesssim 0.4$	0	neither SM-type nor hidden-type Higgs $H_1, H_2$ ( $\sigma_{H_1, H_2} < 3$ )
$0.4 \lesssim \sin^2 \chi \lesssim 0.6$	2	SM + hidden-type Higgs $H_1, H_2$ ( $\sigma_{H_1, H_2} \geq 3$ )
$\sin^2 \chi \gtrsim 0.6$	1	hidden-type Higgs $H_2$ ( $\sigma_{H_1} \leq 1, \sigma_{H_2} \geq 3$ )

# The Higgs sector precision: 7 TeV



## Parameters:


- $\Delta_W$   $\Delta_Z$   $\Delta_t$   $\Delta_b$   $\Delta_\tau$
- effective couplings hopeless

## 2.1-4.9fb<sup>-1</sup>: pre-Moriond

- all errors on couplings <1
- portal precision: 15%


## Extrapolation to 20fb<sup>-1</sup>

- most errors on couplings <0.5
- portal precision 10%
- scaling roughly statistical as expected




# Higgs sector: 14 TeV

14TeV no effective couplings:




14TeV with effective couplings:



- no effective couplings lead to a slight increase of precision
- coupling precision in general better than about 0.3
- portal precision better than 0.1

# The Higgs sector precision: zoom on 125 GeV



$\Delta_{Z/W}$   $\Delta_{\tau/b}$   $\Delta_{b/W}$ :

- coupling ratio
- error reduced also for 7TeV:  
+correl with  $\Delta_b$

$\Delta_H$ :

- one parameter fit
- best precision

$\Delta_Z$   $\Delta_t$   $\Delta_b$   $\Delta_\tau$  :

- direct coupling
- +correl with  $\Delta_b$

$\Delta_\gamma$   $\Delta_g$ :



- effective coupling
- additional contribution BSM
- 14TeV only

# The Higgs sector precision: far far future

For Francois Lediberder et al.

**3000fb<sup>-1</sup>:**

- extrapolate blindly
- full set including effective couplings
- all errors on couplings <20%
- most better than: 15%
- best at 10%



**3000fb<sup>-1</sup>:**

- no effective couplings
- all errors on couplings <15%
- most better than: 10%
- best at 7%


• portal:

• 5%

• reached theory error

• theory error on Xsection (cpl 0.5)

# Conclusions



- couplings known to better than 0.5 ☺
- portal precision: 10%
- post-Moriond being prepared