

A. Azatov, O. Bondu, A. Falkowski, M. Felcini, S. Gascon-Shotkin,  
D. K. Ghosh, G. Moreau, A. Y. Rodriguez-Marrero, and S. Sekmen

# **HIGGS BOSON PRODUCTION VIA NEW VECTOR-LIKE TOP QUARK DECAYS AND DETECTION AT THE LHC**

# To appear in the arXiv on March 30th

Preprint also uploaded to the agenda of this session

<https://indico.cern.ch/getFile.py/access?contribId=41&sessionId=5&resId=0&materialId=slides&confId=162621>

## Higgs boson production via new vector-like top quark decays and detection in diphoton or multilepton plus multijets channels at the LHC

A. Azatov,<sup>1</sup> O. Bondu,<sup>2</sup> A. Falkowski,<sup>3</sup> M. Felcini,<sup>4</sup> S. Gascon-Shotkin,<sup>2</sup>  
D. K. Ghosh,<sup>5</sup> G. Moreau,<sup>3</sup> A. Y. Rodríguez-Marrero,<sup>4</sup> and S. Sekmen<sup>6</sup>

<sup>1</sup>*Dipartimento di Fisica, Università di Roma "La Sapienza", INFN Sezione, I-00185 Roma, Italy*

<sup>2</sup>*Université de Lyon, Université Claude Bernard Lyon 1, CNRSIN2P3,  
Institut de Physique Nucléaire de Lyon, F-69622 Villeurbanne Cedex, France*

<sup>3</sup>*Laboratoire de Physique Théorique, Bât. 210, CNRS,  
Université Paris 11, F-91405 Orsay Cedex, France*

<sup>4</sup>*Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, E-39005 Santander, Spain*

<sup>5</sup>*Department of Theoretical Physics, Indian Association for the Cultivation of Science Kolkata 700 032, India*

<sup>6</sup>*CERN Physics Department, CH-1211 Geneva 23, Switzerland.*

Motivated by the recent stringent constraints from the search for direct Higgs boson production at the LHC, over a wide Higgs mass range, we first build a minimal model of vector-like quarks where the dominant Higgs production process at the LHC, via gluon fusion, can be significantly suppressed. Within this model, compatible with the present experimental constraints, we demonstrate that the Higgs production via a heavy vector-like quark,  $t_2$ , decay,  $pp \rightarrow t_2 \bar{t}_2$ ,  $t_2 \rightarrow th$ , allows to discover a Higgs boson and measure its mass, through the decay channels  $h \rightarrow \gamma\gamma$  or  $h \rightarrow ZZ$ . We also comment on the recent hint in LHC data from a possible  $\sim 125$  GeV Higgs, in the presence of heavy vector-like top quarks.


PACS numbers:

### I. INTRODUCTION


One of the primary goals of the Large Hadron Collider (LHC) is the direct search for the cornerstone of the Standard Model (SM), namely the Higgs boson, or of any signal from alternative Electro-Weak Symmetry Breaking (EWSB)

6 [hep-ph] 29 Mar 2012

# Higgs status: may be... may be not...



Physics Letters B 710 (2012) 49–66



Physics Letters B 710 (2012) 26–48

By end of this year

If a SM like Higgs observed -> ascertain its origin at LHC 14 TeV-> keep looking BSM

If a SM like Higgs excluded -> -----> keep looking BSM

BSM search. including the Higgs sector, very important  
irrespective of the Higgs outcome of this year

→ keep investigating BSM Higgs signatures

→ Higgs produced in the decays of BSM heavy particles

→ Discover Higgs and BSM at once

# Higgs production via heavy top decay

BSM theoretical model with **new heavy top-like states**:


$t_3, t_2, t_1$

$m_{t_3} > m_{t_2} > m_{t_1}$

$t_1 = \text{top quark} ; m_{t_1} = m_t$

In this model, consider the **production and decay of the lightest of the two heavy new states  $t_2$**

$pp \rightarrow t_2 t_2$  cross-section




**Production:**  $pp \rightarrow t_2 t_2$

**Decays:**  $t_2 \rightarrow th, t_2 \rightarrow tZ, t_2 \rightarrow bW$


# Heavy top decays – experimental searches

$pp \rightarrow t_2 t_2 \quad t_2 \rightarrow bW$

$pp \rightarrow t_2 t_2 \quad t_2 \rightarrow tZ$



CERN-PH-EP/2012-081



PRL 107, 271802 (2011)

# Theory framework and benchmarks

see preprint

Parameter Set	A1	A2	B	C	D
$Y / \dot{Y}$	-1.43 / 2	1.02 / -0.1	1.15 / 0.4	1.12 / -0.5	1.05 / -0.3
$Y' / Y''$	1.85 / -1	1 / 0.55	-1.5 / 1.6	1.1 / 1.65	1.7 / 1.9
$m / m'$ (GeV)	0 / 370	0 / 675	0 / 770	0 / 810	80 / 1100
$m''$ (GeV)	510	645	980	850	1100
$m_{t_3}$ (GeV)	722	804	1181	1125	1454
$m_{t_2}$ (GeV)	403	599	626	572	788
$m_h$ (GeV)	125	125	255	320	540
$\sigma_{gg \rightarrow h}^{SM}$ (pb) @ 7 TeV	15.31	15.31	3.18	2.25	0.58
$\sigma_{gg \rightarrow h}^{SM}$ (pb) @ 14 TeV	49.85	49.85	13.50	10.59	3.85
$\sigma_{gg \rightarrow h}^{t'}/\sigma_{gg \rightarrow h}^{SM}$	1.27	1.31	0.45	0.40	0.65
$\sigma_{t_1 t_1 h}^{t'}$ (pb) @ 7 TeV	0.0194	0.076	0.0037	0.0016	$7 \cdot 10^{-4}$
$\sigma_{t_1 t_1 h}^{t'}$ (pb) @ 14 TeV	0.138	0.0539	0.036	0.021	0.015
$\sigma_{t_2 t_2}$ (pb) @ 7 TeV	1.361	0.0936	0.0709	0.136	0.0115
$\sigma_{t_2 t_2}$ (pb) @ 14 TeV	13.53	1.465	1.164	1.975	0.284
$B_{t_2 \rightarrow t_1 h}$ (%)	62.6	82.1	60.8	13.5	43.0
$B_{t_2 \rightarrow t_1 Z}$ (%)	28.6	14.7	25.0	46.1	40.3
$B_{t_2 \rightarrow bW}$ (%)	8.8	3.2	14.2	40.4	16.6
$\Gamma_{t_2}$ (GeV)	4.4	3.5	19.8	6.5	8.8
$\sigma_{i_2 t_2} B_{i_2 \rightarrow bW}^2$ (pb)	0.01	$8 \cdot 10^{-5}$	0.001	0.022	0.000(3)
LHC bound [72]	< 0.26	< 0.14	< 0.14	< 0.16	×
$\sigma_{i_2 t_2} B_{i_2 \rightarrow t_1 Z}^2$ (pb)	0.11	0.002	0.004	0.029	0.002
LHC bound [73]	< 0.5	< 0.4	< 0.4	< 0.4	×
$S / T$	0.05 / 0.05	0.03 / 0.03	-0.01 / 0.23	-0.01 / 0.30	-0.01 / 0.28

light  
Higgs  
allowed

heavier  
Higgs  
allowed

# Search strategy

$pp \rightarrow t_2 t_2 \rightarrow th \text{ bW}, thtZ, th th$

consider

$h \rightarrow \gamma\gamma$  and  $h \rightarrow ZZ \rightarrow 4l$

decays



## 14 TeV

Signal event Yield per fb<sup>-1</sup>

Parameter set	$m_{t_2}$ (GeV)	$m_h$ (GeV)	$\sigma_{\bar{t}_2 t_2}$ (fb)	$B_{t_2 \rightarrow th}$	$B_{t_2 \rightarrow tZ}$	$B_{t_2 \rightarrow bW}$	$B_{h \rightarrow VV}^{SM}$	$f_{h \rightarrow VV}^{t'/SM}$	$Y_S(h \rightarrow VV)$ (fb)		
									$thbW$	$thtZ$	$thth$
Point A1	403	125	$1.353 \cdot 10^4$	0.626	0.286	0.088	$2.29 \cdot 10^{-3}$	0.91	3.11	10.1	22.1
Point A2	599	125	$1.465 \cdot 10^3$	0.821	0.147	0.032	$2.29 \cdot 10^{-3}$	0.90	0.159	0.729	4.07
Point B	626	255	$1.164 \cdot 10^3$	0.608	0.25	0.142	0.298	1.00	59.9	105	256
Point C	572	320	$1.975 \cdot 10^3$	0.135	0.461	0.404	0.309	1.00	66.6	76.0	22.2
Point D	788	540	$0.284 \cdot 10^3$	0.43	0.403	0.166	0.265	1.06	11.4	27.6	29.5

see preprint



# $t_2 t_2 \rightarrow th \text{ bW}, thtZ, th th, h \rightarrow ZZ \rightarrow 4l$



see  
preprint

Parameter set	Signal: $thbW/thtZ/thth, h \rightarrow ZZ$				Total signal $S, M_{4l}(\text{GeV})$ cut			
	$Y_S$ (fb)	$N_{4l}$	$N_{N_j, H_T}$	$N_{2b}$	no mass cut	> 200	> 300	> 500
Point B	59.9 / 105 / 256	8.1 / 23.7 / 37.8	5.5 / 18.0 / 28.9	3.52 / 11.5 / 18.5	33.4	30.4	17.0	4.9
Point C	66.6 / 76.0 / 22.2	7.7 / 15.8 / 3.1	4.8 / 11.3 / 2.2	3.07 / 7.23 / 1.41	11.7	10.7	7.4	2.0
Point D	11.4 / 27.6 / 29.5	1.5 / 6.6 / 4.7	1.1 / 5.3 / 3.9	0.70 / 3.40 / 2.50	6.6	6.2	5.1	2.7
Background					Total background $B, M_{4l}(\text{GeV})$ cut			
Process	$\sigma(\text{fb})$	$N_{4l}$	$N_{N_j, H_T}$	$N_{2b}$	no mass cut	> 200	> 300	> 500
$t\bar{t} + \text{jets}$	$9.19 \cdot 10^5$	4680	1480	27.5	35.5	23.5	13.2	6.8
$t\bar{t}b\bar{b} + \text{jets}$	$2.50 \cdot 10^3$	5.60	3.10	2.0	Statistical $\Delta B, M_{4l}(\text{GeV})$ cut			
$t\bar{t}W + \text{jets}$	$1.99 \cdot 10^2$	1.20	0.40	0.036	no mass cut	> 200	> 300	> 500
$t\bar{t}Z + \text{jets}$	97.3	25.0	9.5	6.0	6.0	4.9	3.6	2.6

# $t_2 t_2 \rightarrow th \text{ bW}, thtZ, th th, h \rightarrow \gamma\gamma$



see preprint

Parameter set	Signal: $thbW/thtZ/thth, h \rightarrow \gamma\gamma$				Total signal $S, M_{2\gamma}(\text{GeV})$ cut	
	$Y_S$ (fb)	$N_{2\gamma}$	$N_{N_1>6}$	$N_{N_1>8}$	> 90	[115,135]
Point A1	3.1 / 10.1 / 22.1	5.4 / 15.4 / 30.4	4.1 / 12.7 / 27.1	1.4 / 6.2 / 14.5	22.1	17.7
Point A2	0.16 / 0.73 / 4.1	0.34 / 1.35 / 6.98	0.27 / 1.17 / 6.6	0.10 / 0.62 / 3.8	4.5	3.2
Background					Tot, background $B, M_{2\gamma}(\text{GeV})$ cut	
Process	$\sigma(\text{fb})$	$N_{M_{2\gamma}>90}$	$N_{N_1>6}$	$N_{N_1>8}$	> 90	[115,135]
$W\gamma\gamma + \text{jets}$	450	422	110	19.6	27.8	4.38
$t\bar{t}\gamma\gamma + \text{jets}$	15.5	11.8	11.3	8.18	Statistical $\Delta B, M_{2\gamma}(\text{GeV})$ cut	
$t\bar{t}W\gamma\gamma + \text{jets}$	0.0678	0.0577	0.0515	0.0272	> 90	[115,135]
					5.3	2.1

# Conclusions

- **New possibilities** to discover a Higgs boson and/or measure its properties  
 $pp \rightarrow t\bar{t} \rightarrow th \text{ bW, thtZ, th th}$   
with  
 $h \rightarrow \gamma\gamma$  and  $h \rightarrow ZZ \rightarrow 4l$  decays
- **Investigate Higgs boson production** over the whole mass range  $\sim 120\text{-}600$  GeV
- **Discover** ( $5\sigma$ ) at the **14 TeV LHC** with luminosities  $\sim 20 - 200 \text{ fb}^{-1}$
- **Measure Higgs and heavy top** masses (+other properties) with  $200 \text{ fb}^{-1}$  or more
- **For details see**

Preprint on arXiv March 30th or

<https://indico.cern.ch/getFile.py/access?contribId=41&sessionId=5&resId=0&materialId=slides&confId=162621>