

## Higgs seaches in CMS

Irakli Svintradze for CMS Collaboration

*Department of Physics, Kansas State University, Manhattan, KS, USA*

---

### Abstract

We present the results of the standard model Higgs boson search in proton-proton collisions at 7 and 8 TeV center of mass energy collected by the Compact Muon Solenoid experiment at the Large Hadron Collider. Data correspond to integrated luminosity of about  $5 \text{ fb}^{-1}$  at 7 TeV and another  $5 \text{ fb}^{-1}$  at 8 TeV. Five decay modes of the Higgs boson are analyzed:  $\gamma\gamma$ ,  $ZZ$ ,  $WW$ ,  $\tau^+\tau^-$ , and  $b\bar{b}$ . The observed excess has a significance of  $5.0 \sigma$  standard deviations at the mass of about 125 GeV, while the expected significance of the standard model Higgs boson at this mass is  $5.8 \sigma$ . Analysis of  $\gamma\gamma$  and  $ZZ$  decay channels, that offer the best mass resolution, yield the measurement of the new particle's mass to be  $125.3 \pm 0.4(\text{stat.}) \pm 0.5(\text{syst.}) \text{ GeV}$ .

**Keywords:** CMS, physics, Higgs

---

### 1. Introduction

Within the standard model (SM), particles acquire mass through interaction with the Higgs field [1, 2, 3, 4, 5, 6, 7, 8, 9]. The precision electroweak measurements imply that the mass of the Higgs boson  $m_H$  is below 152 GeV at 95% confidence level (CL) [10]. For a few decades scientists have been searching for this elusive particle: direct searches performed by LEP collider ruled out the Higgs boson with  $m_H < 114.4 \text{ GeV}$  at 95% CL [11], while Tevatron results exclude the mass range 162–166 GeV at 95% CL [12] and observe an excess of events, recently reported in [13, 14, 15], in the range 120–135 GeV. One of the major goals of LHC is to discover or exclude the Higgs boson. The results from the last year's direct searches at CMS excluded the Higgs boson in the mass range from 127 to 600 GeV at 95% CL, using the data of total integrated luminosity of  $5 \text{ fb}^{-1}$  with a center of mass energy 7 TeV [16]. The ATLAS experiment, using the same amount of data, performed the same search and excluded the Higgs boson at 95% CL in the range of 111.4–116.6, 119.4–122.1, and 129.2 – 541 GeV. Both experiments observed a slight excess of events near 125 GeV [17]. In 2012 the center of mass energy of proton-proton collisions was in-

creased to 8 TeV, and significantly enhanced the sensitivity to the Higgs boson search. The results presented below are a summary of searches performed on combined data, taken in 2011 and 2012. These include searches for the Higgs boson in five decay modes:  $\gamma\gamma$ ,  $ZZ$ ,  $WW$ ,  $\tau^+\tau^-$ , and  $b\bar{b}$ . The combined results indicate an excess of events above the expected background near 125 GeV. The excess has a local significance of 5.0 standard deviations ( $\sigma$ ). This compares well with the expected significance for a SM Higgs boson of  $5.8 \sigma$ . The excess is most prominent in two final states:  $H \rightarrow \gamma\gamma$  with a significance of  $4.1 \sigma$  and  $H \rightarrow ZZ$  with a significance of  $3.2 \sigma$ . The observed new particle must be a boson with spin different from one, as it decays into a pair of photons. As the most sensitivity comes from a di-photon and  $ZZ$  modes, only searches for a Higgs boson in these two decay modes will be described in the details below, while the study of the other three decay modes are described briefly.

### 2. Search for $H \rightarrow \gamma\gamma$

An excellent energy resolution of photons at CMS makes the Higgs boson search to di-photon final state

the most promising channel at low mass [18]. Nevertheless, the search is complicated by a small branching fraction, and a very large background, dominated by two photons from a quantum chromodynamics (QCD) production. Thus, the photon resolution is crucial in this study, so we employ boosted decision tree (BDT) [19, 20] algorithms to improve the photon energy scale using Monte Carlo (MC) simulation and cross check the procedure with  $Z \rightarrow ee$  and  $Z \rightarrow \mu\mu\gamma$  data events. As the resolution of photon energy depends on pseudorapidity, we split events into four non-overlapping event classes based on the photon pseudorapidity and the shape of the energy deposition of the photon candidates. Further improvement is achieved by categorizing events using a multivariate analysis technique (MVA) [21]. We use event-by-event mass resolution, photon identification discriminant, di-photon kinematic variables, and vertex identification probability in the BDT that results in a significant improvement in sensitivity. The further sensitivity is obtained by separately optimizing the vector boson fusion (VBF) production process. Overall, compared to previous studies [21, 22, 23, 24, 25], we achieve an improvement on the combined exclusion sensitivity of approximately 10% in cross section by utilizing BDT, MVA, and the above mentioned categorization.

We illustrate in Fig. 1 an invariant mass of di-photon candidates weighed on the event-by-event basis by  $S/(S+B)$  [26] of the event's category. Here,  $S$  and  $B$  are the number of signal and background events obtained from simultaneous signal plus background best-fit on  $\gamma\gamma$  invariant mass distribution to all categories. In the same figure we also show the unweighted distribution. In both cases we see an obvious excess in data with respect to smoothly falling background from multi-jet events. Here, the background shape is estimated by a polynomial fit to the di-photon mass spectrum in sidebands of the diphoton mass distribution in each of the categories. The background-only hypothesis indicate the expected 95% CL upper limits on the signal strength  $\sigma/\sigma_{SM}$  to be 0.75 near 125 GeV. The results are crosschecked with an independent sideband background model featuring the same observed limit.

The local  $p$ -value is defined as the probability of a background to fluctuate into the observed excess. This quantity, as a function of the mass of the Higgs boson for 7 and 8 TeV data and their combination, is shown in Fig. 2. The minimum local  $p$ -value in data occurs at  $m_H \sim 125$  GeV and has a significance of  $4.1 \sigma$ , while the expected significance is  $2.8 \sigma$ . A production cross section times the relevant branching fractions, relative to the SM expectation is referred to as the signal




Figure 1: The di-photon invariant mass distribution with each event weighted by the  $S/(S+B)$  value of its category. The lines represent the fitted background and signal, and the inset shows the central part of the unweighted invariant mass distribution.

strength and denoted as  $\sigma/\sigma_{SM}$ . The signal strength of best-fit obtained from a SM Higgs boson with mass 125 GeV hypothesis is  $\sigma/\sigma_{SM} = 1.6 \pm 0.4$ .

### 3. Search for $H \rightarrow ZZ$

In the  $H \rightarrow ZZ$  decay mode, the signal has four well-isolated leptons in the final state - making this channel to stand out by having a very distinct signature, that allows a suppression of the most background processes significantly. The later consist mainly of irreducible SM  $ZZ$  production and instrumental  $Z+b\bar{b}$ ,  $t\bar{t}$  production, where a final state contains two true leptons and two  $b$  jets that produce lepton candidates from  $B$ -meson decays. Additionally, the small background comes from  $Z$ +jet and  $WZ$ +jet production, where jets are misidentified as leptons. However, background processes do not affect the sensitivity to the Higgs boson signal as backgrounds result in a smooth four-lepton candidates invariant mass distribution, whereas the Higgs boson signal would result in a narrow peak. Thus, to increase the sensitivity to the Higgs boson production, we focus on optimizing the signal selection efficiency. This is achieved by lowering the thresholds on the transverse momentum  $p_T$  of the lepton candidates identified by very efficient MVA algorithms. We also treat analyses of  $4e$ ,  $4\mu$ , and  $2e2\mu$  final states separately, as these have different background




Figure 2: The local  $p$ -value as a function of  $m_H$  in the  $\gamma\gamma$  decay mode for the combined 7 and 8 TeV data sets. The additional lines show the values for the two data sets taken individually. The dashed line shows the expected local  $p$ -value for the combined data sets, should a SM Higgs boson exist with mass  $m_H$ .

rates and invariant mass resolutions. To further improve the sensitivity to the Higgs boson signal, an algorithm to recover the final state radiation photons is designed and used in this analysis.

The four lepton invariant mass distribution is shown in Fig. 3, where an excess is seen at 125 GeV.

A kinematic discriminant ( $K_D$ ), exploiting the decay kinematics expected for the signal events, is used to provide a significant discriminating power between signal and background. The four lepton system is fully described by five angles and the di-lepton invariant masses (one for each Z boson) [27, 28, 29], thus a probability ratio of the signal and background hypothesis is constructed,  $K_D = \mathcal{P}_{sig}/(\mathcal{P}_{sig} + \mathcal{P}_{bkg})$ , as described in Ref. [30]. We perform six simultaneous two-dimensional (in  $m_{4\ell}$  and  $K_D$  space) maximum-likelihood fits (two data sets and three decay modes) for each value of  $m_H$ .

We show local probability as a function of the mass of the Higgs boson for 7 and 8 TeV data and their combination in Fig. 4. The minimum local  $p$ -value in data occurs at  $m_H = 125.6$  GeV and has a significance of  $3.2\sigma$ , while the expected significance is  $3.8\sigma$ . The best-fit signal strength for a SM Higgs boson mass hypothesis at the same mass point is  $\sigma/\sigma_{SM} = 0.7^{+0.4}_{-0.3}$ .

#### 4. Search for Higgs boson in WW, $\tau\tau$ and $b\bar{b}$ channels

These three channels have less sensitivity to Higgs boson production with the current data samples, and




Figure 3: Distribution of the four-lepton invariant mass for the  $ZZ \rightarrow 4l$  analysis. The points represent data, the filled histograms represent the background, and the open histogram shows the signal expectation for a Higgs boson of mass  $m_H = 125$  GeV, added to the background expectation. The inset shows the  $m_{4\ell}$  distribution after selection of events with  $K_D > 0.5$ , as described in the text.

therefore, we only provide a short summary of searches in these channels.

In  $H \rightarrow WW$  channel, the 95% CL expected and observed limits for the combination of the 7 and 8 TeV analyses are shown in Fig. 5. A broad excess is observed across the mass range. Given the modest mass resolution in this channel, this excess is consistent with a SM Higgs boson with a mass around 125 GeV. This is illustrated by the dotted curve in Fig. 5 showing the median expected limit under the hypothesis of the presence of a SM Higgs boson with mass 125 GeV. The expected significance for a SM Higgs boson with mass 125 GeV is  $2.4\sigma$  and the observed significance is  $1.6\sigma$ .

In  $H \rightarrow \tau\tau$  decay mode, the expected and observed 95% CL limits on the signal strength for the combination of all categories are shown in Fig. 6. The expected and observed limits are 1.3 and 1.1 times the SM Higgs boson cross section at mass 125 GeV, respectively. The expected significance for a SM Higgs boson with mass 125 GeV is  $1.4\sigma$ , and the observed value is zero  $\sigma$ .

In  $H \rightarrow b\bar{b}$  channel, combined results for expected and observed 95% CL limits obtained from the 7 and 8 TeV data sets are displayed in Fig. 7. The expected and observed limits are 1.6 and 2.1 times the SM Higgs boson cross section at mass 125 GeV. The expected significance for a SM Higgs boson of mass 125 GeV cor-




Figure 4: The observed local  $p$ -value for the  $ZZ$  decay mode as a function of the SM Higgs boson mass. The dashed line shows the expected local  $p$ -values for a SM Higgs boson with a mass  $m_H$ .

responds to  $1.9\sigma$ , while the observed one corresponds to  $0.7\sigma$ .

## 5. Combined results

The results from each individual channel analysed are combined using the modified frequentist criterion  $CL_s$  [31, 32]. In combination, we assume the relative branching fractions predicted by the SM and take into account all statistical and systematic uncertainties, as well as the theoretical uncertainties. The  $CL_s$  is shown in Fig. 8 as a function of the Higgs boson mass hypothesis. The observed values are shown by the solid points. The dashed line indicates the median of the expected results for the background-only hypothesis and shaded regions reflect the ranges in which the  $CL_s$  values lie in 68% and 95% CL with the background-only hypothesis. The probabilities of background-only hypothesis to fluctuate into observation for at the 68% (95%) CL are 16% (2.5%) each. The thick horizontal lines indicate  $CL_s$  values of 0.05, 0.01, and 0.001. The mass regions where the observed  $CL_s$  values are below these lines are excluded with the corresponding  $(1 - CL_s)$  confidence levels, resulting in new exclusion range of Higgs boson with  $110 < m_H < 121.5$  GeV at 95% CL. A significant excess is observed in the mass range  $121.5 < m_H < 128$  GeV and the SM Higgs boson cannot be excluded at 95% CL.

In Fig. 9 we show a scan of the local  $p$ -value for each decay modes and combined  $p$ -value. The overall combination has a significance of  $5.0\sigma$  for  $m_H = 125.5$  GeV. The largest contribution comes from  $\gamma\gamma$  and  $ZZ$




Figure 5: The 95% CL limit on  $\sigma/\sigma_{SM}$  for a Higgs boson decaying, via a  $W$  boson pair, to two leptons and two neutrinos, for the combined 7 and 8 TeV data sets. The symbol  $\sigma/\sigma_{SM}$  denotes the production cross section times the relevant branching fractions, relative to the SM expectation. The background-only expectations are represented by their median (dashed line) and by the 68% and 95% CL bands. The dotted curve shows the median expected limit for a SM Higgs boson with  $m_H = 125$  GeV.

channels, as they both are high mass resolution decay modes. The excess indicates the observation of a new particles with mass near 125 GeV. A new particle must be a boson with spin other than one, as it decays into two photons [33, 34].

The mass of the observed particle is determined using the  $ZZ$  and  $\gamma\gamma$  decay modes, the later one is split in untagged and di-jet tagged categories. The precision of this measurement is dominated by  $ZZ$  channel. A fit on invariant mass is performed. In order to decrease the model dependence the fit allows the signal strength to float. The combined best-fit mass is  $m_X = 125.3 \pm 0.4$  (stat.)  $\pm 0.5$  (syst.) GeV. In Fig. 10 we display the two-dimensional 68% CL regions of the signal strength  $\sigma/\sigma_{SM}$  and  $m_X$ . The results from all three channels are compatible within uncertainties.

The scan of signal strength for combined results as a function of the SM Higgs boson mass is provided in Fig. 11. The band corresponds to the  $\pm 1\sigma$  uncertainty. The results indicate that the observed particle is compatible with the SM Higgs boson with mass of 125 GeV within  $\pm 1\sigma$ . The results illustrated in Fig. 12 indicate that the results in all analyzed channels are compatible




Figure 6: The 95% CL limit on the signal strength  $\sigma/\sigma_{\text{SM}}$  for a Higgs boson decaying to  $\tau\tau$ , for the combined 7 and 8 TeV data sets. The symbol  $\sigma/\sigma_{\text{SM}}$  denotes the production cross section times the relevant branching fractions, relative to the SM expectation. The background-only expectations are represented by their median (dashed line) and by the 68% and 95% CL bands. The dotted curve shows the median expected limit for a SM Higgs boson with  $m_H = 125$  GeV.

with each other.

## 6. Conclusions

The results of the SM Higgs boson search in the CMS experiment are presented using the proton proton collisions data at 7 TeV (5  $\text{fb}^{-1}$ ) and 8 TeV (5  $\text{fb}^{-1}$ ). Five decay modes of Higgs boson are analysed -  $\gamma\gamma$ ,  $ZZ$ ,  $WW$ ,  $\tau^+\tau^-$ , and  $b\bar{b}$ . We observe a new particle with the invariant mass of about 125 GeV with a local significance of  $5\sigma$ . This agrees well with the SM Higgs boson expected significance of  $5.8\sigma$ . The excess is the most prominent in  $\gamma\gamma$  and  $ZZ$  decay modes, as these two have the best mass measurement resolution, compared to other channels. A fit to these two channels results in a mass measurement of a new particle to be  $125.3 \pm 0.4$  (stat.)  $\pm 0.5$  (syst.) GeV. Two photon decay indicated that observed particles is a boson with spin different from one. The presented results show that new particles is fully consistent within uncertainties with the SM Higgs boson particle, but the measurements are not very precise yet and there is a chance for this particle not to be the SM Higgs particle.




Figure 7: The 95% CL limit on the signal strength  $\sigma/\sigma_{\text{SM}}$  for a Higgs boson decaying to two b quarks, for the combined 7 and 8 TeV data sets. The symbol  $\sigma/\sigma_{\text{SM}}$  denotes the production cross section times the relevant branching fractions, relative to the SM expectation. The background-only expectations are represented by their median (dashed line) and by the 68% and 95% CL bands. The dotted curve shows the median expected limit for a SM Higgs boson with  $m_H = 125$  GeV.




Figure 8: The  $\text{CL}_s$  values for the SM Higgs boson hypothesis as a function of the Higgs boson mass in the range 110-145 GeV. The background-only expectations are represented by their median (dashed line) and by the 68% and 95% CL bands.




Figure 9: The observed local  $p$ -value for the five decay modes and the overall combination as a function of the SM Higgs boson mass. The dashed line shows the expected local  $p$ -values for a SM Higgs boson with a mass  $m_H$ .




Figure 11: The observed best-fit signal strength  $\sigma/\sigma_{SM}$  as a function of the SM Higgs boson mass in the range 110–145 GeV for the combined 7 and 8 TeV data sets. The symbol  $\sigma/\sigma_{SM}$  denotes the production cross section times the relevant branching fractions, relative to the SM expectation. The band corresponds to the  $\pm 1$  standard deviation uncertainty in  $\sigma/\sigma_{SM}$ .




Figure 10: The 68% CL contours for the signal strength  $\sigma/\sigma_{SM}$  versus the boson mass  $m_x$  for the untagged  $\gamma\gamma$ ,  $\gamma\gamma$  with VBF-like dijet,  $4\ell$ , and their combination. The symbol  $\sigma/\sigma_{SM}$  denotes the production cross section times the relevant branching fractions, relative to the SM expectation. In this combination, the relative signal strengths for the three decay modes are constrained by the expectations for the SM Higgs boson.




Figure 12: Values of  $\sigma/\sigma_{SM}$  for the combination (solid vertical line) and for individual decay modes (points). The vertical band shows the overall  $\sigma/\sigma_{SM}$  value  $0.87 \pm 0.23$ . The symbol  $\sigma/\sigma_{SM}$  denotes the production cross section times the relevant branching fractions, relative to the SM expectation. The horizontal bars indicate the  $\pm 1$  standard deviation uncertainties in the  $\sigma/\sigma_{SM}$  values for individual modes; they include both statistical and systematic uncertainties.

## References

- [1] Englert, F. and Brout, R., "Broken symmetry and the mass of gauge vector mesons", *Phys. Rev. Lett.* 13 (1964) 321. doi:10.1103/PhysRevLett.13.321.
- [2] P. W. Higgs, "Broken symmetries, massless particles and gauge fields", *Phys. Lett.* 12 (1964) 132. doi:10.1016/0031-9163(64)91136-9.
- [3] P. W. Higgs, "Broken symmetries and the masses of gauge bosons", *Phys. Rev. Lett.* 13 (1964) 508. doi:10.1103/PhysRevLett.13.508.
- [4] G. S. Guralnik, C. R. Hagen, T. W. B. Kibble, "Global conservation laws and massless particles", *Phys. Rev. Lett.* 13 (1964) 585. doi:10.1103/PhysRevLett.13.585.
- [5] P. W. Higgs, "Spontaneous symmetry breakdown without massless bosons", *Phys. Rev.* 145 (1966) 1156. doi:10.1103/PhysRev.145.1156.
- [6] T. W. B. Kibble, "Symmetry breaking in non-Abelian gauge theories", *Phys. Rev.* 155 (1967) 1554. doi:10.1103/PhysRev.155.1554.
- [7] S. L. Glashow, "Partial-symmetries of weak interactions", *Nucl. Phys.* 22 (1961) 579. doi:10.1016/0029-5582(61)90469-2.
- [8] S. Weinberg, "A Model of Leptons", *Phys. Rev. Lett.* 19 (1967) 1264. doi:10.1103/PhysRevLett.19.1264.
- [9] A. Salam, "Weak and electromagnetic interactions", in: N. Svartholm (Ed.), *Elementary particle physics: relativistic groups and analyticity*, Almqvist & Wiskell, 1968, p. 367, proceedings of the eighth Nobel symposium.
- [10] ALEPH, CDF, D0, DELPHI, L3, OPAL, SLD Collaborations, the LEP Electroweak Working Group, the Tevatron Electroweak Working Group, and the SLD Electroweak and Heavy Flavour Groups, "Precision Electroweak Measurements and Constraints on the Standard Model", CERN PH-EP-2010-095, "At this time, the most up-to-date Higgs boson mass constraints come from <http://lepewwg.web.cern.ch/LEPEWWG/plots/winter2012/>" (2010). arXiv:1012.2367. URL <http://cdsweb.cern.ch/record/1313716>
- [11] ALEPH, DELPHI, L3, OPAL Collaborations, and LEP Working Group for Higgs Boson Searches, "Search for the standard model Higgs boson at LEP", *Phys. Lett. B* 565 (2003) 61. arXiv:hep-ex/0306033, doi:10.1016/S0370-2693(03)00614-2.
- [12] CDF and D0 Collaborations, "Combination of Tevatron Searches for the Standard Model Higgs Boson in the  $W^+W^-$  Decay Mode", *Phys. Rev. Lett.* 104 (2010) 061802. doi:10.1103/PhysRevLett.104.061802.
- [13] CDF Collaboration, "Combined search for the standard model Higgs boson decaying to a  $b\bar{b}$  pair using the full CDF data set", submitted to *Phys. Rev. Lett.* (2012). arXiv:1207.1707.
- [14] CDF and D0 Collaborations, "Evidence for a particle produced in association with weak bosons and decaying to a bottom-antibottom quark pair in Higgs boson search at the Tevatron", submitted to *Phys. Rev. Lett.* (2012). arXiv:1207.6436.
- [15] D0 Collaboration, "Combined search for the standard model Higgs boson decaying to  $b\bar{b}$  using the D0 Run II data set", submitted to *Phys. Rev. Lett.* (2012). arXiv:1207.6631.
- [16] Chatrchyan, Serguei and others, "Combined results of searches for the standard model Higgs boson in  $pp$  collisions at  $\sqrt{s} = 7$  TeV", *Phys. Lett. B* 710 (2012) 26. arXiv:1202.1488, doi:10.1016/j.physletb.2012.02.064.
- [17] G. Aad, et al., "Combined search for the Standard Model Higgs boson in  $pp$  collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector", *Phys. Rev. D* 86 (2012) 032003. arXiv:1207.0319, doi:10.1103/PhysRevD.86.032003.
- [18] C. J. Seez, T. S. Virdee, L. Di Lella, R. H. Kleiss, Z. Kunszt, "Photon decay modes of the intermediate mass Higgs".
- [19] H.-J. Yang, B. P. Roe, J. Zhu, "Studies of boosted decision trees for MiniBooNE particle identification", *Nucl. Instrum. Meth. A* 555 (2005) 370. doi:10.1016/j.nima.2005.09.022.
- [20] H. Voss, A. Höcker, J. Stelzer, F. Tegenfeldt, "TMVA: Toolkit for Multivariate Data Analysis with ROOT", in: XI Int. Workshop on Advanced Computing and Analysis Techniques in Physics Research, 2007, PoS ACAT:040. arXiv:physics/0703039. URL <http://pos.sissa.it/archive/conferences/050/040/ACAT040.pdf>
- [21] S. Chatrchyan, et al., "Search for the standard model Higgs boson decaying into two photons in  $\gamma\gamma$  collisions at  $\sqrt{s} = 7$  TeV", *Phys. Lett. B* 710 (2012) 403. arXiv:1202.1487, doi:10.1016/j.physletb.2012.03.003.
- [22] S. Chatrchyan, et al., "Search for the standard model Higgs boson in the decay channel  $H \rightarrow ZZ \rightarrow 4\ell$  in  $pp$  collisions at  $\sqrt{s} = 7$ ", *Phys. Rev. Lett.* 108 (2012) 111804. arXiv:1202.1997, doi:10.1103/PhysRevLett.108.111804.
- [23] S. Chatrchyan, et al., "Search for the standard model Higgs boson decaying to  $W^+W_-$  in the fully leptonic final state in  $pp$  collisions at  $\sqrt{s} = 7$ ", *Phys. Lett. B* 710 (2012) 91. arXiv:1202.1489, doi:10.1016/j.physletb.2012.02.076.
- [24] S. Chatrchyan, et al., "Search for neutral Higgs bosons decaying to tau pairs in  $pp$  collisions at  $\sqrt{s} = 7$ ", *Phys. Lett. B* 713 (2012) 68. arXiv:1202.4083, doi:10.1016/j.physletb.2012.05.028.
- [25] S. Chatrchyan, et al., "Search for the standard model Higgs boson decaying to bottom quarks in  $pp$  collisions at  $\sqrt{s} = 7$  TeV", *Phys. Lett. B* 710 (2012) 284. arXiv:1202.4195, doi:10.1016/j.physletb.2012.02.085.
- [26] R. J. Barlow, "Event classification using weighting methods", *J. Comp. Phys.* 72 (1987) 202. doi:10.1016/0021-9991(87)90078-7.
- [27] N. Cabibbo, A. Maksymowicz, "Angular Correlations in  $K_{e4}$  Decays and Determination of Low-Energy  $\pi-\pi$  Phase Shifts", *Phys. Rev. B* 137 (1965) 438, also Erratum, 10.1103/PhysRev.168.1926. doi:10.1103/PhysRev.137.B438.
- [28] Y. Gao, A. V. Gritsan, Z. Guo, K. Melnikov, M. Schulze, et al., "Spin determination of single-produced resonances at hadron colliders", *Phys. Rev. D* 81 (2010) 075022. arXiv:1001.3396, doi:10.1103/PhysRevD.81.075022.
- [29] A. De Rujula, J. Lykken, M. Pierini, C. Rogan, M. Spiropulu, "Higgs look-alikes at the LHC", *Phys. Rev. D* 82 (2010) 013003. arXiv:1001.5300, doi:10.1103/PhysRevD.82.013003.
- [30] S. Chatrchyan, et al., "Search for a Higgs boson in the decay channel  $H \rightarrow ZZ^{(*)} \rightarrow q\bar{q}\ell^-\ell^+$  in  $pp$  collisions at  $\sqrt{s} = 7$  TeV", *JHEP* 04 (2012) 036. arXiv:1202.1416, doi:10.1007/JHEP04(2012)036.
- [31] T. Junk, "Confidence level computation for combining searches with small statistics", *Nucl. Instrum. Meth. A* 434 (1999) 435. doi:10.1016/S0168-9002(99)00498-2.
- [32] A. L. Read, "Presentation of search results: the CLs technique", *J. Phys. G* 28 (2002) 2693. doi:10.1088/0954-3899/28/10/313.
- [33] L. D. Landau, "On the angular momentum of a two-photon system", *Dokl. Akad. Nauk* 60 (1948) 207.
- [34] C. N. Yang, "Selection Rules for the Dematerialization of a Particle into Two Photons", *Phys. Rev.* 77 (1950) 242. doi:10.1103/PhysRev.77.242.
- [35] S. Chatrchyan, et al., "The CMS experiment at the CERN LHC", *JINST* 3 (2008) S08004. doi:10.1088/1748-0221/3/08/S08004.
- [36]