

BSM with FEYNRULES

Towards NLO: status and plans

Benjamin Fuks (IPHC Strasbourg / Université de Strasbourg)

In collaboration with N.D. Christensen and C. Duhr.

ERC Miniworkshop @ CERN
January 27, 2012

Outline

- 1 Status of FEYNRULES.
- 2 FEYNRULES at NLO - Status and plans.
- 3 Summary.


Implementation of BSM theories in Monte Carlo tools.

- A model consists in:
 - * **Particles**,
 - * **Parameters**,
 - * **Interactions** (\equiv Feynman rules).
- The Feynman rules **have to be derived (from a Lagrangian)**.
 - ▶ **Translated in a programming language.**
⇒ **Tedious, time-consuming, error prone.**
 - ▶ **Iterations** for each model.
 - ▶ **Iterations** for each MC tool.
 - ▶ Beware: **Lorentz and color structures**.
 - ▶ Beware: **validation**.

Redundancies of the work.

A framework for LHC analyzes.

[Christensen, de Aquino, Degrande, Duhr, BenjF, Herquet, Maltoni, Schumann (EPJC '11)]



FEYNRULES in a nutshell.

[Christensen, Duhr (CPC '09); Christensen, Duhr, BenjF (in prep)]

- **A framework for LHC analyzes based on FEYNRULES to:**
 - * **Develop new models.**
 - * **Implement (and validate)** new models in Monte Carlo tools.
 - * Facilitate **phenomenological** investigations of the models.
 - * **Test** the models against data.
- **Main features**
 - * FEYNRULES is a MATHEMATICA package.
 - * FEYNRULES derives **Feynman rules from a Lagrangian.**
 - * **Requirements:** locality, Lorentz and gauge invariance.
 - * **Supported fields:** scalar, fermion, vector, tensor, ghost, superfield.
 - * **Interfaces:** export the Feynman rules to Monte Carlo generators. CALCHEP, FEYNARTS, MADGRAPH, SHERPA, WHIZARD
 - * **Universal FEYNRULES output:** MADGRAPH5 and GOSAM.

FEYNRULES-1.6 - status.

● Current public version: 1.6.0.

- * **To be download on <http://feynrules.irmp.ucl.ac.be/>.**
- * Contains the **superspace module**. [Duhr, BenjF (CPC '11)]
- * Contains the **UFO interface** \Rightarrow MADGRAPH5, GOSAM.
[Degrande, Duhr, BenjF, Grellscheid, Mattelaer, Reiter (2011)]
- * Supports **color sextets**.
- * Contains the new **FEYNARTS interface**.
 \Rightarrow **Generic Lorenz structures allowed**.
- * Interfaced to **WHIZARD**. [Christensen, Duhr, BenjF, Reuter, Speckner (2010)]
- * Other interfaces: **CALCHEP/COMPHEP**, **MADGRAPH4**, **SHERPA**.
- * **Manual currently being updated** [Christensen, Duhr, BenjF (in prep)].

● Current online model database.

- * **<http://feynrules.irmp.ucl.ac.be/wiki/ModelDatabaseMainPage/>** .
- * Standard Model and simple extensions (10).
- * Supersymmetric models (4).
- * Extra-dimensional models (4).
- * Strongly coupled and effective field theories (4).

Outline.

- 1 Status of FEYNRULES.
- 2 FEYNRULES at NLO - Status and plans.
- 3 Summary.

NLO calculations with MADGRAPH - AMC@NLO.

● Real emission.

- * Must include the appropriate **subtraction terms**.
⇒ MADFKS [Frederix, Frixione, Maltoni, Stelzer (JHEP '09)].
- * The tree-level Feynman rules are the **only required components**.

😊 No particular problem for BSM ⇒ problem solved. 😊
(Use of FEYNRULES, its interfaces to MC tools)

● One-loop virtual amplitudes.



- * Several algorithms have been proposed in the last few years.
⇒ MADLOOP [Hirschi, Frederix, Frixione, Garzelli, Maltoni, Pittau (JHEP '11)].
⇒ based on OPP reduction [Ossola, Papadopolous, Pittau (NPB '07)].
- * Requirements:
 - ◇ **Tree-level Feynman rules**.
 - ◇ **UV renormalization counterterms**.
 - ◇ **Rational R_2 terms**.

☹ The two latter must be included by hand. ☹

NLO calculations in the context of FEYNRULES.

- Counterterms and R_2 terms.

- *  Non-automatic steps.
- *  Can be derived from the tree-level Lagrangian.

 All the information is already there at the FEYNRULES-level. 

- Automatic renormalization in the $\overline{\text{MS}}$ -scheme with FEYNRULES

- ① Automated extraction of the renormalized Lagrangian ✓.
- ② Modification of the FEYNARTS interface to include counterterms ✓.
- ③ Calculation of the renormalization constants with FORMCALC.
 - ▶ Self-energies: 80% done.
 - ▶ Vertices ✗
- ④ Re-injection in FEYNRULES ✗.

- Automatic R_2 terms ✗.

- The UFO at NLO: basically there [UFO people + Hirschi]

Automatic renormalization with FEYNRULES.

- **Expansion of the renormalization constants (works with full Lagrangians).**
 - * The **type of the interactions** in the loops can be specified.
 - * The **loop-level** can be specified.

ExtractCounterterms [l [s, f], {aS, 1}]

$$\blacktriangleright l_{sf} \rightarrow l_{sf} + \frac{\alpha_s}{4\pi} \left[(\delta Z_{ll}^{L(1)})_{ff'} (P_L)_{ss'} + (\delta Z_{ll}^{R(1)})_{ff'} (P_R)_{ss'} \right] l_{s'f'}$$

ExtractCounterterms [y_d, {{aS, 2}, {aEW, 1}}]

$$\blacktriangleright y_d \rightarrow y_d + \frac{\alpha_s}{2\pi} \delta y_d^{(1,0)} + \frac{\alpha}{2\pi} \delta y_d^{(0,1)} + \frac{\alpha_s^2}{4\pi^2} \delta y_d^{(2,0)} + \frac{\alpha_s \alpha}{4\pi^2} \delta y_d^{(1,1)} + \frac{\alpha_s^2 \alpha}{8\pi^3} \delta y_d^{(2,1)}$$

- **Treatment of the internal parameters.**
 - * Automatic computation of the **relations** among renormalization constants.
 - * **Only** the ren. cnsts of the **external parameters** will have to be computed.

g_s and α_s at first order in QCD.

$$g_s = 2\sqrt{\pi\alpha_s} \quad \Rightarrow \quad \delta g_s^{(1)} = \frac{\sqrt{\alpha_s}}{2\sqrt{\pi}} \delta \alpha_s^{(1)}$$

Outline.

- 1 Status of FEYNRULES.
- 2 FEYNRULES at NLO - Status and plans.
- 3 Summary.

Conclusions.

- FEYNRULES provides a platform to:
 - * **Develop** new models.
 - * Investigate their **phenomenology**.
- FEYNRULES and leading order tools.
 - * **Many interfaces** exist.
 - * **Any model** (renormalizable or not) can be exported to at least one MC.
 - * **MC event generation at LO**: the problem is solved (up to spin-3/2 fields).
- NLO challenges.
 - * **Achievement of the UFO @ NLO format.**
 - ▶ Easter '12.
 - * **Automatic renormalization.**
 - ▶ Summer '12.
 - * **Automatic R_2 terms.**
 - ▶ Summer '12 ?
- Full merging to the NLO tools.