

Detailed Measurements of Bottomonium Suppression in PbPb Collisions at 2.76 TeV with CMS



Guillermo Breto Rangel

(University of California, Davis)

for the CMS Collaboration



Quark Matter conference, Washington DC

14th August, 2012

Quarkonia

1. Quarkonia are very “unusual” hadrons

heavy quark ($Q\bar{Q}$) bound states **stable** under strong decay

- heavy: $m_c \simeq 1.2 - 1.4$ GeV, $m_b \simeq 4.6 - 4.9$ GeV
- stable: $M_{c\bar{c}} - 2M_D \ll 0$ and $M_{b\bar{b}} - 2M_B \ll 0$

What is “usual”?

- light quark ($q\bar{q}$) constituents
- loosely bound, $M_\rho - 2M_\pi \gg 0$, $M_\Phi - 2M_K \simeq 0$
- hadronic size $\sim 1/\lambda_{\text{QCD}} \simeq 1$ fm, independent of mass



(At $T = 0$ Cornell potential gives full spectroscopy)

State	J/ψ	ψ'	Υ	Υ'	Υ''
mass [GeV]	3.10	3.68	9.46	10.02	10.36
ΔE [GeV]	0.64	0.05	1.10	0.54	0.20
radius [fm]	0.25	0.45	0.14	0.28	0.39

Relativistic Heavy Ion Physics

By F Becattini, P Braun-Munzinger, Rainer Fries, C Gale, J. Schaffner-Bielich

Sequential Melting



✓ High temperature QGP: weaker color binding (Debye screening) ([Phys. Lett. B178 \(1986\) 416](#))

- Real $V(T)$


✓ Gluons collide with $Q\bar{Q}$ bound states:
- shorter lifetime \Rightarrow larger spectral widths

(Landau Damping) ([JHEP 0703:054,2007](#))

- Im $V(T)$

✓ Dissociation when $\text{Re } V(T) \sim \text{Im } V(T)$


✓ Quarkonia:
help quantify medium properties (TEMPERATURE)



[Margotta et al. Phys. Rev. D 84, 069902\(E\) \(2011\)](#)

Bottomonium in Heavy Ions collisions

- ✓ Mass of the b-quark is large
- ✓ No B hadron feed down to Υ
- ✓ nPDF effects smaller
- ✓ The relative yields analysis of the excited states / ground state
 - cancels cold nuclear matter effects
 - nPDFs (shadowing, etc)
 - initial parton energy loss
 - final state nuclear absorption (if negligible at LHC energies)
 - carries only effects related to final (hot) medium
- ✓ Regeneration is smaller




CMS

The Compact **MUON**
Solenoid detector

3.8T Superconducting Solenoid

Hermetic Hadronic
Calorimeter (HCAL)
[Scintillators & Brass]

Lead tungstate E/M
Calorimeter (ECAL)



All Silicon Tracker (Pixels and MicroStrips)

Redundant Muon System
(RPCs, Drift Tubes,
Cathode Strip Chambers)

Υ candidate in PbPb at $\sqrt{s_{NN}} = 2.76$ TeV

CMS Experiment at the LHC, CERN

Data recorded: 2010-Nov-12 03:55:57.236106 GMT(04:55:57 CEST)

Run / Event: 150887 / 1792020

**Hardware L1 Trigger +
Software HLT (High Level Trigger)
Dimuon trigger rate ~ 30 Hertz**

Trigger must be:

Fast

Flexible

Efficient (Single Muon Eff. ~ 95 %)

Redundant


$\mu^+\mu^-$ pair mass: **$9.46 \text{ GeV}/c^2$**

p_T : **$0.06 \text{ GeV}/c$**

$\mu^+ : p_T = 4.74 \text{ GeV}/c$


$\mu^- : p_T = 4.70 \text{ GeV}/c$

$\Upsilon(2S+3S)$ Suppression in 2010



[PRL 107\(2011\)052302](https://arxiv.org/abs/1011.5230)

$$\Upsilon(2S + 3S)/\Upsilon(1S)|_{pp} = 0.78^{+0.16}_{-0.14} \pm 0.02$$




$$\Upsilon(2S + 3S)/\Upsilon(1S)|_{PbPb} = 0.24^{+0.13}_{-0.12} \pm 0.02$$

Measured $\Upsilon(2S+3S)$ production relative to $\Upsilon(1S)$

$$\frac{\Upsilon(2S + 3S)/\Upsilon(1S)|_{PbPb}}{\Upsilon(2S + 3S)/\Upsilon(1S)|_{pp}} = 0.31^{+0.19}_{-0.15} \pm 0.03$$

Indication of 2S+3S relative suppression
significance = 2.4 σ

Muon pairs in PbPb at $\sqrt{s_{NN}} = 2.76$ TeV



✓ Kinematic cuts applied to reduce background level

– Single muon:

$$p_T^\mu > 4 \text{ GeV}/c, |\eta^\mu| < 2.4$$


✓ pp - PbPb

Same cuts, same reco. algorithm



Systematics cancel

Fitting Model in 2011

- ✓ Unbinned maximum likelihood fit
 - Six signal and four background parameters float in the fit
- ✓ Signal
 - Three resonances modeled by crystal-ball function: Gaussian resolution and FSR power-law low mass tail
 - Mass differences of three resonances are fixed to PDG value
 - Float FSR & resolution (fixed in 2010 data analysis)
- ✓ Background
 - Exponential x error-function (Erf describes kinematic turn-on)
- ✓ Variations of the models checked as systematics.



Simultaneous Fit 2011



$$\Upsilon(2S+3S)/\Upsilon(1S) |_{\text{PbPb}} / \Upsilon(2S+3S)/\Upsilon(1S) |_{\text{pp}} = 0.15 \pm 0.05 \pm 0.03$$

Observation of 2S+3S relative suppression
(significance $> 5 \sigma$)

$\Upsilon(nS)/\Upsilon(1S)$ Double ratio

Separated $\Upsilon(2S)$ and $\Upsilon(3S)$

(0 - 100) % Centrality Integrated

$$\frac{Y(2S)/Y(1S)|_{\text{PbPb}}}{Y(2S)/Y(1S)|_{\text{pp}}} = 0.21 \pm 0.07 \text{ (stat.)} \pm 0.02 \text{ (syst.)}$$


$$\frac{Y(3S)/Y(1S)|_{\text{PbPb}}}{Y(3S)/Y(1S)|_{\text{pp}}} = 0.06 \pm 0.06 \text{ (stat.)} \pm 0.06 \text{ (syst.)}$$

$$< 0.17 \text{ (95\% C.L.)}$$

$\Upsilon(2S)$ relative ratio to $\Upsilon(1S)$ in PbPb is suppressed compared to same ratio in pp

Systematics Uncertainties:

- ➔ fitting (11%)
- ▶ Final state radiation modeling.
- ▶ Background modeling: like-sign vs track-rotation
- ➔ imperfect acceptance+efficiency cancelation: 1%



$\Upsilon(nS)$ Absolute Suppression

$$R_{AA} = \frac{\mathcal{L}_{pp}}{T_{AA} N_{MB}} \frac{N_{PbPb}(\Upsilon(1S))}{N_{pp}(\Upsilon(1S))} \frac{\epsilon_{pp}}{\epsilon_{PbPb(\text{cent})}}$$

First time the nuclear modification factors are measured for excited Υ states

$$\begin{aligned} R_{AA}(\Upsilon(1S)) &= 0.56 \pm 0.08 \text{ (stat.)} \pm 0.07 \text{ (syst.)} \\ R_{AA}(\Upsilon(2S)) &= 0.12 \pm 0.04 \text{ (stat.)} \pm 0.02 \text{ (syst.)} \\ R_{AA}(\Upsilon(3S)) &= 0.03 \pm 0.04 \text{ (stat.)} \pm 0.01 \text{ (syst.)} \\ &< 0.10 \text{ (95\% C.L.)}. \end{aligned}$$

Υ states are suppressed sequentially:


$$R_{AA}^{\Upsilon(3S)} < R_{AA}^{\Upsilon(2S)} < R_{AA}^{\Upsilon(1S)}$$

✓ Note: Inclusive measurement of $\Upsilon(1S)$ vs. direct production.

- $R_{AA}(\Upsilon(1S))$ -inclusive : Feed-down contributions (χ_b , $\Upsilon(2S)$, $\Upsilon(3S)$).

-If feed-down $\sim 50\%$, $R_{AA}(\Upsilon(1S))$ -inclusive is consistent with suppression of excited states only.

$\Upsilon(nS)$ R_{AA} vs Centrality




Suppression observed to increase with centrality of the collisions

$\Upsilon(2S)$

- ✓ Always more suppressed than $\Upsilon(1S)$
- ✓ Still suppressed in the most peripheral bin (50-100%)

Global Uncertainties
 lumi pp & fitting pp:
 14% $\Upsilon(1S)$
 21% $\Upsilon(2S)$

$\Upsilon(3S)$ Upper Limit




The $\Upsilon(3S)$ peak is barely visible in the PbPb data.

Set upper limits at 95% C.L. using the Feldman Cousins technique. ([arXiv:physics/9711021v2](https://arxiv.org/abs/physics/9711021v2))

$$R_{AA}^{\Upsilon(3S)} < 0.1$$

(0 - 100) % Centrality Integrated

Experimental Comparisons



✓ STAR measured R_{AA} of $\Upsilon(1S+2S+3S)$ combined

$$R_{AA}(\Upsilon(1S + 2S + 3S)) = 0.56 \pm 0.21^{+0.08}_{-0.16}$$

([arXiv:1109.3891v1](https://arxiv.org/abs/1109.3891v1))



✓ CMS: separate R_{AA} for $\Upsilon(1S)$ and $\Upsilon(2S)$ can calculate R_{AA} of $\Upsilon(1S+2S+3S)$:

$$R_{AA}(\Upsilon(1S + 2S + 3S)) = R_{AA}(\Upsilon(1S)) \times \frac{1 + \Upsilon(2S + 3S)/\Upsilon(1S)|_{PbPb}}{1 + \Upsilon(2S + 3S)/\Upsilon(1S)|_{pp}}$$

$$R_{AA}(\Upsilon(1S+2S+3S)) \sim 0.32$$

✓ Similar Suppression Pattern

Theoretical Comparisons




CMS data consistent within uncertainties with range of suppression predicted for both Y(1S) and Y(2S).

Strickland and D. Bazow ([PRL 107 \(2011\) 132301](https://arxiv.org/abs/1011.2711))

Emerick, X.Zhao,R.Rapp ([Eur. Phys. J. A48 \(2012\) 72](https://arxiv.org/abs/1108.3801))

Summary

- ✓ First measurements of the excited Υ states in heavy-ions.
- ✓ Υ (1S) suppression consistent with melting of excited states only.
- ✓ Suppression pattern (sequential “melting”) has been established.
- ✓ Set upper limits for the Υ (3S) state for the first time.



Thank You

Special thanks to:

Zhen Hu

Nuno Leonardo

Torsten Dahms

Jake Anderson

Michael Gardner

Vineet Kumar

Camelia Mironov

Dongho Moon

Mihee Jo

Raphael de Cassagnac

Lorenzo Moneta

Dilepton Group


CMS Collaboration

ARC Members

Convenors


QM12 Organizers

Manuel Calderón




Back-up

Charmonia Comparisons




Muon pairs in PbPb and pp at $\sqrt{s_{NN}} = 2.76$ TeV



The Compact Muon Solenoid


CMS Detector

Pixels
Tracker
ECAL
HCAL
Solenoid
Steel Yoke
Muons



Total weight : 14000 tonnes
Overall diameter : 15.0 m
Overall length : 28.7 m
Magnetic field : 3.8 T

Muon reconstruction in CMS



Global muons reconstructed with information from tracker and muon stations



Muons need to overcome the magnetic field and energy loss in the absorber
→ need a minimum momentum of $p \sim 3-5 \text{ GeV}/c$ to reach the muon stations

Further muon ID based on track quality (χ^2 , # hits,...)

Quarkonia Suppression

We have established the suppression pattern pinning down medium properties.


We have measured the Quarkonia sequential suppression



Significance

Could background fluctuation produce a result as extreme as observed in data?


- Generate pseudo-experiments following the *null-hypothesis* (i.e. no suppression)
- Fit pseudo-data samples with nominal fit
- Count fraction of occurrences for which the ratio (taken as test statistic) is same or lower than observed:
 - p-value: 0.9%
 - 2.4σ (1-sided Gaussian test)



[PRL 107 \(2011\) 052302](#)

Acceptance/Efficiency

Acceptance



Efficiency

