



Contribution ID: 359

Type: Oral Presentation

Jet-Tagged Back-Scattering Photons For Quark Gluon Plasma Tomography

Wednesday, August 15, 2012 9:30 AM (20 minutes)

We investigate the correlations of photons produced by back scattering of fast partons in quark gluon plasma with their away-side jets. Back scattering with photon emission, or jet-photon conversion, was originally proposed as a novel source of photons in Phys. Rev. Lett. 90, 132301 (2003).

The unique appeal of this photon source lies in the fact that its photons carry information about both the medium via a $T^2 \log 1/T$ dependence of the yield and about the energy loss of partons before the back scattering occurs. Attempts to identify this source in experiment through inclusive direct photon spectra or direct photon v_2 at intermediate PT at RHIC have been inconclusive so far. We show that the capability to measure jets in coincidence with photons at the upgraded STAR or SPHENIX experiment, or at one of the LHC experiments, offers a unique opportunity to identify back scattering photons at large photon momenta. Jet-triggered back-scattering photons can be distinguished from bremsstrahlung through their strong correlation with the given trigger ET, and from prompt hard photons through the energy loss of their parent parton.

We demonstrate with leading and next-to-leading order calculations that jet-triggered direct photon spectra and nuclear modification factors in nuclear collisions as a function of photon PT show a distinct feature around the trigger ET due to back-scattering photons.

The height and width of this structure are correlated with the medium temperature and parton energy loss spectrum, respectively.

Author: Prof. FRIES, Rainer (Texas A&M University)

Co-authors: SRIVASTAVA, Dinesh (Variable Energy Cyclotron Centre, Kolkata); DE, Somnath (Variable Energy Cyclotron Centre, Kolkata)

Presenter: Prof. FRIES, Rainer (Texas A&M University)

Session Classification: Parallel 3C: Electro-Weak Probes (Chair J. Kapusta)