



Contribution ID: 119

Type: Oral Presentation

## Thermal dileptons in high-energy heavy ion collisions with 3+1D relativistic hydrodynamics

*Wednesday, August 15, 2012 8:30 AM (20 minutes)*

The penetrating nature of dileptons makes them suitable probes to explore the properties of the strongly-interacting medium created in relativistic nuclear collisions. This study investigates thermal dilepton production using MUSIC (a Monotone Upstream-centered Scheme for Ion Collisions): a 3+1D hydrodynamic simulation with or without shear viscosity. We utilize dilepton emission rates that are derived from in-medium hadronic spectral functions, and from pQCD. In addition to the invariant mass and momentum distributions, the elliptic flow of lepton pairs is calculated, and the effects of a finite shear viscosity coefficient are also analyzed. We present results appropriate for measurements by the PHENIX and STAR collaborations, and make predictions for the LHC.

**Primary author:** VUJANOVIC, Gojko (McGill University)

**Co-authors:** SCHENKE, Bjoern (Brookhaven National Laboratory); GALE, Charles (McGill University); YOUNG, Clint (McGill University); RAPP, Ralf (Texas A&M University); JEON, Sangyong (McGill University)

**Presenter:** VUJANOVIC, Gojko (McGill University)

**Session Classification:** Parallel 3C: Electro-Weak Probes (Chair J. Kapusta)

**Track Classification:** Electroweak probes