Quark Matter 2012



Contribution ID: 428

Type: Poster

Balance function studies for non-identified particles in Pb-Pb collisions at √sNN = 2.76 TeV

Thursday 16 August 2012 16:00 (2 hours)

The possible creation of a strongly interacting deconfined phase (Quark-Gluon plasma) in relativistic heavy ion collisions would be measurable in a delayed hadronization time. It was proposed to test this hypothesis via the measurement of correlations between positive and negatively charged pairs pairs as a function of rapidity, the so-called Balance functions, which was done at SPS and RHIC energies. We will present first studies of Balance functions at LHC energies with the ALICE experiment. In Pb–Pb collisions at $\sqrt{sNN} = 2.76$ TeV the width of Balance functions in pseudorapidity and azimuthal angle for non-identified charged particles shows a clear centrality dependence. In addition a comparison to other experiments (NA49, STAR) at lower c.m. energies and different models (Hijing, AMPT, Blast wave) will be presented.

Primary author: ALICE, Collaboration (CERN, Geneva, Switzerland)
Co-author: RODRIGUEZ MANSO, Alis (NIKHEF (NL))
Presenter: RODRIGUEZ MANSO, Alis (NIKHEF (NL))
Session Classification: Poster Session Reception

Track Classification: Correlations and fluctuations