


DI-BOSONS AND ANOMALOUS COUPLINGS AT D0

Heidi Schellman, Northwestern University




Run II Integrated Luminosity

19 April 2002 -30 September 2011



DØ Detector




Lepton ID in the DØ Detector

4



Liquid Argon Calorimeter

5





- Liquid argon active medium and uranium/copper absorber
- Hermetic with coverage for: $|\eta| < 4.2$
- Longitudinal and Transverse segmentation $\Delta\eta \times \Delta\varphi \times X^0 = 0.1 \times 0.1 \times \sim 1$
(0.05×0.05 in third EM layer, near shower maximum)

Lagrangian for neutral (ZZ γ /Z $\gamma\gamma$)

6

$$L_{\gamma ZV} = -ie \left[(h_1^V F^{\mu\nu} + h_3^V \tilde{F}^{\mu\nu}) Z_\mu \frac{(\square + m_V^2)}{M_Z^2} V_\nu + (h_2^V F^{\mu\nu} + h_4^V \tilde{F}_{\mu\nu}) Z^\alpha \frac{(\square + m_V^2)}{M_Z^4} \partial_\alpha \partial_\mu V_\nu \right]$$

CP conserving $\rightarrow h_{3,4}^V$ couplings ($V = \gamma, Z$)



$\Delta \neq 0$
ANOMALOUS
COUPLINGS

SM : $h_{3,4}^V = 0$

SM Deviations : $\Delta h_{3,4}^V = h_{3,4}^V - 0$



Lagrangian for charged (WW γ /WWZ)

7

$$\frac{L_{WWV}}{g_{WWV}} = ig_1^V (W_{\mu\nu}^* W^\mu V^\nu - W_\mu^* V_\nu W^{\mu\nu}) + i\kappa_V W_\mu^* W_\nu V^{\mu\nu} + i \frac{\lambda_V}{M_W^2} W_{\lambda\mu}^* W_\nu^\mu V^{\nu\lambda}$$

$$- g_4^V W_\mu^* W_\nu (\partial^\mu V^\nu + \partial^\nu V^\mu) + g_5^V \epsilon^{\mu\nu\lambda\rho} (W_\mu^* \partial_\lambda W_\nu - \partial_\lambda W_\mu^* W_\nu) V_\rho + i\tilde{\kappa}_V W_\mu^* W_\nu \tilde{V}^{\mu\nu} + i \frac{\tilde{\lambda}_V}{M_W^2} W_{\lambda\mu}^* W_\nu^\mu \tilde{V}^{\nu\lambda}$$

EM gauge inv. ($g_1^Y = 1$), C and P conserving \Rightarrow 5 couplings: $\kappa_V, \lambda_V, g_1^Z$



SM : $g_1^Z = \kappa_V = 1, \lambda_V = 0$


SM Deviations:

$\Delta g_1^Z = g_1^Z - 1, \Delta \kappa_V = \kappa_V - 1, \Delta \lambda_V = \lambda_V - 0$

$W\gamma$ and $Z\gamma$

8

- D0 has excellent photon identification due to fine transverse and longitudinal segmentation in the calorimeter.
- Neural network discriminant yields $> 90\%$ purity in $Z\gamma$





$W\gamma \rightarrow l\nu\gamma$ Production



PRL 107, 241803 (2011)


9






$$\sigma_{W\gamma} \times \text{BR}(W \rightarrow l\nu) = 7.6 \pm 0.4(\text{stat}) \pm 0.6(\text{syst}) \text{ pb}$$

SM@NLO: $\sigma = 7.6 \pm 0.2 \text{ pb}$

(photon $E_T > 15 \text{ GeV}$, $dR_{(l\gamma)} > 0.7$)



Anomalous $W\gamma$ couplings



95% CL limits on TGCs:

$-0.4 < \Delta\kappa_\gamma < 0.4, -0.08 < \lambda_\gamma < 0.07$


$Z\gamma \rightarrow \nu\nu\gamma$

[Phys. Rev. Lett. 102, 201802 \(2009\)](#)




11

Select interactions with large, significant
Missing transverse momentum



$Z\gamma \rightarrow \nu\nu\gamma$ avoids radiation off of the Z!





Limits on the anomalous
couplings of Z's to
photons.

$Z\gamma \rightarrow ll\gamma$ Production



Phys. Rev. D 85, 052001 (2012)

12



$$\sigma_{Z\gamma} \times \text{BR}(Z \rightarrow ll) = 1089 \pm 40_{\text{(stat)}} \pm 65_{\text{(syst)}} \text{ fb}$$

$$\text{SM@NLO : } \sigma = 1096 \pm 34 \text{ fb}$$

$$\sigma_{Z\gamma} \times \text{BR}(Z \rightarrow ll) = 288 \pm 15_{\text{(stat)}} \pm 11_{\text{(syst)}} \text{ fb}$$



$$M_{ll\gamma} > 110 \text{ GeV (FSR removal)}: \quad \text{SM@NLO : } \sigma = 294 \pm 10 \text{ fb}$$

$Z\gamma \rightarrow \mu\mu\gamma$

13

	$\ell\ell\gamma$	7.2 fb^{-1}
	$\nu\nu\gamma$	3.6 fb^{-1}
	Λ	1.5 TeV
$ h_{03}^Z <$		0.026
$ h_{04}^Z <$		0.0013
$ h_{03}^\gamma <$		0.027
$ h_{04}^\gamma <$		0.0014

Fit the p_T distribution to limit anomalous couplings
 Combine with the $\nu\nu\gamma$ channel for improved limits.




Combine with $\nu\nu\gamma$

$WZ \rightarrow l\bar{v}l\bar{l}$ Production

Phys. Rev. D 85, 112005 (2012)

14

Three high p_T (isolated) leptons ($\mu\mu\mu$, eee , $ee\mu$, $\mu\mu e$) +MET



$$\sigma_{WZ} = 4.5^{+0.6}_{-0.7} (\text{stat + syst}) \text{ pb}$$

SM@NLO: $\sigma = 3.21 \pm 0.19 \text{ pb}$
 $(60 < M_{ll} < 120 \text{ GeV})$

ZZ → llll Production



15


Cross section for the ZZ production using fully leptonic final states

- Four high p_T leptons ($\mu\mu\mu\mu$, $eeee$, $ee\mu\mu$)

$ZZ \rightarrow llll$
 10 events
 .37 background

$$\sigma_{ZZ} = 1.26^{+0.47}_{-0.37} (\text{stat + syst}) \text{ pb}$$

SM@NLO: $\sigma = 1.40 \pm 0.10 \text{ pb}$



$ZZ \rightarrow vvll$ Production



Phys. Rev. D 85, 112005 (2012)




16

Challenges in $vvll$ final states:

- MET reconstruction
- WW background

$$p'_T = p_T^{\ell\ell} + 2[p_T^\delta + p_T^{\text{recoil}} + p_T^{\text{trkjets}}]$$

Optimal variable is a modified p_T with negative corrections proportional to resolutions (ie. $p_T^\delta < 0$)



Result:
vvll (8.6 fb^{-1}):
 $\sigma_{ZZ} = 1.64^{+0.46}_{-0.46} (\text{stat+syst}) \text{ pb}$
SM@NLO: $\sigma = 1.40 \pm 0.10 \text{ pb}$

$ZZ \rightarrow vvll$ Production



Phys. Rev. D 85, 112005 (2012)




17

Challenges in $vvll$ final states:

- MET reconstruction
- WW background

$$p'_T = p_T^{\ell\ell} + 2[p_T^\delta + p_T^{\text{recoil}} + p_T^{\text{trkjets}}]$$

Optimal variable is a modified p_T with negative corrections proportional to resolutions (ie. $p_T^\delta < 0$)



Combined:
llll (6.4 fb^{-1})+vvll (8.6 fb^{-1}):


$$\sigma_{ZZ} = 1.44^{+0.35}_{-0.34} (\text{stat+syst}) \text{ pb}$$

SM@NLO: $\sigma = 1.40 \pm 0.10 \text{ pb}$


Tevatron WZ and ZZ cross sections

18

WZ



ZZ




Summary of Bosons at the Tevatron + friends

19

Tevatron $p\bar{p}$ at $\sqrt{s} = 1.96$ TeV + LHC 7 TeV

March 2012



Summary

20




- Diboson cross sections have been measured in many all-leptonic channels
- Future updates
 - ▣ Full statistics ZZ
 - ▣ Full statistics WW
 - ▣ Anomalous couplings using all channels

BACKUP SLIDES

Older D0 limits on anomalous couplings from $W\gamma$, WW , WZ



22



Can be interpreted as measurements of the magnetic dipole and quadrupole moments.

$$\begin{aligned} \mu_W &= (1 + \kappa + \lambda) \frac{e}{2M_W} = 2.02^{+0.08}_{-0.09} \frac{e}{2M_W} \\ q_W &= -(\kappa - \lambda) \frac{e}{M_W^2} = 1.00 \pm 0.09 \frac{e}{M_W^2} \end{aligned}$$

Prospects for the future

23

- D0 and CDF are updating to the full statistical sample.
 - Done for WZ and ZZ channels
- Combining channels and experiments will increase the TGC sensitivity by a factor of 3-5.
- LHC experiments have 10 times the cross section → 10 fb^{-1} of data → factor of 10 in statistics and 3 in sensitivity.


Motivation for Diboson Studies



Probe of the EWSB mechanism

- Test of the SM
- Indirect searches for New Physics

Cross sections, Kinematic distributions, Trilinear Gauge Boson Couplings (TGCs)



Important background to:


- Top
- Higgs
- Beyond the SM

Good understanding is highly valuable

Proving ground for analysis techniques
and statistical treatment used in the

Tevatron Higgs searches




Complementary to Higgs production
(same final states/challenges)



D0 – set charged TGC limits using pt in 4 channels



25



Charged Triple Gauge Couplings

Probed by WW , WZ , and $W\gamma$ production


General Lagrangian has **14** parameters

Assume EM gauge invariance and C and P conservation

\Rightarrow 5 TGC parameters:

$$g^1_z, \kappa_\gamma, \kappa_z, \lambda_\gamma, \lambda_z$$

g^1 and κ are 1 in the SM, the rest are zero



Neutral Triple Gauge Couplings


Probed by ZZ and $Z\gamma$ production

General Lagrangian has **8** TGC parameters

Assume CP conservation

\Rightarrow 4 non-SM TGC parameters:

$$h^3_\gamma, h^3_Z, h^4_\gamma, h^4_Z \text{ all 0 in SM}$$




$\nu\nu\gamma$ candidate event



27

Run 225055 Evt 44315577 Sun Sep 10 03:18:04 2006

ET scale: 114 GeV



Run 225055 Evt 44315577 Sun Sep 10 03:18:04 2006

E scale: 130 GeV

