


DI-BOSONS AND ANOMALOUS COUPLINGS AT D0

Heidi Schellman, Northwestern University




Run II Integrated Luminosity

19 April 2002 - 30 September 2011



DØ Detector



Lepton ID in the DØ Detector


4

muon system

Coverage to $\eta=2$

shielding

electronics





Lagrangian for neutral ($ZZ\gamma/Z\gamma\gamma$)

6

$$L_{gZV} = -ie \hat{e} \left(h_1^n F^{mn} + h_3^n \tilde{F}^{mn} \right) Z_m \frac{(\square + m_V^2)}{M_Z^2} V_m + \left(h_2^V F^{mn} + h_4^V \tilde{F}^{mn} \right) Z^a \frac{(\square + m_V^2)}{M_Z^4} \partial_a \partial_m V_n \dot{u}$$

CP conserving $\rightarrow h_{3,4}^V$ couplings ($V = \gamma, Z$)



$\Delta \neq 0$
ANOMALOUS
COUPLINGS

SM : $h_{3,4}^V = 0$

SM Deviations : $\Delta h_{3,4}^V = h_{3,4}^V - 0$



Lagrangian for charged (WW γ /WWZ)

7

$$\frac{\mathcal{L}_{WWV}}{g_{WWV}} = \boxed{ig_1^V(W_{mn}^* W^m V^n - W_m^* V_n W^{mn}) + ik_V W_m^* W_n V^{mn} + i\frac{l_V}{M_W^2} W_{lm}^* W_n^m V^{nl}}$$

$$-g_4^V W_m^* W_n (\eta^m V^n + \eta^n V^m) + g_5^V e^{mnlr} (W_m^* \eta_l W_n - \eta_l W_m^* W_n) V_r + ik_V W_m^* W_n \tilde{V}^{mn} + i\frac{\tilde{l}_V}{M_W^2} W_{lm}^* W_n^m \tilde{V}^{nl}$$

EM gauge inv. ($g_1^V = 1$), C and P conserving \Rightarrow 5 couplings: $\kappa_V, \lambda_V, g_1^Z$



$\Delta \neq 0$
ANOMALOUS
COUPLINGS

SM: $g_1^Z = \kappa_V = 1, l_V = 0$


SM Deviations:

$\Delta g_1^Z = g_1^Z - 1, \Delta \kappa_V = \kappa_V - 1, \Delta \lambda_V = \lambda_V - 0$

$W\gamma$ and $Z\gamma$



8

- D0 has excellent photon identification due to fine transverse and longitudinal segmentation in the calorimeter.
- Neural network discriminant yields $> 90\%$ purity in $Z\gamma$



PRL 107, 241803 (2011)


9






$$\sigma_{W\gamma} \times \text{BR}(W \rightarrow l\nu) = 7.6 \pm 0.4(\text{stat}) \pm 0.6(\text{syst}) \text{ pb}$$

$$\text{SM@NLO} : \sigma = 7.6 \pm 0.2 \text{ pb}$$

$$(\text{photon } E_T > 15 \text{ GeV}, dR_{(l\gamma)} > 0.7)$$



Anomalous $W\gamma$ couplings



95% CL limits on TGCs:

$$-0.4 < \Delta K_\gamma < 0.4, -0.08 < \lambda_\gamma < 0.07$$


$Z\gamma \rightarrow \nu\nu\gamma$




[Phys. Rev. Lett. 102, 201802 \(2009\)](#)

11



Select interactions with large, significant
Missing transverse momentum



$Z\gamma \rightarrow \nu\nu\gamma$ avoids radiation off of the Z!



Limits on the anomalous
couplings of Z's to
photons.



$$\sigma_{Z\gamma} \times \text{BR}(Z \rightarrow \Pi) = 1089 \pm 40_{\text{(stat)}} \pm 65_{\text{(syst)}} \text{ fb}$$

$$\text{SM@NLO} : \sigma = 1096 \pm 34 \text{ fb}$$

$$\sigma_{Z\gamma} \times \text{BR}(Z \rightarrow \Pi) = 288 \pm 15_{\text{(stat)}} \pm 11_{\text{(syst)}} \text{ fb}$$



$$M_{ll\gamma} > 110 \text{ GeV (FSR removal)} : \text{SM@NLO} : \sigma = 294 \pm 10 \text{ fb}$$

$Z\gamma \square \mu\mu\gamma$

13

	$ll\gamma$	7.2 fb^{-1}
	$\nu\nu\gamma$	3.6 fb^{-1}
	Λ	1.5 TeV
$ h_{03}^Z $	$<$	0.026
$ h_{04}^Z $	$<$	0.0013
$ h_{03}^\gamma $	$<$	0.027
$ h_{04}^\gamma $	$<$	0.0014



Fit the pt distribution to limit anomalous couplings
Combine with the $\nu\nu\gamma$ channel for improved limits.



$WZ \rightarrow |v\bar{v}|$

Phys. Rev. D 85, 112005 (2012)

Three high p_T (isolated) leptons ($\mu\mu\mu$, eee , $ee\mu$, $\mu\mu e$) +MET



$$\sigma_{WZ} = 4.5^{+0.6}_{-0.7} (\text{stat} + \text{syst}) \text{ pb}$$

$$\text{SM@NLO: } \sigma = 3.21 \pm 0.19 \text{ pb}$$

$$(60 < M_{||} < 120 \text{ GeV})$$



ZZ → llll Production

15


Cross section for the ZZ production using fully leptonic final states

- Four high p_T leptons ($\mu\mu\mu\mu$, $eeee$, $ee\mu\mu$)

*ZZ → llll
10 events
.37 background*

$$\sigma_{ZZ} = 1.26^{+0.47}_{-0.37} \text{ (stat + syst) pb}$$

$$\text{SM@NLO: } \sigma = 1.40 \pm 0.10 \text{ pb}$$



ZZ → llνν






Phys. Rev. D 85, 112005 (2012)

Challenges in ννll final states:

- MET reconstruction
- WW background

$$p_T^l = p_T^{\ell\ell} + 2[p_T^\delta + p_T^{\text{recoil}} + p_T^{\text{trkjets}}]$$

Optimal variable is a modified p_T with negative corrections proportional to resolutions (ie. $p_T^\delta < 0$)



Result:

ννll (8.6 fb⁻¹):

$$\sigma_{ZZ} = 1.64^{+0.46}_{-0.46} \text{ (stat+syst) pb}$$

$$\text{SM@NLO: } \sigma = 1.40 \pm 0.10 \text{ pb}$$

ZZ → llνν






Phys. Rev. D 85, 112005 (2012)

Challenges in ννll final states:

- MET reconstruction
- WW background

$$p_T^l = p_T^{\ell\ell} + 2[p_T^\delta + p_T^{\text{recoil}} + p_T^{\text{trkjets}}]$$

Optimal variable is a modified p_T with negative corrections proportional to resolutions (ie. $p_T^\delta < 0$)




Combined:
llll (6.4 fb⁻¹) + ννll (8.6 fb⁻¹):

$$\sigma_{ZZ} = 1.44^{+0.35}_{-0.34} \text{ (stat+syst) pb}$$


SM@NLO: $\sigma = 1.40 \pm 0.10$ pb

Tevatron WZ and ZZ cross sections

WZ



ZZ




Summary of Bosons at the Tevatron + friends

19

Tevatron $p\bar{p}$ at $\sqrt{s} = 1.96$ TeV + LHC 7 TeV

March 2012



Summary

20




- Diboson cross sections have been measured in many all-leptonic channels
- Future updates
 - ▣ Full statistics ZZ
 - ▣ Full statistics WW
 - ▣ Anomalous couplings using all channels

BACKUP SLIDES

Older D0 limits on anomalous couplings from $W\gamma$, WW , WZ



22



Can be interpreted as measurements of the magnetic dipole and quadrupole moments.

$$\mu_W = (1 + \kappa + \lambda) \frac{e}{2M_W} = 2.02^{+0.08}_{-0.09} \frac{e}{2M_W}$$

$$q_W = -(\kappa - \lambda) \frac{e}{M_W^2} = 1.00 \pm 0.09 \frac{e}{M_W^2}$$

Prospects for the future

23

- D0 and CDF are updating to the full statistical sample.
 - ▣ Done for WZ and ZZ channels

- Combining channels and experiments will increase the TGC sensitivity by a factor of 3-5.

- LHC experiments have 10 times the cross section $\rightarrow 10 \text{ fb}^{-1}$ of data \rightarrow factor of 10 in statistics and 3 in sensitivity.


Motivation for Diboson Studies



Probe of the EWSB mechanism

- Test of the SM
- Indirect searches for New Physics

Cross sections, Kinematic distributions, Trilinear Gauge Boson Couplings (TGCs)




Important background to:

- Top
- Higgs
- Beyond the SM

Good understanding is highly valuable

Proving ground for analysis techniques and statistical treatment used in the Tevatron **Higgs searches**





Complementary to Higgs production (same final states/challenges)



DØ – set charged TGC limits using pt in 4 channels



25



Charged Triple Gauge Couplings

Probed by *WW, WZ, and W γ production*


General Lagrangian has **14** parameters

Assume EM gauge invariance and C and P conservation

\Rightarrow 5 TGC parameters:

$$g^1_{Z'}, \kappa_{\gamma'}, \kappa_{Z'}, \lambda_{\gamma'}, \lambda_Z$$

g^1 and κ are 1 in the SM, the rest are zero



Neutral Triple Gauge Couplings


Probed by *ZZ and Z γ production*

General Lagrangian has **8** TGC parameters

Assume CP conservation

\Rightarrow 4 non-SM TGC parameters:

$$h^3_{\gamma'}, h^3_{Z'}, h^4_{\gamma'}, h^4_Z \text{ all 0 in SM}$$




$\nu\nu\gamma$ candidate event



27

Run 225055 Evt 44315577 Sun Sep 10 03:18:04 2006

ET scale: 114 GeV



Run 225055 Evt 44315577 Sun Sep 10 03:18:04 2006

E scale: 130 GeV

