



Thomas Muller, Institut für Experimentelle Kernphysik, KIT



Tevatron:

- Run 1:  $\sqrt{s} = 1.8 \text{ TeV}$  (1992-1996)  
Top quark discovery in 1995 with  $65\text{pb}^{-1}$  (around 20 events each experiment)
- Run 2:  $\sqrt{s} = 1.96 \text{ TeV}$  (2001-2011)  
 $12 \text{ fb}^{-1}$  delivered, on tape  $10 \text{ fb}^{-1}$   
 $8.7 \text{ fb}^{-1}$  being analysed so far




LHC:

- $\sqrt{s} = 7 \text{ TeV}$  (2010-2011)  
 $5.7 \text{ fb}^{-1}$  delivered, on tape  $5 \text{ fb}^{-1}$   
Around  $1M t\bar{t}$  pairs produced per exp.
- Results in  $\sim 60k$  reconstructed  $t\bar{t}$  events in  $e/\mu + \text{jets}$  or in dilepton final state
- $\sqrt{s} = 8 \text{ TeV}$  (2012 ongoing)  
More than  $1M t\bar{t}$  pairs produced  
 $2.8 \text{ fb}^{-1}$  being analysed so far

# All Around the Top

## TOP PRODUCTION

- Cross sections
- Spin-correlations
- Production asymmetries
- Resonances  $X \rightarrow t\bar{t}$ ,  $Y \rightarrow tb$ ,  $Z \rightarrow t\text{jet}$
- Fourth generation  $t'$



The top quark is very special:

- Yukawa couplings close to 1
- Most massive particle known
- The only „free“ quark
- Appears as signal, background and maybe decay product of new states

## THE PARTICLE

- Mass (matter vs. anti-matter)
- Life-time / width
- Charge
- Radiation of Bosons

## TOP DECAY

- Charged Higgs
- $W$  helicity
- Anomalous couplings
- CKM matrix elements

Quarks	$u$	$c$	$t$
up			top
Leptons	$d$	$s$	$b$
down			bottom
	$\nu_e$	$\nu_\mu$	$\nu_\tau$
e- Neutrino	$\mu$ - Neutrino	$\tau$ - Neutrino	
	$e$	$\mu$	$\tau$
electron	muon	tau	
I	II	III	
The Generations of Matter			

# Top Contributions at ICHEP

## TR4 - Top Quark Physics - Roo 216 (09:00-18:00)




time	[id]	title	presenter
09:00	[185]	Inclusive top quark pair production cross - section (ATLAS)	Mr. DERUE, Frederic
09:15	[582]	Top quark pair production cross section at CMS	Dr. RODRIGUEZ MARRERO, Ana
09:30	[186]	Differential top quark pair production (ATLAS)	Dr. CHILDEERS, Taylor
09:45	[586]	Differential cross sections in top pair events at CMS	Dr. ALDAYA MARTIN, Maria
10:00	[689]	Cross section measurements of top quark production at CDF	Mr. CORBO, Matteo
10:15	[83]	NRQCD matching coefficient at next-to-next-to-next-to-leading order	Dr. MARQUARD, Peter
10:30		Refreshment Break	
11:00	[534]	Measurements of the inclusive cross section and of differential distributions in top quark pair production (D0)	Prof. SCHWANENBERGER, Christian
11:15	[187]	Measurements of single top quark production (ATLAS)	Mr. STURM, Philipp
11:30	[594]	Single top production in CMS	Dr. BENELLI, Gabriele
11:45	[538]	Measurements of single top quark production cross sections and $ V_{tb} $ in ppbar collisions at $\sqrt{s}=1.96$ TeV (D0 and CDF)	Dr. PETERS, Yvonne
12:00	[189]	Measurement of the top quark mass (ATLAS)	Dr. SALAMANNA, Giuseppe
12:15	[588]	Measurements of the top quark mass (CMS)	Dr. STADIE, Hartmut
12:30	[590]	Measurement of the top-antitop mass difference (CMS)	Mr. VAN ONSEM, Gerrit
12:45		Lunch Break	
14:00	[693]	Top quark mass measurements at CDF	Dr. LEE, Hyun Su
14:15	[543]	Measurement of the top quark mass in ppbar collisions using events with two leptons (D0)	Dr. BRANDT, Oleg
14:30	[536]	Tevatron and LHC top mass combinations	Dr. DELIOT, Frederic
14:45	[424]	Top Precision Studies at Linear Colliders	Dr. VOS, Marcel
15:00	[852]	Spin correlation and W helicity in top quark events with ATLAS	Dr. JÜNGST, Markus
15:15	[592]	Spin correlations and W helicity in top events with CMS	Dr. SUMOWIDAGDO, Suharyo
15:30		Refreshment Break	
16:00	[541]	Spin correlation in ttbar production (D0)	Dr. PETERS, Yvonne
16:15	[533]	Measurement of top quark properties - electric charge and width (D0)	Prof. SCHWANENBERGER, Christian
16:30	[545]	Combination of CDF and D0 measurements of the W boson helicity in top quark decays	VARNES, Assoc. Prof. Erich Ward
16:45	[190]	Other top quark properties in ATLAS	Dr. LIMOSANI, Antonio
17:00	[601]	Other top quark properties in CMS	Ms. KÜSSEL, Yvonne
17:15	[690]	Top quark properties at CDF	Dr. OH, Youngdo
17:30	[60]	TTbar Spin Correlations at Hadron Colliders	Dr. PARKE, Stephen
17:45	[867]		

## TR4 - Top Quark Physics and TR11 - Particle Astrophysics & Cosmology - Room 216 (09:00-18:00)

time	[id]	title	presenter
09:00	[191]	Charge asymmetry in top pairs at ATLAS	Dr. GIORDANI, Mario
09:15	[587]	Measurement of the charge asymmetry in top quark pair production in pp collisions (CMS)	Dr. CHWALEK, Thorsten
09:30	[684]	Asymmetry measurements in t-tbar at CDF	Dr. HAYS, Chris
09:45	[532]	Measurement of the forward-backward charge asymmetry in top quark pair production (D0)	Dr. GROHSJEAN, Alexander Josef
10:00	[474]	Top quark forward-backward asymmetry from gauged flavor symmetry	Prof. BABU, Kaladi S.
10:15	[318]	Dynamical Origin of the Correlation between the Top Quark Production Asymmetries $A_{\{FB\}^{\{S\}}}$ and $A_{\{FB\}^{\{ell\}}}$	Dr. BERGER, Edmond
10:30		Refreshment Break	
11:00	[513]	Diagnosing top-quark Forward-Backward Asymmetry	Dr. GUPTA, Sudhir Kumar
11:15	[188]	Searches in s-channel single top quark production at ATLAS	Dr. ALVAREZ GONZALEZ, Barbara
11:30	[853]	FCNC in top quark production and decay at ATLAS	Dr. CRISTINZIANI, Markus
11:45	[599]	Search for FCNC in top pair events in pp collisions (CMS)	Dr. CHAO, Yuan
12:00	[52]	CP violation in top-quark physics	Prof. VALENCIA, German
12:15	[495]	Search for a Narrow ttbar Resonance in ppbar Collisions at $\sqrt{s} = 1.96$ TeV (Combined D0, CDF)	ASSOC. PROF. VARNES, Erich
12:30		Lunch Break	
14:00	[585]	Measurement of the top pair invariant mass distribution and search for New Physics (CMS)	Prof. BLEKMAN, Freya
14:15	[192]	Searches for ttbar resonances (ATLAS)	Dr. VOS, Marcel
14:30	[76]	Z' signals in polarised top-antitop final states	Mr. MIMASU, Ken
14:45	[443]	A charged Z' to explain the apparent disagreement in top-antitop asymmetries between Tevatron and LHC	Ms. COLUCCIO LESKOW, Estefania
15:00	[46]	Top Decays with Flavor Changing Neutral Higgs Interactions at the LHC	Prof. KAO, Chung
15:15	[540]	Search for anomalous Wtb couplings in ppbar collisions at $\sqrt{s} = 1.96$ TeV (D0)	Prof. BLOOM, Kenneth



Apologies for not mentioning all results

# 1. PAIR PRODUCTION OF TOP QUARKS



# Decay Channels

## Top Pair Decay Channels



- $t \rightarrow Wb$



Events classified by W decay

- “Lepton [e, $\mu$ ] + jets” (34%)  
 $tt \rightarrow bl\nu b\bar{q}\bar{q}'$


- “Dilepton [e, $\mu$ ]” (6%)  
 $tt \rightarrow bl\nu bl\nu$

- “All jets” (46%)  
 $tt \rightarrow b\bar{q}\bar{q}' b\bar{q}\bar{q}'$

- “Tau + jets” (15%)  
 $tt \rightarrow b\tau\nu b\bar{q}\bar{q}'$



# 1.1 Cross Section Measurements of $t\bar{t}$ Production



Finding the top:




Signal:

- Triggering on lepton
- High missing transverse energy ( $E_T$ )
- High  $E_T$  jets, central and spherical
- Two b-jets (displaced vertex)


Background:


- Dilepton: Z+Jets, Single Top (tW), QCD, W+Jets
- Lepton+Jets: W+Jets, Single Top
- All Hadronic: QCD multi jet events





Determination of the cross section

$$\sigma_{t\bar{t}} = \frac{N_{obs} - N_{bgd}}{\varepsilon_{t\bar{t}} \cdot \int L dt}$$





**CDF Run II****Conf. Note 10878**

- Consistency amongst various channels
- Limitation from systematic uncertainties (JES, b-tab, W+jets)
- Sensitive to NNLO predictions (Bernreuter, Czakon, Mitov, arXiv:1204.5201)

**D0 Run II****arXiv:1105.5834**

**ATLAS (ATLAS-CONF-2012-024)**

- Combination done as product of the individual likelihoods of each channel
- $\sigma = 177 \pm 3 \text{ (stat)} \pm 7 \text{ (syst.)} \pm 7 \text{ (lum.) pb}$

**ATLAS Preliminary****CMS (CMS-PAS-TOP-11-024)**

- Combination done using a binned maximum likelihood fit
- $\sigma = 165.8 \pm 2.2(\text{stat}) \pm 10.6(\text{syst.}) \pm 7.8(\text{lum.}) \text{pb}$



Results are compatible with NNLO calculations

# Top-Antitop Production Cross Sections

**CMS (CMS-PAS-TOP-12-006/7)**

- New measurement at 8 TeV !
- Lepton + jets and di-lepton channels combined:

$$\sigma = 227 \pm 3 \text{ (stat)} \pm 11 \text{ (syst.)} \pm 10 \text{ (lum.) pb}$$






**CMS**

$$\sigma(8\text{TeV})/\sigma(7\text{TeV}) = 1.41 \pm 0.11; \text{ no correlation assumed}$$




# Top-Antitop Differential Cross Sections

- Measure differential cross sections in the lepton+jet channel
- Important test of QCD
- Event selections similar to the cross section analyses
- Bin-to-bin unfolding to parton level at CMS

$$\frac{1}{\sigma} \frac{d\sigma^i}{dX} = \frac{1}{\sigma} \frac{N_{data}^i - N_{bkg}^i}{\epsilon \cdot \mathcal{L}}$$



**ATLAS  
Paper in Prep.**





**CMS  
PAS-TOP-11-013**

In conclusion, tt production is well described by SM calculations

# 1.2 Top-Antitop Charge Asymmetry


NLO QCD: interference of higher order diagrams leads to asymmetry for  $t\bar{t}$  produced through  $q\bar{q}$  annihilation:

- Top quark is emitted preferentially in direction of the incoming quark
- Antitop quark opposite
- Production through new processes may lead to different asymmetries



- At Tevatron: define forward-backward asymmetry


$$A^{t\bar{t}} = \frac{N(\Delta y > 0) - N(\Delta y < 0)}{N(\Delta y > 0) + N(\Delta y < 0)}$$





- At LHC: define asymmetry in the widths of rapidity distributions of t, t̄

$$A_c = \frac{N(\Delta |y| > 0) - N(\Delta |y| < 0)}{N(\Delta |y| > 0) + N(\Delta |y| < 0)}$$



$$\Delta|y| = |y_t| - |y_{\bar{t}}|$$




# Asymmetries at the Tevatron



## CDF-Note 10807



$$A_{FB} = 0.162 \pm 0.041(\text{stat}) \pm 0.022(\text{syst})$$



Measured asymmetry on detector level after bkg subtraction:

$$A_{FB} \text{ det} = 0.092 \pm 0.037 \text{ (stat+syst)}$$

$$\text{MC@NLO: } A_{FB} \text{ det} = 0.024 \pm 0.007$$


Measured asymmetry on parton level:

$$A_{FB} = 0.196 \pm 0.065 \text{ (stat+syst)}$$

D0 results in the di-lepton channel:



$$A_{FB} = 0.118 \pm 0.032$$

## Summary:



Both CDF and D0 see significant asymmetry in  $t\bar{t}$  production in all channels with strong dependence on  $m_{t\bar{t}}$ , in conflict with the SM

# Asymmetries at the LHC




**CMS PAPER TOP-11-030**

ATLAS-CONF-2012-057

- ATLAS:  $A_C = 0.029 \pm 0.018 \text{ (stat.)} \pm 0.014 \text{ (syst.)}$
- CMS: Corrected:  $A_C = 0.004 \pm 0.010 \text{ (stat.)} \pm 0.011 \text{ (syst.)}$
- Theory (Kühn, Rodrigo):  $A_C = 0.0115 \pm 0.0006$

# Differential asymmetries at the LHC



Asymmetry measured in pT, y or invariant mass of the top pair system

Measured asymmetries are compared to

- SM prediction at NLO [1]
- SM simulation at NLO (POWHEG)
- BSM prediction with an effective axial-vector coupling of the gluon at the one-loop level (EAG) [2]; can explain the strong dependence of AFB on  $m(t\bar{t})$  as seen by CDF

[1] Kühn, Rodrigo - arXiv:1109.6830 [2] Gabrielli, Racioppi, Raidal - PRD 85 (2012) 074021, arXiv:1112.5885; arXiv:1203.1488


At LHC data are compatible with SM and with BSM prediction.  
Tevatron results display much larger asymmetry than SM

An LHC run at low energies would be helpful (once we explored the energy frontier).


# 1.3 Search for $t\bar{t}$ Resonances

CDF:

lepton+jets selection, no b-tagging requirements




[Phys. Rev. D 84, 072004 \(2011\)](#)




$M_X > 900 \text{ GeV} @ 95\% \text{ C.L.}$

DØ:







lepton+jets selection, at least one b-tagged jet



[arXiv:111.1271v1](#)



$M_X > 835 \text{ GeV} @ 95\% \text{ C.L.}$

**CMS TOP-11-009/010, EXO-11-006/093****Topcolor Z'****Extra-dimensions****arXiv:1205.5371****So far data compatible with SM up to 1...1.6 TeV**

## 2. SINGLE TOP PRODUCTION


Observation of single top production:

- cross section  $\propto V_{tb}^2$
- study top-polarization and EWK top interaction


Test of non-SM phenomena:

- 4th generation
- FCNC couplings
- $W'$ ,  $H^\pm$
- anomalous  $W_{tb}$  couplings


s-channel



t-channel



Wt-channel



Main backgrounds:

- s-channel: Top pair,  $W + (\text{HF})$  jets, QCD
- t-channel: Top pair,  $W + (\text{HF})$  jets, QCD
- Wt-channel: Top pair,  $Z + (\text{HF})$  jets, QCD


Signal – background discrimination:

- Tevatron: multivariate methods (neural networks, boosted decision trees, matrix element method)
- LHC: cut-based or multivariate method



Collider	s-channel: $\sigma_{tb}$	t-channel: $\sigma_{tqb}$	Wt-channel: $\sigma_{tw}$
Tevatron: $p\bar{p}$ (1.96 TeV)	1.05 pb	2.08 pb	0.22 pb
LHC: pp (7 TeV)	4.6 pb	66 pb	15.7 pb

# Single Top at the Tevatron

## CDF Note 10878



## PLB 705, 313 (2011)



$$\sigma_s = 1.81^{+0.63}_{-0.58} \text{ pb}$$

$$\sigma_t = 1.49^{+0.47}_{-0.42} \text{ pb}$$

$V_{tb} > 0.78$  @ 95% C.L.

$$\sigma_s = 0.98 \pm 0.63 \text{ pb}$$

$$\sigma_t = 2.90 \pm 0.59 \text{ pb}$$

$V_{tb} > 0.79$  @ 95% C.L.

$$|V_{tb,meas}|^2 = \frac{\sigma_{meas}}{\sigma_{SM}} \cdot |V_{tb,SM}|^2$$

- No assumption about number of generations
- Assumption:  $|V_{ts}|^2 + |V_{td}|^2 \ll |V_{tb}|^2$



# Single Top at the LHC

ATLAS:  $\sigma = 83 \pm 4 \text{ (stat.)} \pm 20\text{-}19 \text{ (syst.) pb}$

ATLAS-CONF-1205.3130

CMS:  $s = 70.2 \pm 5.2 \text{ (stat.)} \pm 10.4 \text{ (syst.)} \pm 3.4 \text{ (lumi.) pb}$

CMS-PAS-TOP-11-021




# Single Top: tW-channel

Final state: 2 isolated leptons, 1 b-tagged jet and MET (as dilepton top pair but with 1 b jet less)


## ATLAS ([arXiv:1205.5764](https://arxiv.org/abs/1205.5764))

- Isolated lepton  $p_T > 25 \text{ GeV}/c$
- $\cancel{E}_T > 50 \text{ GeV}$
- No b tagging requirement
- MVA (BDT) based analysis



## CMS ([CMS-PAS-TOP-11-022](https://cds.cern.ch/record/1402220))

- Isolated lepton  $p_T > 20 \text{ GeV}/c$
- $\cancel{E}_T > 30 \text{ GeV}$
- 2nd b-jet veto is applied for signal region
- Cut based analysis




ATLAS:  $\sigma = 16.8 \pm 2.9 \text{ (stat.)} \pm 4.9 \text{ (syst.) pb}$

CMS:  $\sigma = 22^{+9}_{-7} \text{ (stat. + syst.) pb}$

# Search for tb Resonances



- Right-handed  $W'_R$  with SM-like couplings chosen as benchmark model
- Similar signature as single top s-channel (1 lepton, MET, 2 jets and at least 1  $b$  tag)



CMS-PAS-EXO-12-001

**CMS**



- BDT used to discriminate signal from background



<http://arxiv.org/abs/1205.1016>

**ATLAS**



- Invariant mass of  $tb$  system used as discriminating variable



Observed limits (ATLAS, CMS):  $m_{W'_R} > 1.13 / 1.85 \text{ TeV}$


### 3. DECAY PHYSICS: W HELICITY

Three possible helicities:



$$F_0 = \frac{m_t^2}{2M_W^2 + m_t^2}$$

Distribution of Angle  $\theta^*$  between charged lepton in W system and W-Boson in Top-Quark system:



$$\frac{dN_{h_w=-1}}{d(\cos\theta^*)} \sim \frac{3}{8}(1 - \cos\theta^*)^2$$

**SM:**  
 $F_- = 0.31$

$$\frac{dN_{h_w=0}}{d(\cos\theta^*)} \sim \frac{3}{4}(1 - \cos^2\theta^*)$$

$F_0 = 0.69$

$$\frac{dN_{h_w=+1}}{d(\cos\theta^*)} \sim \frac{3}{8}(1 + \cos\theta^*)^2$$




$F_+ = 0.001$

A. Czarnecki, J. G. Körner, J. H. Piclum, Phys. Rev. D 81, 111503 (2010)

# Tevatron Results on W Helicity

With three helicity fractions, there are two independent quantities to measure (3rd fraction is fixed since  $\sum f = 1$ )

- We choose to measure  $f_0$  and  $f_+$
- Can either measure both fractions simultaneously (2D fit)
- or fix one fraction to its SM value and measure the other (1D fit)




## Tevatron Combo


$$\begin{aligned} \text{2D: } & f_0 = 0.722 \pm 0.062 \pm 0.052 \\ & f_+ = -0.033 \pm 0.034 \pm 0.031 \\ \text{1D: } & f_0 = 0.682 \pm 0.035 \pm 0.046 \\ & f_+ = -0.015 \pm 0.018 \pm 0.030 \end{aligned}$$

**PRD85, 071106 (2012)**

# LHC Results




*Combine  
l+jets and  
dilepton*




$$F_0 = 0.67 \pm 0.03 \text{ (stat.)} \pm 0.06 \text{ (syst.)},$$

$$F_L = 0.32 \pm 0.02 \text{ (stat.)} \pm 0.03 \text{ (syst.)},$$

$$F_R = 0.01 \pm 0.01 \text{ (stat.)} \pm 0.04 \text{ (syst.)}.$$



**CMS: l+jets**



$$F_0 = 0.567 \pm 0.074 \text{ (stat.)} \pm 0.047 \text{ (syst.)}$$


$$F_L = 0.393 \pm 0.045 \text{ (stat.)} \pm 0.029 \text{ (syst.)},$$

# 4. TOP QUARK PROPERTIES


# 4.1 Top Mass Measurements

Measurement of the top quark mass:





- Template method:  
fit MC generated distributions assuming different  $M_{top}$  to data
- Matrix element method:  
probability based on LO  $t\bar{t}$  matrix element using full kinematics of the event
- Ideogram method  
event likelihood computed as a convolution of a Gaussian resolution function with a Breit-Wigner (signal)
- Calibration curve method  
di-lepton channel using  $mt^2$



## Sources of uncertainties at Tevatron





# JES vs $M_{\text{top}}$ in all four experiments




# Top Quark Mass Combinations

## – Tevatron combination and perspectives



Expect to reach precision of 0.7-0.8 GeV

## – LHC combination and perspectives



TeV:  $m_t^{\text{comb}} = 173.18 \pm 0.56 \text{ (stat)} \pm 0.75 \text{ (syst)} \text{ GeV}$   
 $= 173.18 \pm 0.94 \text{ GeV}$

LHC:  $m_{\text{top}} = 173.3 \pm 0.5 \text{ (stat)} \pm 1.3 \text{ (syst)} \text{ GeV}$   
 $= 173.3 \pm 1.4 \text{ GeV}$


**ATLAS-CONF-2012-095**

**CMS PAS TOP-12-001**

## 4.2 $t\bar{t}$ Mass Difference

- CPT theorem predicts masses of particles and antiparticles to be equal
  - Top-quark only particle with color charge to test this invariance

- Analysis Principle:



- Analysis of  $\mu+jets$  events

- 1 isolated muon, 4 high- $p_T$  jets
- Kinematic event reconstruction of 12 permutations
- Measurement of top- and antitop-quark masses with Ideogram method
  - Taking b-tagging information into account to reduce background events (mainly  $W+jets$ )

Test CPT invariance in the top sector

$$\Delta M_t = M_{top} - M_{\bar{top}}$$



- Reconstruction of the hadronic side: compare  $\ell^++\text{jets}$  and  $\ell^-\text{-jets}$  events
- Use kinematic fit, and event-per-event likelihood for  $\ell^-$  and  $\ell^+$  separately

$$\Delta m_t = -0.44 \pm 0.46 \text{ (stat.)} \pm 0.27 \text{ (syst.) GeV}$$

Most systematic effects cancel out !  
 → the measurement is stat. limited

World's best so far

Consistent with SM,  
 Consistent between e and  $\mu$





CMS-TOP-11-019

## 4.3 Width of the Top Quark

### t-channel cross section:

$$\sigma(p\bar{p} \rightarrow tqb + X) = 2.90 \pm 0.59 \text{ pb}$$

$m_t = 172.5 \text{ GeV}$




Phys. Rev. D84 012008 (2011)

$$\Gamma_t = 2.00^{+0.47}_{-0.43} \text{ GeV}$$

### partial decay width:

$$R = 0.90 \pm 0.04 \text{ (stat+syst)}$$



$$\begin{aligned}\tau_{\text{top}} &\propto \left(\frac{M_W}{M_{\text{top}}}\right)^3 \\ \tau_{\text{top}} &\approx 4.7 \cdot 10^{-25} \text{ s}\end{aligned}$$

$$\tau_t = (3.29^{+0.90}_{-0.63}) \times 10^{-25} \text{ s}$$

⇒ most precise determination

Phys. Rev. D 85, 091104(2012)

# 4.4 $t\bar{t}+V$ Production

Trilepton channel:  $\sigma(t\bar{t}Z \rightarrow l + \text{jets} + (Z \rightarrow ll))$

Same-sign dilepton channel:  $\sigma(t\bar{t}V \rightarrow l + \text{jets} + (W \rightarrow l\nu) \text{ or } (Z \rightarrow ll))$




with  $l = e$  or  $\mu$

■ Selection:

- 3 leptons: 2 opposite-charge and same-flavor leptons ( $Z^{\text{cand}}$ )
- 3 jets (2  $b$  tagged),  $H_T > 120$  GeV

■ Background estimation from data:

- $t\bar{t}$ : cross-flavor dilepton events
- $Z + \text{jets}$  and diboson: trilepton, no  $b$  tag events



First measurement of  $t\bar{t}V$ :

■ Result combining all 7 channels:

$$\sigma(t\bar{t}V) = 0.51^{+0.15}_{-0.13} (\text{stat.})^{+0.04}_{-0.02} (\text{syst.}) \text{ pb}$$

→ Significance of  $4.67 \sigma$

**CMS-TOP-12-014**

# CONCLUSIONS

Seventeen years after its discovery, top quark physics is ever increasing in fascination.

The Tevatron will leave a legacy in precision measurements, analysis methods and searches at the energy frontier.

The LHC has had a phantastic start and has been taking over from the Tevatron. The LHC has become a top factory.

Theory has been accompanying us in this exciting programme, being a match in terms of precision and paving the way in predicting new phenomena.

So far, agreement with SM predictions is astounding except for one mystery which seems to solidify. We need to get hold of the asymmetry in all possible ways.

**Exciting times are awaiting us !**

Supported by the State of Baden-Württemberg, HGF, the DFG and the BMBF



# APENDIX

# Top Quark Polarisation


In BSM possibility of top polarisation, for instance through axi-gluon or  $W'$  couplings.

D. Krohn, T. Liu, J. Shelton et al.,  
Phys.Rev. D84 (2011) 074034

Angular distribution of charged leptons sensitive to polarisation:

$$\frac{1}{\Gamma} \frac{d\Gamma}{d \cos \theta_{i,n}} = \frac{1}{2} (1 + 2\mathcal{P}_n \kappa_i \cos \theta_{i,n})$$

$n$  direction of the top in the  $t\bar{t}$  rest frame  
 $\kappa_i = 1$  for leptons




**CMS PAS TOP-12-016**

$P_n = -0.009 \pm 0.029 \pm 0.041$

Consistent with Standard Model

# Ratio $R_t = \sigma_t / \sigma_{\bar{t}}$


- $R_t$  is sensitive to the u/d PDFs; naïve expectation  $R_t = 2$
- Measurement of  $\sigma_t$  using binned max. likelihood fit to NN output in 2-jets and 3-jets data split according to charge of lepton



$$\sigma_t = 53.2 \pm 10.8 \text{ pb}$$

$$\sigma_{\bar{t}} = 29.5^{+7.4}_{-7.5} \text{ pb}$$





$$R_t = 1.81^{+0.23}_{-0.22}$$




**ATLAS-CONF-2012-056**

Correlations between the individual cross section measurements are taken into account, leading to a reduction of the uncertainty


The measurement is in agreement with the predictions based on various global PDF sets that range from 1.86 to 2.07.








3 leptons + 0 jets



3 leptons + 1 jet



3 leptons +  $\geq 2$  jets



arXiv:1206.0257





Christian Schwanenberger





# Spin Correlations

- Spins of  $t$  and  $\bar{t}$  are predicted to be correlated in SM
  - Top decay before hadronization allows to measure top spin from its decay products
  - At LHC at low  $m(t\bar{t})$   $t\bar{t}$  production dominated by like-helicity gluon pairs  $\rightarrow$  like-helicity  $t\bar{t}$  pairs.
  - In dilepton final states this results in correlations between the leptons in the azimuthal angle  $\Delta\phi$  in the lab frame (Mahlon & Parke)

- Complementary between Tevatron and LHC




# Spin correlation at the LHC



Phys. Rev. Lett. 108  
212001 (2012)


$$A = \frac{N(\uparrow\uparrow) + N(\downarrow\downarrow) - N(\uparrow\downarrow) - N(\downarrow\uparrow)}{N(\uparrow\uparrow) + N(\downarrow\downarrow) + N(\uparrow\downarrow) + N(\downarrow\uparrow)}.$$



SM:  $f = 1.0, A = 0.31$

$f = 0.74 \pm 0.08 \text{ (stat)} \pm 0.24 \text{ (syst)},$   
 $A = 0.24 \pm 0.02 \text{ (stat)} \pm 0.08 \text{ (syst)}$

CMS-PAS-TOP-12-004





Correlation values in agreement with SM assumptions

# Top Quark Cross Sections (Th.)

Cross sections (pb) [ $m_{top} = 173$ GeV]	s-channel	t-channel	tW channel	top pair
Tevatron: ppbar@1.96TeV	$1.046^{+0.002}_{-0.01}{}^{+0.06}_{-0.056}$	$2.08^{+0.00}_{-0.04} \pm 0.12$	$0.22 \pm 0.08$	$7.08^{+0.00}_{-0.24}{}^{+0.36}_{-0.27}$
LHC: pp @ 7 TeV	$4.56 \pm 0.07^{+0.18}_{-0.17}$	$65.9^{+2.1}_{-0.7}{}^{+1.5}_{-1.7}$	$15.6 \pm 0.4^{+1.0}_{-1.2}$	$163^{+7}_{-5} \pm 9$
LHC: pp @ 8 TeV	$5.55 \pm 0.08 \pm 0.21$	$87.2^{+2.8}_{-1.0}{}^{+2.0}_{-2.2}$	$22.2 \pm 0.6 \pm 1.4$	$234^{+10}_{-7} \pm 12$

N. Kidonakis  
[arxiv.org/pdf/1205.3453v1](http://arxiv.org/pdf/1205.3453v1.pdf) (2012)


# Top Couplings from production



$$|V_{tb}| = 0.92 +0.10-0.08 \text{ (stat.+sys.)} \pm 0.05 \text{ (theory)}$$

Other single top studies:  
FCNC single top quark production


Search for:  
 $qg \rightarrow t \rightarrow W(\rightarrow \ell\nu)b$



mT+ETmiss>60 GeV

Require only 1 jet (b-tagged)

$$\sigma(qg \rightarrow t) \cdot B(t \rightarrow Wb) < 3.9 \text{ pb (95% CL)}$$



$$B(t \rightarrow ug) < 5.7 \cdot 10^{-5}, B(t \rightarrow cg) < 2.7 \cdot 10^{-4}$$

CMS-TOP-11-028 is 0.34%,