


Searches for SUSY in events with 3rd generation particles at CMS

On behalf of the CMS Collaboration



Altan CAKIR
DESY

ICHEP 2012, 05 July 2012,
Melbourne, Australia

SUSY in the 3rd Generation


- SUSY can solve many problems intrinsic to SM:
 - Hierarchy problem
 - Unification of forces at a high energy scale
- If R-parity is conserved: Lightest SUSY particle (LSP) is stable
 - ➔ natural Dark Matter (DM) candidate

In the third generation SUSY particles ➔ sizeable mixing

\tilde{t}_1 and \tilde{b}_1 can be lighter than the other squarks
 $\tilde{\tau}_1$ can be lighter than the other sleptons

➔ 3rd generation sfermions:


- Can be produced in pairs or appear in the gluino cascade decay
- Produce *b*, *t* or τ in their decay




Searching for 3rd Generation SUSY in CMS*

☞ Search for SUSY in Final States with a **Single Lepton, B-jets**, and Missing Transverse Energy in Proton-Proton Collisions at $\sqrt{s} = 7$ TeV – [PAS-SUS-11-028](#)

☞ Search for SUSY in events with a **single lepton** and jets using templates
[PAS-SUS-11-027](#)

 Search for physics beyond the standard model in events with **tau leptons** in the presence of multijets and large momentum imbalance in pp collisions at $\sqrt{s} = 7$ TeV [PAS-SUS-12-004](#)

 Search for new physics in events with **same-sign dileptons and b-tagged jets** in pp collisions at $\sqrt{s} = 8$ TeV [CMS-SUS-12-017](#)

 Search for new physics in events with **b-quark jets** and missing transverse energy in pp collisions at $\sqrt{s} = 7$ TeV [PAS-SUS-12-003](#)



☞ *Razor, For Alpha_{Tb} and M_{T2b} analyses → see Will Reece and Seema Sharma's talks!*

* <https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsSUS>

Search for SUSY in Final States with a Single Lepton, B-jets, and Missing Transverse Energy in Proton-Proton Collisions at $\sqrt{s} = 7$ TeV

PAS-SUS-11-028

☞ The analysis is performed in three channels according to the number of b-tags: **exactly one b-tag, exactly two b-tags and three or more b-tags.**






☞ $H_T > 375 \text{ GeV}$, $N_{\text{jets}} \geq 4$ and one exact isolated lepton (e, μ) (left)
 $\rightarrow N_{\text{bjets}} \geq 1$ required for H_T plot (right)

Search for SUSY in Final States with a Single Lepton, B-jets, and Missing Transverse Energy in Proton-Proton Collisions at $\sqrt{s} = 7$ TeV

PAS-SUS-11-028


Background Estimation Method - Factorization




➡ The measured and the predicted number of events in signal region (D) are in good agreement.

Interpretation of the results

cMSSM



Simplified Model



- ☞ Limits are set using the **CLs** method with a test statistic given by a profile likelihood ratio.
- ☞ The limits are based on ≥ 1 btag and ≥ 3 btag for cMSSM and SMS models, respectively.

☑ No deviation from the SM has been found

☑ Upper limits have been set on production cross-sections for both models.

PAS-SUS-11-028

Search for SUSY in events with a single lepton and jets using templates


PAS-SUS-11-027

☞ The analysis is done as a function of the number of identified b-quark (including 0-btag) jets in the event.


Methodology for Background Estimation

The dominant SM backgrounds (Wjets, TTjets) can be obtained from data:

- Use hadronic component of events
- Fit the parameters of a model for the genuine MET in a control region defined by H_T
- Apply individual MET models for W^+ jets, W^- jets and TTjets
- Use W+jets/TTJets ratios for events with 0,1 or ≥ 2 tagged bjets determined using template fit.




The fitted parameter α as obtained from data:




cakir@cern.ch



Interpretation of the results



☞ The observed and expected median limits are based on **all btags bins** (0,1,2 btag).



☞ The signal region is defined by $H_T > 750$ GeV and $MET > 250$ GeV with **at least 2 b-jet bin**.

☑ No excess has been observed



☑ The results with $n_{bjet} \geq 2$ are used for interpretation in simplified model.

PAS-SUS-11-027

Search for physics SUSY in events with tau leptons in the presence of multijets and large momentum imbalance at $\sqrt{s}=7$ TeV



PAS-SUS-12-004

☞ The analysis is performed with one or more hadronically decaying τ -leptons, highly energetic jets and large momentum imbalance in the final state.



Interpretation of the results


cMSSM



- ✓ Single tau analysis sensitive to lower m_0
- ✓ Tau-tau analyses sensitive to higher m_0

PAS-SUS-12-004

Interpretation of the results



☞ 95% CL cross section upper limits for the limits on the mass of the gluino and the LSP (left) and cross section upper limits as a function of gluino mass in the GMSB

☑ No excess beyond the SM expectations has been found for both single and di-tau final states.



PAS-SUS-12-004

Search for new physics in events with same-sign dileptons and b-tagged jets in pp collisions at $\sqrt{s} = 8$ TeV

PAS-SUS-12-017



Dominant SM backgrounds:

- $t\bar{t}$ bar with “fake” leptons \rightarrow fake ratio / isolation extrapolation
- Charge mis-reconstruction \rightarrow use Z^0 's for charge
- Rare SM processes with high $P_{T,l}$ leptons and bjets \rightarrow estimate from MC




Background Estimation

- ☞ Define pre-selection regions in $MET - H_T$
 - ✓ Validate data-driven background estimates with $\sim 10-100$ events
- ☞ Define search regions by adding MET, H_T requirements \rightarrow Data driven techniques




Interpretation of the results

Simplified Model



gluino \rightarrow virtual top squarks



gluino \rightarrow on-shell top squarks

- Glueballs have been excluded with masses up to approximately 880 GeV
- Lower limit on the bottom squark mass of 408 GeV.

PAS-SUS-12-017



\rightarrow For multiple bottom final states \rightarrow see Pablo Arbol's talk!

Altan CAKIR | Searches for SUSY in events with 3rd generation particles at CMS


ICHEP 2012, Melbourne, Australia | Page 13



Search for new physics in events with b-quark jets and missing transverse energy in pp collisions at $\sqrt{s} = 7$ TeV


PAS-SUS-12-003

☞ The SM background estimates from the data-based background procedures in comparison with the observed number of events in data.



Search for new physics in events with b-quark jets and missing transverse energy in pp collisions at $\sqrt{s} = 7$ TeV

Simplified Models



☞ 95% CL observed cross section upper limits for multi top and bottom quarks final states.

☑ No excess beyond the SM expectations

PAS-SUS-12-003

☑ Set limits on new physics in the context of the b-jet rich simplified model

Summary

👉 Variety of searches for SUSY events with 3rd generation squarks and sleptons

👉 Exploring signatures with heavy quarks or tau leptons using **L = 4.98/fb data at $\sqrt{s} = 7$ TeV (2011)** and **L = 3.95/fb data at $\sqrt{s} = 8$ TeV (2012)** with CMS

☑ 1 lepton + 1 and 3 bjets \rightarrow cMSSM and simplified model (multiple top quarks)

☑ 1 lepton + 0,1,2 bjets \rightarrow cMSSM and simplified model (multiple top quarks)

☑ 1 or 2 tau leptons \rightarrow GMSB scenario and simplified model (multiple tau leptons)


☑ 2 lepton (SS) + 2 bjets \rightarrow cMSSM and simplified model (multiple top quarks)

☑ 0 lepton + 1,2,3 bjets \rightarrow cMSSM and simplified models (multiple top and bottom quarks)

✓ No significant excess observed over SM expectations \rightarrow Limits on the masses of the sparticles in a various SUSY scenarios




Compact Muon Solenoid (CMS) Experiment



cakir@cern.ch




B-Tagging Schema



B-Tagging Algorithm

"Track Counting" algorithm: This is a very simple tag, exploiting the long lifetime of B hadrons. It calculates the signed impact parameter significance of all good tracks, and orders them by decreasing significance. Its b tag discriminator is defined as the significance of the N'th track. It comes in two variations for $N = 2$ (high efficiency) or $N = 3$ (high purity).

"Combined Secondary Vertex" algorithm: This sophisticated and complex tag exploits all known variables, which can distinguish b from non-b jets. Its goal is to provide optimal b tag performance, by combining information about impact parameter significance, the secondary vertex and jet kinematics.



- > Factorization method (also called ABCD method) is used:
- > Hypothesis: H_T and Y_{MET} are **not strongly correlated**



$$k := \frac{N_A \times N_D}{N_B \times N_C} \quad \hat{N}_D := k \frac{N_B \times N_C}{N_A}$$

*K ≠ 1 to account for correlation
where K is taken from MC*

1.20 ± 0.13

Control
regions

Signal
region

Regions Boundaries

H_T Y_{MET}

A: $375 < H_T < 650$ $3.25 < Y_{MET} < 5.5$

B: $650 < H_T$ $3.25 < Y_{MET} < 5.5$

C: $375 < H_T < 650$ $5.5 < Y_{MET}$


D: $650 < H_T$ $5.5 < Y_{MET}$

Variation	$\Delta\kappa$ (0 b-tags)	$\Delta\kappa$ (1 b-tag)	$\Delta\kappa$ (2 b-tags)	$\Delta\kappa$ (≥ 3 b-tags)	$\Delta\kappa$ (≥ 1 b-tags)
JES	±7.5%	±2.2%	±1.4%	±4.0%	±1.5%
JER	±4.2%	±1.7%	±1.8%	±5.5%	±1.1%
p_T^{lepton}	±0.6%	±1.5%	±0.7%	±1.2%	±0.7%
Uncl. energy	±3.1%	±0.3%	±0.7%	±0.8%	±0.4%
Pile-up	±1.7%	±0.5%	±1.1%	±0.9%	±0.8%
B-tag SF	±0.3%	±0.1%	±0.1%	±0.1%	±0.0%
Mis-tag SF	±0.0%	±0.1%	±0.0%	±0.1%	±0.1%
Cross-sect. var.	±3.4%	±1.0%	±2.0%	±1.4%	±0.4%
0b-data	±10.0%	±10.0%	±10.0%	±10.0%	±10.0%
Total syst. uncert.	±14.1%	±10.6%	±10.5%	±12.3%	±10.3%
Stat. error	±11.8%	±4.9%	±4.6%	±6.2%	±3.3%


Variation	ΔN_D (0 b-tags)	ΔN_D (1 b-tag)	ΔN_D (2 b-tags)	ΔN_D (≥ 3 b-tags)	ΔN_D (≥ 1 b-tags)
JES	±26.3 %	±20.9 %	±17.9 %	±17.1 %	±19.6 %
JER	±7.7 %	±6.1 %	±7.0 %	±9.5 %	±6.7 %
p_T^{lepton}	±2.3 %	±1.8 %	±2.0 %	±2.2 %	±1.9 %
Uncl. energy	±2.1 %	±0.3 %	±0.3 %	±0.3 %	±0.3 %
Pile-up	±0.2 %	±0.7 %	±0.3 %	±0.2 %	±0.5 %
B-tag SF	±2.3 %	±0.9 %	±3.8 %	±7.4 %	±1.5 %
Mis-tag SF	±1.9 %	±0.6 %	±1.1 %	±5.7 %	±1.2 %
Model uncert.	±16.0 %	±16.0 %	±16.0 %	±16.0 %	±16.0 %
Lep. trig. & ID	±3.0 %	±3.0 %	±3.0 %	±3.0 %	±3.0 %
Lumi. uncert.	±2.2 %	±2.2 %	±2.2 %	±2.2 %	±2.2 %
Total uncert.	±32.2 %	±27.3 %	±25.7 %	±27.3 %	±26.6 %
Stat. error	±8.4 %	±3.4 %	±3.1 %	±4.3 %	±2.2 %

☞ Systematical uncertainties for cMSSM and simplified model



CMSSM $\tan\beta=10$, CMS Simulation $\sqrt{s} = 7$ TeV, ≥ 1 b-tag





$pp \rightarrow \tilde{g}\tilde{g}, \tilde{g} \rightarrow t\bar{t}\tilde{\chi}^0$; $m(\tilde{q}) \gg m(\tilde{g})$



$pp \rightarrow \tilde{g}\tilde{g}, \tilde{g} \rightarrow t\bar{t}\tilde{\chi}^0$; $m(\tilde{q}) \gg m(\tilde{g})$










⌚ | Searches for SUSY in events with 3rd generation particles at CMS



- It is the complete MET model for an HT interval/bin.



$$\mathcal{M}_i(x) \sim x \exp(-\alpha x^{0.5}) \times (1 + \text{erf}(x; b_0 + b_1 H_{T,i}, c_0 + c_1 H_{T,i})) \times (1 - \text{erf}(x; b_0 + b_1 H_{T,i+1}, c_0 + c_1 H_{T,i+1})).$$






Source	μ channel			e channel		
	total	0-tag	≥ 1-tag	total	0-tag	≥ 1-tag
Jet and E_T^{miss} scale	6.0 %	7.5 %	7.2 %	3.1 %	5.6 %	2.1 %
W polarization (1), ±10%	0.5 %	0.6 %	0.1 %	1.3 %	1.8 %	0.2 %
W ⁻ polarization (2), ±5%	0.3 %	0.5 %	0.1 %	0.5 %	0.5 %	0.2 %
W ⁺ polarization (2), ±5%	0.1 %	0.2 %	0.1 %	0.1 %	0.1 %	0.1 %
W polarization (3), ± 10%	0.0 %	0.1 %	0.0 %	0.5 %	0.6 %	0.2 %
vary lep. eff. at low p_T	0.4 %	0.3 %	0.6 %	0.6 %	1.3 %	0.7 %
vary lep. eff. in endcaps	0.2 %	0.2 %	0.1 %	0.6 %	0.8 %	0.4 %
vary pile-up	0.1 %	0.1 %	0.2 %	0.3 %	1.5 %	0.4 %
Non-leading bkg ± 50%	0.7 %	0.4 %	0.4 %	4.0 %	3.0 %	6.2 %
dilep. contr ± 50%	0.1 %	0.5 %	0.7 %	0.6 %	1.2 %	0.6 %
$\sigma(t\bar{t})$, ± 32%	1.2 %	2.3 %	1.6 %	0.7 %	1.8 %	2.0 %
$\sigma(W+\text{jets})$, ± 32%	1.3 %	2.9 %	2.3 %	2.6 %	1.6 %	2.8 %
exponent $t\bar{t}$ ± 10%	1.6 %	0.2 %	5.3 %	1.8 %	0.3 %	4.8 %
exponent W ⁺ +jets ± 10%	3.5 %	4.4 %	1.3 %	3.6 %	4.6 %	1.5 %
exponent W ⁻ +jets ± 10%	0.7 %	0.8 %	0.3 %	0.9 %	1.4 %	0.9 %
α slope $t\bar{t}$	11.0 %	2.4 %	29.3 %	14.8 %	5.0 %	34.3 %
α slope W ⁺ +jets	15.9 %	20.6 %	6.0 %	16.5 %	22.2 %	5.1 %
α slope W ⁻ +jets	4.9 %	8.2 %	2.0 %	5.6 %	8.7 %	0.5 %
Variation of Erfc.	4.1 %	4.6 %	2.9 %	3.1 %	3.2 %	2.7 %

Source	μ channel					e channel				
	total	0-tag	1-tag	≥ 1-tag	≥ 2-tag	total	0-tag	1-tag	≥ 1-tag	≥ 2-tag
W+jets/ $t\bar{t}$ ratio	2.9 %	2.1 %	6.1 %	4.8 %	2.4 %	1.1 %	2.4 %	2.6 %	2.3 %	2.3 %
b-tagging efficiency ±1 σ	2.0 %	1.5 %	2.2 %	1.3 %	5.1 %	2.2 %	1.6 %	0.8 %	1.7 %	3.6 %
mistag rate ±1 σ	0.4 %	0.4 %	0.7 %	0.9 %	0.6 %	0.3 %	0.4 %	0.4 %	0.2 %	0.1 %





$$\Delta\phi_N^{\min} \equiv \min(\Delta\phi_i / \sigma_{\Delta\phi,i})$$

$$\sigma_{T_i}^2 \approx \sum_n (\sigma_{p_{T,n}} \sin \alpha_n)^2$$




$$\sigma_{\Delta\phi,i} = \arctan(\sigma_{T_i} / E_T^{\text{miss}})$$

SB=Side Band, LSB=Low Side Band
LDP=Low Delta Phi, SIG=Signal

QCD background

$$N_{SIG}^{QCD} = \frac{N_{LSB}}{N_{LSB-LDP}} \times (N_{SIG-LDP} - N_{SIG-LDP}^{top,MC} - N_{SIG-LDP}^{EW,MC}),$$

$$N_{SB}^{QCD} = \frac{N_{LSB}}{N_{LSB-LDP}} \times (N_{SB-LDP} - N_{SB-LDP}^{top,MC} - N_{SB-LDP}^{EW,MC}),$$




cakir@cern.ch



Top and W+jets background (nominal)

$$N_{SIG}^{top+W} = \frac{N_{SIG-SL}}{N_{SB-SL}} \times (N_{SB} - N_{SB}^{Z \rightarrow \nu\bar{\nu}} - N_{SB}^{QCD} - N_{SB}^{other, MC}).$$



Z +jets background $Z \rightarrow \mu\mu$ -to- $\nu\nu$ replacement




$$BR(Z \rightarrow \nu\bar{\nu}) / BR(Z \rightarrow \mu^+\mu^-) = 5.95 \pm 0.02$$


$$\epsilon = \mathcal{A} \cdot \epsilon_{trig} \cdot \epsilon_{\ell reco}^2 \cdot \epsilon_{\ell sel}^2$$

	1BL	1BT	2BL	2BT	3B
Scale factors	20 (17)	20 (17)	61 (49)	61 (49)	105 (144)
$Z \rightarrow \ell^+\ell^-$ background	8 (10)	8 (10)	8 (10)	8 (10)	8 (10)
Acceptance	3 (3)	8 (6)	3 (3)	4 (4)	3 (3)
Lepton selection efficiency	4 (5)	4 (5)	4 (5)	4 (5)	4 (5)
Trigger efficiency	5 (5)	5 (5)	5 (5)	5 (5)	5 (5)
MC closure	19 (11)	19 (11)	19 (11)	19 (11)	19 (11)
Total	30 (24)	30 (25)	65 (52)	65 (52)	107 (145)



Top and W+jets background (E_T^{miss} -reweighting)

1. top or W+jets events in which exactly one W decays into an e or μ , or into a τ that decays into an e or μ , while the other W (if any) decays hadronically;
2. top or W+jets events in which exactly one W decays into a hadronically-decaying τ , while the other W (if any) decays hadronically;
3. $t\bar{t}$ events in which both W bosons decay into an e, μ or τ , with the τ decaying either leptonically or hadronically.





Signal region		H_T (GeV)	E_T^{miss} (GeV)	N_{bjets}
1b-loose	1BL	> 400	> 250	≥ 1
1b-tight	1BT	> 500	> 500	≥ 1
2b-loose	2BL	> 400	> 250	≥ 2
2b-tight	2BT	> 600	> 300	≥ 2
3b	3B	> 400	> 250	≥ 3



PAS-SUS-12-004 (needs to be improved – later!)

Background Estimate for Single Tau Analysis

Several energy corrections will be applied before the computation of HT and MHT.




$$f_{event}^{corr} = \frac{p_{\mu}^W \times \epsilon_{\tau}^{ID} \times f_{\tau}^{bf(hadr)}}{\epsilon_{\mu}^{reco} \times \epsilon_{\mu}^{iso}}$$

- Muon reconstruction efficiency
- Muon isolation efficiency
- Muons produced in tau decays
- Tau reconstruction efficiency
- Tau hadronic branching fraction

Estimate of Fake-Tau Background

The fake rates f_i of jet i are used as individual events weights in the following form:

$$f_{event}^{corr} = 1 - \prod_i (1 - f_i), \quad f_i = f_i(p_T^i, \eta^i)$$



Simplified Model for multi tau states

PAS-SUS-12-004 (needs to be improved – later!)

Background Estimate for Di-Tau Analysis

We measure correction factors and/or selection efficiencies in control regions (CRs) and use these values to extrapolate to the region where we expect to observe our signal. A novel approach is to use the observed jet multiplicity in each CR along with measured $\text{jet} \rightarrow \tau h$ misidentification rate to calculate the yield in the signal regions (SR).

$$N_{Background}^{SR} = N_{Background}^{CR} [\alpha_{\tau\tau} \mathcal{P}(0) + \alpha_{\tau j} \mathcal{P}(1) + \alpha_{jj} \mathcal{P}(2)]$$

tt estimation

$$N_{t\bar{t}}^{Signal} = A_{\tau+j} \frac{N_{t\bar{t}}^{CR}}{P(2 \text{ b-jets})} \epsilon^{\tau \text{ iso}} \sum_{N=1}^{\infty} P(N) \sum_{n=1}^N C(N, n) f^n (1-f)^{N-n} \\ + A_{j+j} \frac{N_{t\bar{t}}^{CR}}{P(2 \text{ b-jets})} \sum_{M=2}^{\infty} P(M) \sum_{m=2}^M C(M, m) f^m (1-f)^{M-m}$$

$Z \rightarrow \nu\nu$ estimation

$$N_{Z \rightarrow \nu\nu + \text{jets}}^{Signal} = \frac{N_{Z \rightarrow \mu\mu + \text{jets}}^{CR}}{A_{\mu}^2 \epsilon_{\mu}^2} \frac{B(Z \rightarrow \nu\nu)}{B(Z \rightarrow \mu\mu)} \frac{\epsilon_{H_T}^{Trigger}}{\epsilon_{\mu\tau}^{Trigger}} \epsilon^{H_T} \sum_{N=2}^{\infty} P(N) \sum_{n=2}^N C(N, n) f^n (1-f)^{N-n}$$

$Z \rightarrow \tau\tau$ estimation

$$N_{Z \rightarrow \tau\tau}^{Signal} = N_{Z \rightarrow \mu\mu}^{CR} R \epsilon^{H_T} \frac{A_{\tau}^2 \epsilon_{\tau}^2}{A_{\mu}^2 \epsilon_{\mu}^2} \\ + N_{Z \rightarrow \mu\mu}^{CR} R \epsilon^{H_T} \frac{A_{\tau}^2 (2\epsilon_{\tau}(1-\epsilon_{\tau}))}{A_{\mu}^2 \epsilon_{\mu}^2} \sum_{N=1}^{\infty} P(N) \sum_{n=1}^N C(N, n) f^n (1-f)^{N-n} \\ + N_{Z \rightarrow \mu\mu}^{CR} R \epsilon^{H_T} \frac{2A_{\tau}(1-A_{\tau})\epsilon_{\tau}}{A_{\mu}^2 \epsilon_{\mu}^2} \sum_{M=1}^{\infty} P(M) \sum_{m=1}^M C(M, m) f^m (1-f)^{M-m} \\ + N_{Z \rightarrow \mu\mu}^{CR} R \epsilon^{H_T} \frac{(1-A_{\tau})^2}{A_{\mu}^2 \epsilon_{\mu}^2} \sum_{K=2}^{\infty} P(K) \sum_{k=2}^K C(K, k) f^k (1-f)^{K-k}$$

W+Jets estimation

$$N_{W+\text{jets}}^{Signal} = A_{\tau+j} \frac{N_{W+\text{jets}}^{\text{After subtraction}}}{P(0 \text{ b-jets})} \epsilon^{\tau \text{ iso}} \sum_{N=1}^{\infty} P(N) \sum_{n=1}^N C(N, n) f^n (1-f)^{N-n} \\ + A_{j+j} \frac{N_{W+\text{jets}}^{\text{After subtraction}}}{P(0 \text{ b-jets})} \sum_{M=2}^{\infty} P(M) \sum_{m=2}^M C(M, m) f^m (1-f)^{M-m}$$




cakir@cern.ch



$$R = \frac{B(Z \rightarrow \tau\tau) B(\tau \rightarrow \tau h)}{B(Z \rightarrow \mu\mu)} \frac{\epsilon_{H_T}^{Trig}}{\epsilon_{\mu\tau}^{Trig}}$$

Altan CAKIR | Searches for SUSY in events with 3rd generation particles at CMS


ICHEP 2012, Melbourne, Australia | Page 32









	SR0	SR1	SR2	SR3	SR4	SR5	SR6	SR7	SR8
No. of jets	≥ 2	≥ 2	≥ 2	≥ 2	≥ 2	≥ 2	≥ 2	≥ 3	≥ 2
No. of btags	≥ 2	≥ 2	≥ 2	≥ 2	≥ 2	≥ 2	≥ 2	≥ 3	≥ 2
Lepton charges	$++/--$	$++/--$	$++$	$++/--$	$++/--$	$++/--$	$++/--$	$++/--$	$++/--$
E_T^{miss}	$> 0 \text{ GeV}$	$> 30 \text{ GeV}$	$> 30 \text{ GeV}$	$> 120 \text{ GeV}$	$> 50 \text{ GeV}$	$> 50 \text{ GeV}$	$> 120 \text{ GeV}$	$> 50 \text{ GeV}$	$> 0 \text{ GeV}$
H_T	$> 80 \text{ GeV}$	$> 80 \text{ GeV}$	$> 80 \text{ GeV}$	$> 200 \text{ GeV}$	$> 200 \text{ GeV}$	$> 320 \text{ GeV}$	$> 320 \text{ GeV}$	$> 200 \text{ GeV}$	$> 320 \text{ GeV}$
Charge-flip BG	1.32 ± 0.28	1.04 ± 0.22	0.52 ± 0.11	0.05 ± 0.01	0.35 ± 0.08	0.11 ± 0.03	0.02 ± 0.01	0.01 ± 0.01	0.18 ± 0.05
Fake BG	5.89 ± 3.78	4.46 ± 2.68	1.86 ± 1.12	0.33 ± 0.36	2.46 ± 2.16	0.77 ± 0.82	0.20 ± 0.33	0.08 ± 0.52	1.36 ± 1.12
Rare SM BG	4.92 ± 2.57	4.44 ± 2.32	2.95 ± 1.59	1.01 ± 0.62	2.95 ± 1.56	1.77 ± 1.03	0.71 ± 0.51	0.24 ± 0.40	2.24 ± 1.27
Total BG	12.13 ± 4.58	9.94 ± 3.55	5.33 ± 1.95	1.39 ± 0.72	5.76 ± 2.67	2.64 ± 1.32	0.93 ± 0.61	0.33 ± 0.66	3.78 ± 1.69
Event yield	13	11	0	1	4	2	1	1	4
N_{UL} (13% unc.)	11.4	10.0	5.1	4.1	6.6	5.1	3.8	3.4	6.2
N_{UL} (20% unc.)	11.7	10.4	5.3	4.2	6.7	5.2	3.9	3.6	6.4
N_{UL} (30% unc.)	12.5	11.1	5.6	4.4	7.0	5.4	4.1	3.8	6.8



Events are collected with dilepton triggers, with one lepton of $P_T > 17 \text{ GeV}$ and the second lepton of $P_T > 8 \text{ GeV}$. The trigger efficiency is measured to be $95 \pm 3\%$ for ee events, $92 \pm 3\%$ for $e\mu$ events, and $88 \pm 3\%$ for $\mu\mu$ events from studies of events collected with hadronic triggers.



- Lepton Selection Efficiency for LM9 benchmark point: 1450, 175, 0, 50, +, m_0 , $m_{1/2}$, A_0 , $\tan\beta$ and μ parameter, respectively