

5th International workshop on heavy quark production in heavy-ion collisions

Wednesday 14 November 2012 - Saturday 17 November 2012

Academic Building, Utrecht University



Book of Abstracts

Contents

Welcome	1
Coherence effects and broadening in medium-induced QCD radiation off a massive $q\bar{q}$ antenna	1
Effects of momentum dependent drag and equation of state on the heavy flavour suppression	1
b-Jet tagging in proton-(anti)proton collisions	2
Recent developments in lattice studies for quarkonia	2
Suppression and elliptic flow of open charmed mesons	2
Bottomonium Suppression in Heavy Ion Collisions	3
Recent results on heavy flavour production from LHCb	3
The STAR Heavy Flavor Upgrades	3
Radiative Contributions to Heavy Quark Energy Loss in a Langevin Approach	4
Precision quarkonium studies using multi-purpose fixed-target experiments	4
Measurement of the J/ψ elliptic flow at forward rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ALICE experiment	5
Quarkonium production measurements in Pb-Pb collisions with the ALICE experiment at the LHC	5
Exclusive coherent production of heavy vector mesons in nucleus-nucleus collisions	6
Production of two $c\bar{c}$ pairs and two identical D mesons - evidence for double parton scattering mechanism.	6
Radiative energy loss and damping effects	8
Open charmed mesons at LHC	8
Measurement of heavy-flavour decay muon production at forward rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with ALICE	8
D meson nuclear modification factors and D meson elliptic flow in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ALICE detector	9

Feasibility of the detection of D0 mesons in the NA61/SHINE experiment at the CERN SPS	9
Open and hidden charm dynamics	10
The chiral transition and equation of state	10
Heavy flavour measurements with ALICE in view of the inner tracker upgrade	10
b-jet Identification in PbPb Collisions with CMS	11
Measurement of the nuclear modification factor and the elliptic flow of electrons from heavy flavour decays in Pb-Pb collisions at 2.76 TeV with ALICE	11
Open charmed mesons production at STAR	12
Heavy quark quenching from RHIC to LHC	12
J/ψ production in STAR	13
Coherent J/ψ photoproduction in ultra-peripheral Pb-Pb collisions at ALICE	14
Bottom electron production in 7 TeV pp collisions at ALICE	14
Heavy flavor spectra in AA collision within a Langevin approach	15
System Size, Energy and Rapidity Dependence of Quarkonia Production Measured by the PHENIX Detector.	15
Measurement of charm and bottom contributions to electrons from heavy quark decay at RHIC-PHENIX experiment	16
Heavy flavour correlation measurements in pp and Pb-Pb collisions with ALICE	16
Cold Nuclear Matter Effects on Heavy Quarks from PHENIX	17
Measurements of Non-photonic Electron Production and Azimuthal Anisotropy in Au+Au Collisions at STAR	17
D meson azimuthal anisotropy measured with ALICE experiment	17
Upsilon PbPb from Alice	18
CMS	18
Upsilon production in STAR	18
The CBM heavy-quark program	18
Energy loss effects in cold nuclear matter	19
Theory overview	19
ATLAS	19
Heavy flavour particle correlations in Phenix	19
QCD factorization theorems for production of heavy quarkonia	19

AdS/CFT	19
PHENIX upgrades	20
Heavy flavour perspectives with CBM	20
Theory summary	20
Experiment summary	20
Recent results on quarkonium and open bottom suppression in PbPb collisions at the LHC with CMS	20
Nonperturbative Heavy-Flavor Diffusion and Hadronization in a Hydrodynamic Description of Heavy-Ion Collisions	21
Heavy-quarkonium suppression in p-A collisions from parton energy loss	21
Welcome, Head of the Department of Physics and Astronomy	22
Heavy Quark Production in Pb-Pb Collisions at the LHC with the ATLAS Detector	22
Gauge/gravity, thermalization and energy loss	22
Quarkonium at finite temperature	22
Future of Heavy Quark Measurement at sPHENIX	23
Heavy Flavor Dileptons at PHENIX	23
From the RAA to azimuthal correlations: what can we learn from heavy-quark observables?	24
Upsilon measurements in STAR	24

Welcome - Board: LOC / 0

Welcome

Corresponding Author: a.mischke@uu.nl

Welcome and announcements

HQ prod in jets and b-tag / 5

Coherence effects and broadening in medium-induced QCD radiation off a massive $q\bar{q}$ antenna

Author: Hao Ma^{None}

Co-authors: Carlos Albert Salgado Lopez ¹; Konrad Tywoniuk ²; Nestor Armesto Perez ¹; Yacine Mehtar-Tani

¹ *Universidade de Santiago de Compostela (ES)*

² *Universidade de Santiago de Compostela*

Corresponding Author: hao.ma@usc.es

Studies of medium-induced QCD radiation usually rely on the calculation of single-gluon radiation spectrum off an energetic parton traversing an extended colored medium. Recently, the importance of interference effects between emitters in the medium has been explored. In this work we extend previous studies by calculating the single-gluon coherent spectrum off an antenna consisting of a massive quark-antiquark pair. Interferences dominate the spectrum of soft gluons, which are mainly emitted outside of the cone made by the antenna opening angle, while the antenna results in a superposition of independent emitters above a critical gluon energy scale. We study the interplay between the dead-cone effect and medium-induced jet broadening in both cases of soft and hard gluons and present results on energy loss distributions.

Lattice QCD / 6

Effects of momentum dependent drag and equation of state on the heavy flavour suppression

Author: Surasree Mazumder¹

Co-authors: Jan-e Alam ²; Trambak Bhattacharyya ³

¹ *Variable Energy Cyclotron Centre, Kolkata*

² *Variable Energy Cyclotron Centre, Kolkata*

³ *Variable Energy Cyclotron Centre*

Corresponding Author: surasree@vecc.gov.in

The momentum dependence of the drag and diffusion coefficients of heavy quarks propagating through a quark gluon plasma (QGP) have been evaluated. The effects of the equation of state and momentum dependence of transport coefficients on the suppression of heavy flavours in QGP have been studied. We observe that the momentum dependence of the transport coefficients plays crucial role in the suppression of the heavy quarks and, consequently, in discerning the properties of the QGP using heavy flavors as a probe. We use the experimental data on nuclear suppression of heavy flavours and hadronic multiplicity from RHIC and LHC to constrain the initial condition of the QGP

expected to be formed at nuclear collision at RHIC energies.

Again, we have calculated drag and diffusion coefficients using Hard Thermal Loop (HTL) approximations. In this case we have shielded the infra-red divergence by the effective gluon propagator dictated by HTL perturbation theory.

Summary:

We have found out quantitatively that radiative energy loss plays more dominant role in the nuclear modification factor than the collisional loss.

We have obtained the value of the initial entropy density, which varies from 20 to 59/fm³ depending on the value of the velocity of sound suggested by lattice QCD. Our investigation leads to a conservative value of the initial entropy density of ~20/fm³ with a corresponding initial temperature of ~210 MeV, which is well above the value of the transition temperature predicted by lattice QCD.

As for the HTL approximated transport coefficients, we have found that there are considerable differences between drag and diffusion obtained using Debye mass as the infra-red shield and that calculated using HTL approximated gluon self energy as the shield.

HQ prod in jets and b-tag / 8

b-Jet tagging in proton-(anti)proton collisions

Author: Frank Filthaut¹

¹ *Radboud University Nijmegen*

Corresponding Author: frank.filthaut@cern.ch

A large part of the high-pt physics programme at the Tevatron and LHC colliders relies on the identification of jets originating from the fragmentation of b quarks. In my presentation I will discuss the salient features of the b-jet tagging algorithms developed to achieve this goal, and also discuss the methods that have been developed to calibrate the performance of these algorithms.

Lattice QCD / 9

Recent developments in lattice studies for quarkonia

Author: Olaf Kaczmarek¹

¹ *University of Bielefeld*

Corresponding Author: okacz@physik.uni-bielefeld.de

I will give a review on recent developments in lattice QCD studies for quarkonia in the quark gluon plasma medium. I will discuss recent progress in the extraction of spectral properties from lattice QCD calculations of hadronic correlation functions. Besides medium modifications of bound states and their dissociation in the plasma I will focus on transport coefficients, like heavy quark diffusion constants, extracted from different correlation functions on the lattice. Present limitations and future perspectives for studies of quarkonia and related transport coefficients on the lattice will be discussed.

Open charm / 10

Suppression and elliptic flow of open charmed mesons

Author: Jan Uphoff¹

¹ *Goethe University Frankfurt*

Corresponding Author: uphoff@th.physik.uni-frankfurt.de

For the first time in ultra-relativistic heavy ion collisions it is possible at the LHC and also at RHIC to measure the suppression and flow of D mesons directly without any contribution from B mesons, giving direct and unbiased insight to the charm quark interactions with the quark gluon plasma. Model calculations and predictions for the nuclear modification factor and elliptic flow of D mesons in ultra-relativistic heavy ion collisions are reviewed and compared to experimental data. Furthermore, the D meson results are put into context with other heavy flavor observables.

Quarkonia / 11

Bottomonium Suppression in Heavy Ion Collisions

Author: Michael Strickland¹

¹ *Gettysburg College*

Corresponding Author: mstrickl@gettysburg.edu

The thermal suppression of heavy quark bound states represents an ideal observable for determining if one has produced a quark gluon plasma in ultrarelativistic heavy-ion collisions. In recent years, however, a paradigm shift has taken place in the theory of quarkonium suppression due to new first principles calculations of the thermal widths of these states. These thermal widths are large, eg $O(20-100 \text{ MeV})$ for the Upsilon, and cause in-medium suppression of the states at temperatures below their traditionally defined disassociation temperatures. In order to apply the newly developed understanding to phenomenology, however, one must make detailed space-time models of the plasma including the effects of finite shear viscosity. These effects include not only the modification of the time evolution of the temperature of the system, flow, etc., but also the emergence of potentially large local momentum-space anisotropies which can affect the in-medium properties of the states. I will present comparisons of theory with data from RHIC 200 GeV/nucleon Au-Au collisions and LHC 2.76 TeV/nucleon Pb-Pb collisions as a function of number of participants, rapidity, and transverse momentum.

Summary / 12

Recent results on heavy flavour production from LHCb

Authors: Nicholas BROOK¹; Nicholas Brook²

¹ *BRISTOL*

² *University of Bristol (GB)*

Corresponding Authors: n.brook@bristol.ac.uk, nick.brook@cern.ch

During 2011, the LHCb experiment accumulated $\sim 1 \text{ fb}^{-1}$ of data in proton-proton collisions data at 7 TeV, collecting a sample rich in B mesons and baryons. The data provide a wealth of new measurements as well as probes of QCD theory predictions. We present recent results in quarkonium and b and c hadron production, as well as studies of these states' properties such as masses and decay asymmetries.

Future experiments and upgrades / 13

The STAR Heavy Flavor Upgrades

Author: Flemming Videbaek¹

¹ *BNL*

Corresponding Author: videbaek@bnl.gov

The STAR have already reported on results in pp and AA on heavy flavor production. In order to perform high precision measurements of heavy flavor STAR has embarked on an upgrade program. The primary motivation for the Heavy Flavor Tracker (HFT) is perform measurements of heavy quark collectivity and to separate bottom and charm quark energy loss in the QCD medium. This will be achieved by the HFT by extending the capabilities of STAR by measurements of displaced vertices and direct topological identification of open charm. The motivation for the Muon Telescope Detector (MTD) is to provide muon identification at mid-rapidity for measurements of J/psi and Upsilon.

These detectors will greatly enhance the STAR physics capability to measure heavy quark collectivity and correlations using topologically reconstructed charmed hadrons and heavy quark decay electron-muon correlations and bottom quark production. In addition, measurements of the quarkonium muon decay channels will enable us to separate Upsilon 1S from 2S and 3S states in p+p and A+A collisions. An overview of the upgrades, their expected performance and current status will also be presented. The STAR Collaboration should complete the Heavy Flavor Tracker (HFT) and the Muon Telescope Detector (MTD) upgrades by 2014.

Open charm / 14

Radiative Contributions to Heavy Quark Energy Loss in a Langevin Approach

Author: Steffen A. Bass¹

Co-author: Shanshan Cao¹

¹ *Duke University*

Corresponding Author: bass@phy.duke.edu

Over the past year our group has studied heavy quark energy loss in the context of a Langevin approach. Main emphasis of our study was the question of heavy quark thermalization in a QGP medium as well as the dependence of the most common observables associated with heavy quark energy loss (R_{AA} and v_2) on various parameters of the medium evolution, to which our Langevin approach is coupled.

Here, in addition to reviewing the results of the aforementioned studies, we will introduce an algorithm that incorporates the radiative energy loss of heavy quarks in a QGP medium into the current Langevin framework by treating gluon radiation as an additional force to the heavy quarks traversing the medium. We examine the corresponding effects on observables such as the elliptic flow and the nuclear modification factor, and compare our simulations to the experimental results at the LHC. Our study constitutes an essential contribution to a more quantitative understanding of the energy loss of heavy quarks propagating through hot and dense nuclear medium.

New ideas from theory / 15

Precision quarkonium studies using multi-purpose fixed-target experiments

Author: Jean-Philippe Lansberg¹

¹ *IPN Orsay, Paris Sud U. / IN2P3-CNRS*

Corresponding Author: lansberg@in2p3.fr

I will briefly review the most important quarkonium-production results at fixed-target experiments (NA50, NA60, Hera-B, E866, ...) reported in the last decade. Then I will move onto the discussion of the many quarkonium-physics opportunities offered by A multi-purpose Fixed-Target Experiment (AFTER) using the proton and lead Large Hadron Collider (LHC) beams extracted by a bent crystal. With a foreseen integrated luminosity of 0.5 fb⁻¹ per year on a typical 1 cm-long target, AFTER would provide a quarkonium observatory, where their production can be analysed in great details in pp, pd and pA collisions at $\sqrt{s_{NN}} = 115$ GeV and at $\sqrt{s_{NN}} = 72$ GeV in PbA collisions. By instrumenting the target-rapidity region, the large negative-xF domain can be accessed for the first time, greatly extending previous measurements by Hera-B and E866. Such analyses should help resolving the quarkonium-production controversies and clear the way for gluon PDF extraction via quarkonium studies. The nuclear target-species versatility provides a unique opportunity to study nuclear matter and the features of the hot and dense matter formed in PbA collisions.

Quarkonia / 16

Measurement of the J/ψ elliptic flow at forward rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ALICE experiment

Author: Indranil Das¹

Co-author: Collaboration ALICE²

¹ *Universite de Paris-Sud 11 (FR)*

² .

Corresponding Author: indranil.das@cern.ch

Heavy quark resonances, produced in high energy heavy-ion collisions, are important observables for the study of Quantum Chromodynamics (QCD) interactions at extreme energy-densities. Lattice QCD calculations predict a phase transition of the nuclear matter to Quark Gluon Plasma (QGP). The suppression (due to color screening mechanism) or enhancement (due to regeneration mechanism) of charmonium resonances (J/ψ) have been proposed as a signature of QGP. ALICE has measured the J/ψ in the forward rapidity region via the dimuon ($\mu^+\mu^-$) decay channel in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, the result shows less suppression with respect to that measured at lower energies of SPS and RHIC. The elliptic flow measurement of J/ψ provides an important observation to test the degree of thermalization of J/ψ in heavy-ion collisions.

In this talk, the elliptic flow of J/ψ measured in the forward rapidity ($-4.0 < y < -2.5$) of ALICE via $\mu^+\mu^-$ decay channel will be discussed for Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. The recent analysis of the p_T integrated flow for different centrality bin will be presented. We will also discuss differential flow as function of transverse momentum. The results will be compared with the previous measurements at RHIC and theoretical calculations.

Quarkonia / 17

Quarkonium production measurements in Pb-Pb collisions with the ALICE experiment at the LHC

Author: Lizardo Valencia Palomo¹

¹ *Universite de Paris-Sud 11 (FR)*

Corresponding Author: lizardo.valencia.palomo@cern.ch

ALICE (A Large Ion Collider Experiment), one of the four main experiments at the Large Hadron Collider (LHC), was designed and built to perform dedicated research on heavy ion collisions to study the Quark Gluon Plasma (QGP), a deconfined state of strongly interacting QCD matter.

As heavy flavours are produced on a very short time-scale in the initial hard scattering processes, they can be used to characterize the hot and dense medium formed in high-energy heavy-ion collisions through their modified yield as compared to pp collisions.

In ALICE, quarkonium production is measured via the dielectron and dimuon decay channels at central ($|y| < 0.9$) and forward ($2.5 < y < 4$) rapidity, respectively.

In this talk the latest results on quarkonium production by the ALICE experiment in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV will be presented.

HF particle correlations / 18

Exclusive coherent production of heavy vector mesons in nucleus-nucleus collisions

Author: Wolfgang Schaefer¹

Co-authors: Anna Cisek¹; Antoni Szczurek¹

¹ *Institute of Nuclear Physics PAN*

Corresponding Author: wolfgang.schafer@ifj.edu.pl

High energy heavy ions are the source of a flux of Weizsäcker-Williams photons, which can be utilized to study the photoproduction of vector mesons on nuclear targets.

We discuss how information on the small- x gluon distribution in the nucleus can be obtained.

We present our calculations based on a k_{\perp} -factorization approach which allows us to construct the unintegrated gluon of a nucleus from the free-nucleon one.

Saturation effects are incorporated by an explicit treatment of the $q\bar{q}g$ -Fock state. If time permits, we also briefly discuss the role of the latter in incoherent diffraction on nuclei.

The presentation will be based on our recent publication

“Exclusive coherent production of heavy vector mesons in nucleus-nucleus collisions at LHC.”
A. Cisek, W. Schaefer (Cracow, INP), A. Szczurek (Cracow, INP & Rzeszow U.), Phys.Rev. C86 (2012) 014905.

HF decays leptons / 19

Production of two $c\bar{c}$ pairs and two identical D mesons - evidence for double parton scattering mechanism.

Author: Antoni Szczurek¹¹ I

Corresponding Author: antoni.szczurek@ifj.edu.pl

We discuss charm production at LHC. The production of single $c\bar{c}$ pairs is calculated in the k_t -factorization approach. We use several unintegrated gluon distributions from the literature. Some of them include effect of small- x saturation and fullfil Balitsky-Kovchegov evolution equation. The hadronization is included with the help of fragmentation functions found for the production of c (\bar{c}) in e^+e^- collisions. Differential distributions for several charmed mesons will be presented and compared to recent results of the ALICE and LHCb collaborations. Some missing strength is identified. Different schemes of fragmentation are discussed.

We concentrate on production of two pairs of $c\bar{c}$ within a simple formalism of double-parton scattering (DPS). We perform calculation both in collinear and k_t -factorization approaches. Surprisingly large cross sections, comparable to single-parton scattering (SPS) contribution, are predicted for LHC energies.

Both total inclusive cross section as a function of energy and differential distributions are shown.

We include recently discussed evolution of double partons in the case of two scales.

We discuss perspectives how to identify the double scattering contribution. We find much larger cross section for large rapidity distance between charm quarks from different hard parton scatterings compared to single scattering.

Predictions for two $c\bar{c}$ pair production in single-parton scattering will be presented.

We present also first results for the $c\bar{c}c\bar{c}$ production in the k_t -factorization approach. The results are compared with those for the collinear approach.

Predictions for the production of different pairs of charm mesons ($D^0\bar{D}^0$, $D^0\bar{D}^0$, etc.) are presented for the kinematics of ATLAS, CMS, ALICE and LHCb experiment. Differential distributions of single meson provided that both mesons are measured by the detector, distribution in azimuthal angle between two mesons and in dimeson invariant mass are calculated and will be compared with recent experimental data. Predictions for future measurements are given.

LITERATURE:

\item M. {\L}uszczak, R. Maciu{\L}a and A. Szczurek, "Production of two $c\bar{c}$ pairs in double-parton scattering", Phys. Rev. {\bf D85} (2012) 094034.

\item W. Sch{\L}afer and A. Szczurek, "Production of two $c\bar{c}$ pairs in gluon-gluon scattering in high energy proton-proton scattering", Phys. Rev. {\bf D85} (2012) 094029.

\item R. Maciu{\L}a and A. Szczurek, a paper in preparation.

HF decays leptons / 20**Radiative energy loss and damping effects****Author:** Thierry Gousset¹¹ *Subatech***Corresponding Author:** gousset@subatech.in2p3.fr

The effect of the damping of radiation in an absorptive medium on radiative energy loss is considered. First, the radiation spectrum for a fast parton going through a dense medium is estimated from considerations on the formation time in the QCD context. In a second stage, such a derivation is connected to a full calculation carried out in electrodynamics.

Open charm / 21**Open charmed mesons at LHC****Author:** Rafal Maciula¹**Co-author:** Antoni Szczurek²¹ *Institute of Nuclear Physics PAN*² *I***Corresponding Author:** rafal.maciula@ifj.edu.pl

I will discuss production of open charmed mesons in proton-proton collisions at the LHC. The cross section for inclusive production of $c\bar{c}$ pairs is calculated in the framework of the k_{\perp} -factorization approach. Taking wide range of x values necessary for the calculation we use and test several unintegrated gluon distributions from the literature. Some of them include effect of small- x saturation and fullfil Balitsky-Kovchegov evolution equation. Theoretical uncertainties of the model related to the choice of renormalization and factorization scales as well as due to the quark mass are also discussed. Results from the k_{\perp} -factorization approach are compared to NLO parton model and FONLL predictions. The hadronization of charm quarks is included with the help of different fragmentation functions found for the production of charm in e^+e^- collisions. Sensitivity of our predictions to the choice of the model of fragmentation is also shown. Inclusive differential distributions in transverse momentum and (pseudo)rapidity of several charmed mesons (D^0 , D^{\pm} , $D^{*\pm}$, D_S^{\pm}) will be presented and compared to recent results of the ALICE, ATLAS and LHCb collaborations. Furthermore, I will also consider production of different pairs of charmed mesons ($D^0\bar{D}^0$, D^0D^- , $D^+D_S^-$ etc.) in the unique kinematics of forward rapidities in the LHCb experiment. Kinematical correlations in azimuthal angle $\varphi_{D\bar{D}}$ and invariant mass $M_{D\bar{D}}$ distributions will be presented and compared to LHCb data.

HF decays leptons / 22**Measurement of heavy-flavour decay muon production at forward rapidity in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with ALICE****Author:** Philippe Pillot¹¹ *Laboratoire de Physique Subatomique et des Technologies Associees*

Corresponding Author: pillot@subatech.in2p3.fr

Heavy flavours (charm and beauty) are expected to provide essential information on the properties of the high-density strongly-interacting system formed in the early stages of high-energy heavy-ion collisions. In particular, the investigation of modifications of heavy-flavour transverse momentum yield will shed light on the nature of the parton-medium interactions, while the measurement of their collective flow provides insights on the possible thermalization of heavy quarks in the medium.

ALICE is the experiment at the LHC mainly dedicated to the study of nucleus-nucleus collisions. At forward rapidity ($2.5 < y < 4$), the production of open heavy flavours is measured via their semi-muonic decay channels in the Forward Muon Spectrometer. After a short description of the apparatus, the latest results on the pT-differential nuclear modification factor RAA and elliptic flow of muons from heavy-flavour decays in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV will be presented.

Open charm / 23

D meson nuclear modification factors and D meson elliptic flow in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ALICE detector

Author: Alessandro Grelli¹

¹ *University of Utrecht (NL)*

Corresponding Author: alessandro.grelli@cern.ch

The properties of the hot and dense QCD medium formed in ultra-relativistic heavy-ion collisions, as well as the mechanism of in-medium partonic energy loss, can be accessed via the measurement of the nuclear modification factor of particle production. The measurement of D meson production provides key tests of parton energy-loss models, which predict that charm quarks should experience less in-medium energy loss than light quarks and gluons. Moreover if in-medium hadronization is the dominant mechanism of charm hadron formation at low $p_{\{T\}}$ then the production of charm strange hadrons, like $D_{\{s\}}^{\{+\}}$, is expected to be largely enhanced. Finally the elliptic flow v_2 of D meson compared to that of light hadrons is expected to bring insights into the degree of thermalization of charm quarks within the quark gluon plasma. The ALICE experiment has measured the production of prompt $D^{\{0\}}$, $D^{\{+\}}$ and $D^{\{*\{+\}}$ and $D^{\{+\{s\}}$ mesons in *pp* and *Pb-Pb* collisions at the LHC at $\sqrt{s} = 7$ and 2.76 TeV and at $\sqrt{s_{NN}} = 2.76$ TeV, respectively, via the exclusive reconstruction of their hadronic decay. The $p_{\{T\}}$ -differential production yields in the range $1 < p_{\{T\}} < 36$ GeV/c ($4 < p_{\{T\}} < 12$ GeV/c in the case of $D^{\{+\{s\}}$) at central rapidity, $|y| < 0.5$, were used to calculate the nuclear modification factor. A suppression of a factor 4-5 for $p_{\{T\}} \geq 10$ GeV/c in the 7.5% most central collisions was observed. The measurement of the D meson elliptic flow in semi-central (30-50%) Pb-Pb collisions will be presented.

Cold nuclear matter / 24

Feasibility of the detection of D0 mesons in the NA61/SHINE experiment at the CERN SPS

Author: Yasir Ali¹

Co-author: Pawel Staszal¹

¹ *University of Jageollanian*

A feasibility study of D0 meson (Open charm) measurements by its decay into two daughter particles, $D0 \rightarrow K^+ \pi^-$, in central Pb+Pb collisions at the CERN SPS energies will be presented. The study for the NA61/SHINE experimental setup supplemented with a future vertex detector that will allow for a precise track and vertex reconstruction at the target proximity. In order to generate the physical input we use the AMPT (A MultiPhase Transport model) event generator with multiplicities of D0 and D0bar mesons scaled to the multiplicities predicted by the HSD model. We employed the GEANT4 application to describe particle transport through the experimental setup. The presentation will discuss obtained results focusing on the predicted experimental yields of D0 mesons and comparing different analysis strategies. It will also address the issue of the vertex detector optimization regarding its geometry and applied detection technologies.

Quarkonia / 25

Open and hidden charm dynamics

Author: Elena Bratkovskaya¹

¹ *FIAS*

Corresponding Author: elena.bratkovskaya@th.physik.uni-frankfurt.de

Based on the Parton-Hadron-String Dynamics (PHSD) transport approach we study the suppression pattern of charmonia at SPS and RHIC with respect to centrality and rapidity employing various model concepts such as variants of the comover absorption' model or the charmonium melting' scenario. A detailed comparison to the RHIC experimental data demonstrates that non-hadronic interactions are mandatory to describe the narrowing of the J/Ψ rapidity distribution from pp to central Au+Au collisions. The Ψ' to J/Ψ ratio is found to be crucial in disentangling the different charmonium absorption scenarios especially in the RHIC energy range. Also we investigate the contribution from open and hidden charm and bottom to the dilepton production from SPS to LHC energies.

Lattice QCD / 26

The chiral transition and equation of state

Author: Szabolcs Borsanyi¹

¹ *University of Wuppertal*

Corresponding Author: borsanyi@uni-wuppertal.de

I present various bulk observables from lattice QCD that characterize the chiral and deconfinement aspects of the transition at finite temperature. I discuss the efforts towards a mapping of the QCD phase diagram at small chemical potentials, and present preliminary results on the effect of the dynamical charm quark.

Future experiments and upgrades / 27

Heavy flavour measurements with ALICE in view of the inner tracker upgrade

Author: Chiara Bianchin¹

¹ *Acad. of Sciences of the Czech Rep. (CZ)*

Corresponding Author: chiara.bianchin@cern.ch

The ALICE experiment at the LHC studies Pb-Pb and pp collisions with the aim of investigating the properties of the high-density state of strongly-interacting matter, expected to be produced in Pb-Pb collisions.

Heavy quarks are sensitive probes to test the medium properties, since they are formed at shorter time scale with respect to the deconfined state.

The present measurements of the nuclear modification factor (R_{AA}) and azimuthal anisotropy of heavy flavour mesons are limited by the large combinatorial background, in the low p_T region.

The measurements of heavy-flavour baryons, interesting to assess the thermalization of heavy quarks in the medium and discriminate among thermal and coalescence models, is difficult to be achieved with the current tracking and impact parameter resolution due to the small displacement (tens of microns for the case of the Λ_c) of the decay tracks from the primary vertex.

The Upgraded Inner Tracking System will have greatly improved features in terms of: determination of the distance of closest approach (dca) to the primary vertex, standalone tracking efficiency at low p_T , momentum resolution and readout capabilities. The large benefit of the upgraded detector on the heavy-flavour physics performance will be presented in this talk.

HQ prod in jets and b-tag / 28

b-jet Identification in PbPb Collisions with CMS

Author: Matthew Nguyen¹

¹ *Ecole Polytechnique (FR)*

Corresponding Author: matthew.nguyen@cern.ch

The flavor dependence of jet quenching is a powerful handle to discriminate models of parton energy loss in heavy ion collisions. While there is evidence for a strong energy loss of heavy quarks from single particle measurements, heavy flavor tagging of fully reconstructed jets has thus far not been achieved in heavy ion collisions. In this talk we demonstrate the capacity of CMS to identify jets initiated by bottom quarks using displaced vertices reconstructed in the silicon tracking system. Identification of b-jets is shown to be feasible even in the dense environment of PbPb collisions. We discuss the status and prospects for measurements of identified b-jets in PbPb collisions with CMS.

HF decays leptons / 29

Measurement of the nuclear modification factor and the elliptic flow of electrons from heavy flavour decays in Pb-Pb collisions at 2.76 TeV with ALICE

Author: Bogdan Theodor Rascanu¹

¹ *Johann-Wolfgang-Goethe Univ. (DE)*

Corresponding Author: bogdan.theodor.rascanu@cern.ch

Theodor Rascanu for the ALICE collaboration

In heavy-ion collisions, charm and beauty quarks are produced in the initial hard scattering processes. They propagate then through the created Quark Gluon Plasma. Therefore they are a sensitive tool to probe the medium's properties.

One way to measure heavy-quarks is via electrons from the semileptonic decay channels of open charm and beauty hadrons. The suppression of heavy flavour electrons, quantified by the nuclear modification factor R_{AA} , carries information about the heavy quark energy loss mechanism in the medium.

At low transverse momentum, the level of thermalization of heavy quarks can be studied via the azimuthal anisotropy of the heavy flavour electron emission in the transverse plane, the elliptic flow v_2 . At high pt, v_2 provides insight on the path length dependence of parton energy-loss.

In this talk we present R_{AA} and v_2 measurements of electrons from heavy flavour decays at central rapidity with ALICE in central (0-10%) and semi-central (20-40%) Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, respectively. The electrons are identified using the Time Of Flight detector and Time Projection Chamber at low momenta, as well as the Electromagnetic Calorimeter at higher momenta. The latest detector also provides the possibility to trigger on electrons in the collision.

We compare the results to different theoretical models.

Open charm / 30

Open charmed mesons production at STAR

Author: Jaroslav Bielcik¹

¹ *Czech Technical University (CZ)*

Corresponding Author: jaroslav.bielcik@fifi.cvut.cz

The properties of the hot and dense nuclear matter produced at RHIC in heavy ion collisions can be investigated in multiple ways by heavy flavor production.

In this talk, we will present the STAR results of open charm hadron production at mid-rapidity in $p + p$ and Au+Au collisions at

$\sqrt{s_{NN}} = 200$ GeV. Open charm mesons were reconstructed

directly via hadronic decay channels with daughter particles

identified by TPC and TOF detectors. With abundant statistics of

Au+Au collisions collected by STAR in the year 2010 and 2011, the D-meson is measured at p_T from 0 to 8 GeV in minimum bias Au+Au

collisions. The centrality dependence of D-meson p_T spectra as well as the nuclear modification factor will be presented. The measurement of the D^0 elliptic flow in 200 GeV Au+Au collisions

will be reported. Finally, we will discuss the open charm hadron

measurement in $\sqrt{s} = 500$ GeV $p + p$ collisions to study the energy dependence of charm production.

HF decays leptons / 33

Heavy quark quenching from RHIC to LHC

Author: Pol Gossiaux¹

Co-authors: Joerg Aichelin²; Marcus Bluhm³; Marlene Nahrgang; Thierry Gousset¹

¹ *Subatech*

² *Unknown*

³ *SUBATECH Nantes*

Corresponding Author: gossiaux@subatech.in2p3.fr

Recently, we have proposed a microscopic approach for the quenching and thermalisation of heavy quarks (HQ) in URHIC \cite{Gossiaux:2008,Gossiaux:2009,Gossiaux:2010}, assuming that they interact with light partons through both elastic and radiative processes evaluated by resorting to some parameterization of the running coupling constant, while those partons are spatially distributed along hydrodynamical evolution of the hot medium. This approach is able to explain successfully several observables measured at RHIC, such as a) the nuclear modification factor and the elliptic flow of non-photonic single electrons and b) the nuclear modification factor of D mesons presented by STAR at QM 2012. We will discuss the predictions of our model for D and B mesons production in URHIC at LHC energies and confront them with experimental results obtained so far by ALICE and CMS collaborations for Pb-Pb collisions at $\sqrt{s} = 2.76$ TeV.

Since \cite{Bluhm:2011}, our interest has been triggered on the effect of an absorptive medium on radiative energy loss and more specifically on its implications on the standard LPM \cite{LPM} effect. We have advocated that the large time needed for the photon formation in Bremsstrahlung from ultrarelativistic charges is not affordable if damping is taken into account. Similar effect manifests itself in QCD, as we have recently advocated in \cite{Bluhm_2012}. It has possible implications on the quenching of D mesons in URHIC that will be discussed in this talk, focusing on observables such as nuclear modification factor and correlations.

%

\begin{thebibliography}{9}

\bibitem{Gossiaux:2008}

P.B. Gossiaux, J. Aichelin, Phys. Rev. C{\bf 78}, 014904 (2008), [hep-ph/0802.2525].

\bibitem{Gossiaux:2009}

P.B. Gossiaux, R. Bierkaendt, J. Aichelin, Physical Review C{\bf 79} (2009) 044906

\bibitem{Gossiaux:2010}

P.B. Gossiaux, V. Guiho, J. Aichelin, J. Phys. G: Nucl. Part. Phys. {\bf 37} (2010) 094019

\bibitem{Gossiaux:2011}

P.B. Gossiaux et al., arXiv:1102.1114

\bibitem{Bluhm:2011}

M. Bluhm, P.B. Gossiaux, and J. Aichelin, arXiv:1106.2856, PRL {\bf 107} (2011) 265004

\bibitem{LPM}

L.D. Landau and I. Ya. Pomeranchuk, Dokl. Akad. Nauk SSSR {\bf 92} (1953) 535; ibid. {\bf 92} (1953) 735.

\bibitem{Bluhm_2012}

M. Bluhm, P. B. Gossiaux, T. Gousset, J. Aichelin,[arXiv:1204.2469v1]

\end{thebibliography}

Quarkonia / 34

J/ψ production in STAR

Author: Barbara Trzeciak¹

¹ *Warsaw University of Technology*

Corresponding Author: barbara.trzeciak@gmail.com

Barbara Trzeciak
for the STAR Collaboration
Warsaw University of Technology / Lawrence Berkeley National Laboratory

The suppression of charmonia production in high energy nuclear collisions relative to proton-proton collisions due to the color screening was proposed as a signature of the formation of Quark-Gluon Plasma. However, there are other effects that may affect the observed charmonia production, such as cold nuclear matter effects, final state nuclear absorption, statistical coalescence of $c\bar{c}$ pairs. Studies of J/ψ production and J/ψ elliptic flow (v_2) in heavy-ion collisions can provide insight into the properties of the hot and dense medium created in relativistic heavy-ion collisions at RHIC. Moreover, measurements of the J/ψ production for different colliding systems, centralities and collision energies may help to understand J/ψ production mechanisms as well as the medium properties. The J/ψ production in $p+p$ collisions is equally important, as the J/ψ production mechanism in elementary collisions is not yet known. Of particular interest in $p+p$ collisions is the J/ψ cross-section and spin alignment (J/ψ polarization). At high- p_T , this can provide discrimination power between different J/ψ production models.

In this talk results on J/ψ production via the dielectron decay channel in Au+Au, d +Au and $p+p$ collisions at midrapidity at $\sqrt{s_{NN}} = 200$ GeV in STAR experiment in broad p_T range, $0 < p_T < 14$ GeV/ c , will be presented. The J/ψ nuclear modification factor in Au+Au and d +Au collisions as a function of centrality and p_T , and the J/ψ v_2 in Au+Au collisions will be shown. The J/ψ polarization measurement will be presented in $p+p$ collisions for J/ψ p_T from 2 to 6 GeV/ c .

Quarkonia / 35

Coherent J/ψ photoproduction in ultra-peripheral Pb-Pb collisions at ALICE

Author: Veronica Canoa Roman¹

¹ *Centro Invest. Estudios Avanz. IPN (MX)*

Corresponding Author: veronica.canoa.roman@cern.ch

The ALICE experiment at the CERN Large Hadron Collider allows the study of vector meson production in ultra-peripheral heavy-ion collisions (UPC). Due to the strong electromagnetic fields gamma-gamma and gamma-nucleus processes can be studied.

We report the first results obtained at ALICE on the coherent J/ψ photo-production cross section, a process sensitive to the gluon distribution of the nuclei, in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV and its rapidity dependence measured at both forward rapidities ($\mu+\mu-$ channel) and central rapidities ($e+e-$ and $\mu+\mu-$ channels)

HF decays leptons / 36

Bottom electron production in 7 TeV pp collisions at ALICE

Author: Tomas Aronsson¹

¹ *Yale University (US)*

Corresponding Author: tomas.aronsson@cern.ch

Tomas Aronsson for the ALICE collaboration

High-energy heavy-ion collisions at the LHC allow for the study of the properties of the quark-gluon plasma (QGP). Heavy quarks, charm and bottom, produced in the initial hard scattering processes of the collision are excellent probes of the QGP. When heavy quarks traverse the QGP they are expected to lose energy and such energy loss is predicted to be smaller than for gluons and light quarks. Heavy-flavour production can be studied using electrons from semi-leptonic decays of D and B mesons. The separation of electrons from these two sources (charm and bottom) is of crucial importance to address the expected mass dependence of energy loss. In this talk, we present the

measurement of electrons from bottom decay in the transverse momentum range of 1-11 GeV/c in 7 TeV pp collisions at the LHC. Two methods are used to identify electrons from bottom decay; a track impact-parameter study based on the displacement of the decay vertex from the collision vertex, and a two-track secondary vertex b-tagging algorithm used to select displaced decay vertices. The cross section for bottom decay electrons is compared to FONLL pQCD predictions and it will serve as a reference for studies of B meson suppression in Pb-Pb collisions.

Summary:

Bottom electron production in 7 TeV pp collisions at ALICE, Tomas Aronsson for the ALICE collaboration

Open charm / 37

Heavy flavor spectra in AA collision within a Langevin approach

Author: Andrea Beraudo¹

Co-authors: Arturo De Pace²; Francesco Prino¹; Marco Monteno¹; Marzia Nardi³; alfredo molinari⁴; wanda maria alberico⁵

¹ *Universita e INFN (IT)*

² *INFN*

³ *Unknown*

⁴ *Universita' di Torino*

⁵ *universita' di Torino e INFN*

Corresponding Author: andrea.beraudo@cern.ch

We present the updated outcomes of a theoretical study of heavy-flavor spectra in AA collisions. The core of the analysis is represented by the study of the propagation of heavy (c and b) quarks in the Quark Gluon Plasma: their dynamics is described through a relativistic Langevin equation, whose transport coefficients are evaluated within a thermal-field-theory approach. Results obtained with lattice-QCD transport coefficients will be also shown. Our Langevin approach, at variance with radiative energy-loss calculations, would lead (asymptotically) the heavy quarks to thermal equilibrium.

The numerical solution of the Langevin equation enters into a multi-step setup including

- Initial hard production of the q-qbar pairs, given by the POWHEG-BOX event generator (based on NLO pQCD);
- For the AA case: Langevin dynamics in the QGP background (evolving according to hydrodynamics);
- Hadronization according to the most up-to-date branching fractions and fragmentation functions;
- Decays into the experimentally accessible channels (open charm hadrons, displaced J/psi, heavy-flavor electrons).
- Evaluation of inclusive spectra (in pp and AA), R_{AA} and v_2 (in AA) and comparison with the most recent experimental results obtained in Pb-Pb collisions at the LHC at 2.76 TeV.

Our analysis represents an improvement of the calculation presented in Eur.Phys.J. C71 (2011) 1666. In particular measurements of exclusive open-charm spectra which became available at the LHC (at 2.76 and 7 TeV) allow now to fix the pp benchmark with tighter constraints.

Quarkonia / 38

System Size, Energy and Rapidity Dependence of Quarkonia Production Measured by the PHENIX Detector.

Author: Cesar Luiz da Silva¹

¹ *Los Alamos National Lab*

Corresponding Authors: durham@lanl.gov, slash@bnl.gov

The nuclear modification of quarkonia in heavy ion collisions involves a set of physics parameters like energy density, path length and initial state effects which can be controlled experimentally with variations in the beam energy, use of different species and rapidity ranges. The PHENIX detector recently collected high statistics data using combinations of heavy ion species and beam energies in a broad rapidity range. This talk will present the most recent measurements of J/ψ , ψ' , χ_c and Υ production in d+Au, Au+Au, Cu+Cu, Cu+Au collisions at $\sqrt{s_{NN}} = 200$ GeV, 62 GeV and 39 GeV in mid- ($|y| < 0.35$) and forward rapidities ($1.2 < |y| < 2.2$).

HF decays leptons / 39

Measurement of charm and bottom contributions to electrons from heavy quark decay at RHIC-PHENIX experiment

Author: Ryohji Akimoto¹

¹ *University of Tokyo*

Corresponding Author: r.akimoto@cns.s.u-tokyo.ac.jp

Heavy quark (charm and bottom) is a good probe to study the interaction between partons and quark gluon plasma (QGP). Heavy quarks are created mainly by initial hard scatterings and interaction between heavy quarks and QGP during full space-time evolution is clearly reflected in the final states.

A silicon vertex tracker (VTX) was installed at the PHENIX detector in 2011. The VTX enables us to measure charm and bottom contributions individually to electrons from heavy quark decay, and therefore to access the information of charm and bottom behavior inside QGP, which leads to greater understanding of the interaction.

In the presentation, the results of the electron measurement from charm and bottom decays for both p+p and Au+Au collisions will be presented.

HF particle correlations / 40

Heavy flavour correlation measurements in pp and Pb-Pb collisions with ALICE

Author: Sarah La Pointe¹

¹ *University of Utrecht (NL)*

Corresponding Author: s.lapointe@cern.ch

In high energy heavy-ion collisions heavy-flavour quarks are primarily produced in the early stage of the collision, through initial hard scatterings. This makes them an important probe of the high energy density QCD matter that is formed in the collision, since they should then experience the full evolution of the system. Heavy quarks can be studied through the measurements of open heavy-flavour hadrons and single leptons from heavy-flavour decays. Azimuthal angular correlations of single electrons and charged hadrons can be utilized to estimate the relative contribution of charm and beauty hadrons to the measured heavy-flavour single electron yield. In addition, an understanding on the production and fragmentation of heavy quarks can be examined through azimuthal angular correlations of heavy-flavour hadrons.

In this talk preliminary results on the relative contribution of electrons from beauty decays to the heavy-flavour single electron yield will be shown, along with the charm and beauty production cross section in pp collisions at $\sqrt{s} = 2.76$ TeV. These measurements will be compared to FONLL calculations. We also show results from the correlation analysis of D mesons and hadrons performed using pp collisions at $\sqrt{s} = 7$ TeV. Finally we discuss the status of heavy-flavour electron and charged hadron correlations in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV.

Cold nuclear matter / 41

Cold Nuclear Matter Effects on Heavy Quarks from PHENIX

Author: J. Matthew Durham¹

¹ *Los Alamos National Laboratory*

Corresponding Author: durham@lanl.gov

The flexibility of the Relativistic Heavy Ion Collider has allowed the PHENIX Collaboration to perform detailed measurements of heavy quark production in p+p, d+Au, Cu+Cu, and Au+Au collisions at 200 GeV, probing a wide range of temperature and collision geometry. Studies of d+Au collisions have shown significant and surprising cold nuclear matter effects on both charmonium and open heavy flavor, which imply that the initial state baseline for interpreting heavy quark transport in the hot partonic medium formed in Au+Au collisions is highly modified from the elementary p+p or pQCD shape. This talk will discuss recent measurements of cold nuclear matter effects on heavy quarks, and how they may influence our understanding of the cold nuclear environment and the interpretation of Au+Au data.

HF decays leptons / 42

Measurements of Non-photonic Electron Production and Azimuthal Anisotropy in Au+Au Collisions at STAR

Author: Mustafa Mustafa¹

¹ *Purdue University*

Heavy quarks are produced early in the heavy-ion collisions and are expected to interact with the created strongly interacting partonic medium differently from light quarks. Therefore, they can shed new light on understanding the medium's properties. Simultaneous measurements of spectra and azimuthal anisotropy of electrons from semileptonic decays of heavy flavor hadrons provide experimental insight to distinguish different energy loss theoretical models and to characterize heavy quarks medium interaction. During RHIC 2010 run, STAR has collected a large amount of minbias, central and high tower triggered data in Au+Au collisions at $\sqrt{s_{NN}} = 39, 62.4$ and 200 GeV with detector configuration for minimum photonic conversion background. From this data we will report a new high precision measurement of non-photonic electrons mid-rapidity invariant yield, improved nuclear modification factor and v_2 in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. We will also report measurements of mid-rapidity invariant yield at $\sqrt{s_{NN}} = 62.4$ and v_2 at $\sqrt{s_{NN}} = 39$ and 62.4 GeV.

Open charm / 43

D meson azimuthal anisotropy measured with ALICE experiment

Author: Andrea Festanti¹

¹ *Universita e INFN (IT)*

Corresponding Author: andrea.festanti@cern.ch

The ALICE experiment at the LHC is dedicated to the study of ultra-relativistic heavy-ion collisions, with the aim of investigating the high-density color deconfined state of strongly interacting matter that is expected to be formed in these collisions.

Heavy quarks serve as a probe of the dynamics of the medium since they are produced at the early stages of the collisions and they propagate through the created matter.

The D meson azimuthal anisotropy (v_2) is expected to bring insights into the degree of thermalization of charm quarks within the quark-gluon plasma.

A non-zero v_2 at low transverse momentum indicates a collective motion of charm quarks with respect to the bulk of created matter, while at high transverse momentum v_2 is sensitive to the path length dependence of the charm quark energy loss within the medium.

The measurement of D_0 , D^+ and D^+ elliptic

flow and D_0 RAA versus event

plane in semi-central Pb{Pb collisions at

$\sqrt{s_{NN}} = 2.76$ TeV will be pre-

sented. D mesons have been reconstructed via their hadronic decay channels

($D_0 \rightarrow K^0 \pi^+$, $D^+ \rightarrow K^0 \pi^+$ and $D^+ \rightarrow D^0 \pi^+$) in the central rapidity region.

44

Upsilon PbPb from Alice

45

CMS

Corresponding Author: torsten.dahms@cern.ch

46

Upsilon production in STAR

Author: Manuel Calderon De La Barca¹

¹ *UC Davis*

Corresponding Author: mcalderon@ucdavis.edu

Open charm / 47

The CBM heavy-quark program

Corresponding Author: i.vassiliev@gsi.de

The Compressed Baryonic Matter (CBM) experiment at FAIR is designed to explore the QCD phase diagram in the region of high baryon densities and moderate temperatures. The key CBM observables include particles containing hidden charm – J/ψ and Y' , open charm – D^0 , D^+ , D^* and L_c , low-mass vector mesons decaying into leptons and multi-strange hyperons. Particularly demanding is the measurement of open charm particles with very low multiplicities, which is based on the real time selection of displaced vertices with an accuracy of about $50 \mu\text{m}$. In the talk we discuss the problems of the detection of heavy-quark particles in fixed target experiments with relativistic heavy ion collisions.

48

Energy loss effects in cold nuclear matter

49

Theory overview

50

ATLAS

51

Heavy flavour particle correlations in Phenix

Corresponding Author: jason.kamin@cern.ch

New ideas from theory / 52

QCD factorization theorems for production of heavy quarkonia

Corresponding Author: jqiu@bnl.gov

53

AdS/CFT

54

PHENIX upgrades

Author: Cesar Luiz da Silva¹

¹ *Los Alamos National Lab*

Corresponding Author: slash@bnl.gov

Future experiments and upgrades / 55

Heavy flavour perspectives with CBM

Corresponding Author: v.friese@gsi.de

Summary / 56

Theory summary

Author: Hendrik van Hees¹

¹ *Justus-Liebig-Universität Giessen*

Corresponding Author: hendrik.vanhees@theo.physik.uni-giessen.de

Summary / 57

Experiment summary

Corresponding Author: enrico.scomparin@cern.ch

Quarkonia / 58

Recent results on quarkonium and open bottom suppression in PbPb collisions at the LHC with CMS

Author: Torsten Dahms¹

¹ *Ecole Polytechnique (FR)*

Corresponding Author: torsten.dahms@cern.ch

The Compact Muon Solenoid (CMS) has measured various quarkonium states via their decays into muon pairs in pp and PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. Quarkonia are especially relevant for studying the quark-gluon plasma since they are produced at early times of the collision and propagate through the medium, mapping its evolution. The most recent results on the production of prompt J/psi, psi(2S) and the first three Y states will be presented. At high transverse momenta ($p_T > 6.5$ GeV) and midrapidity ($|\eta| < 1.6$), psi(2S) are found to be more suppressed in PbPb collision than J/psi. Also the nuclear modification factors of the three Y states are found to decrease with decreasing binding energy, as expected in a sequential melting scenario. Furthermore, the nuclear modification factor of non-prompt J/psi from b-hadron decays gives access to study the energy loss of b-quarks in the quark-gluon plasma.

HF decays leptons / 59

Nonperturbative Heavy-Flavor Diffusion and Hadronization in a Hydrodynamic Description of Heavy-Ion Collisions

Author: Min He¹

Co-authors: Rainer Fries²; Ralf Rapp²

¹ Hunan University, China

² Texas A&M University

Corresponding Author: mhe@comp.tamu.edu

We compute open heavy-flavor (HF) transport in relativistic heavy-ion collisions by combining a strong-coupling treatment in both macro- and microscopic dynamics (hydro and nonperturbative diffusion interactions) [1]. The hydrodynamic bulk evolution is quantitatively constrained by bulk and multi-strange hadron spectra and elliptic flow [2]. In the Quark-Gluon Plasma (QGP) phase, heavy-quark diffusion coefficients are taken from a non-perturbative T -matrix approach. The latter leads to resonance formation close to T_c which is implemented as a hadronization (recombination) mechanism on a hydrodynamic hypersurface. In the hadronic phase, the diffusion of HF mesons is obtained from effective hadronic theory [3]. We compute observables at RHIC and LHC for both HF mesons and non-photonic electrons [4,5]. In particular, we suggest the R_{AA} and v_2 of the D_s mesons as a unique observable due to the coupling of charm to the strangeness enhancement in AA collisions [4], which allows to quantitatively test key components of our approach, including recombination and hadronic diffusion effects.

References:

- [1] M. He, R. J. Fries and R. Rapp, Phys.Rev. C86,014903 (2012).
- [2] M. He, R. J. Fries and R. Rapp, Phys. Rev. C85, 044911 (2012).
- [3] M. He, R. J. Fries and R. Rapp, Phys. Lett. B701, 445 (2011).
- [4] M. He, R. J. Fries and R. Rapp, arXiv:1204.4442 [nucl-th].
- [5] M. He, R. J. Fries and R. Rapp, arXiv:1208.0256 [nucl-th].

Cold nuclear matter / 60

Heavy-quarkonium suppression in p-A collisions from parton energy loss

Author: Stephane Peigne¹

¹ Subatech

Corresponding Author: peigne@subatech.in2p3.fr

The effects of parton energy loss in cold nuclear matter on heavy-quarkonium suppression in p-A collisions are discussed. It is shown from first principles that at large quarkonium energy E in the nucleus rest frame, the medium-induced energy loss is proportional to E . Using this result, a phenomenological model depending on a single free parameter is able to reproduce J/ψ and Upsilon suppression data in a broad xF-range and at various center-of-mass energies. These results strongly support energy loss as the dominant effect in heavy-quarkonium suppression in p-A collisions. Predictions for J/ψ and Upsilon suppression in p-Pb collisions at the LHC are presented.

Welcome / 61

Welcome, Head of the Department of Physics and Astronomy

HF decays leptons / 62

Heavy Quark Production in Pb-Pb Collisions at the LHC with the ATLAS Detector

Author: Mariusz Przybycien¹

¹ *AGH University of Science and Technology (PL)*

Corresponding Author: mariusz.przybycien@cern.ch

Bottom quarks are important probes to study the hot, dense medium produced in the heavy ion collisions. These heavier quarks are produced at a relatively early stage of the nucleus-nucleus collisions and may have reduced gluon radiation due to the suppression of small angle gluon radiation known as the 'dead cone effect'. Because of the heavy mass of b-hadrons, muons from semi-leptonic b-hadron decays tend to have a larger angle with respect to the jet axis. This information can be used to tag b-jets. In this talk, we present the inclusive heavy flavor suppression in Pb+Pb collisions at 2.76 TeV, which was obtained by studying single muons decaying semi-leptonically from the b- and c-quark containing hadrons.

New ideas from theory / 63

Gauge/gravity, thermalization and energy loss

Author: Wilke van der Schee¹

¹ *Utrecht University*

Holography provides an excellent tool to study (certain) strongly coupled gauge theories, which is especially useful if the state is very far from equilibrium. In this talk I will review current progress on thermalization, which is modelled by black hole formation, and highlight recent progress on energy loss using holography. Unfortunately I will also have to point out many caveats and uncertainties in these computations, in particular in relation to the energy loss of an energetic quark.

Quarkonia / 64

Quarkonium at finite temperature

Author: Nora Brambilla¹

¹ *Institut fuer Theoretische Physik*

Corresponding Author: nora.brambilla@cern.ch

I will review the effective field theory description of quarkonium interaction at finite temperature and its phenomenological implications.

Future experiments and upgrades / 65

Future of Heavy Quark Measurement at sPHENIX

Author: Cesar Luiz da Silva¹

¹ *Los Alamos National Lab*

Corresponding Authors: slash@bnl.gov, durham@lanl.gov

The PHENIX Collaboration is designing a new detector configuration (sPHENIX) to continue the studies of the recent discoveries at RHIC. The new detector will be based on a strong solenoid magnet with full azimuthal calorimetry and tracking coverage as well as displaced-vertex measurement capabilities. A new landmark of sQGP interaction, color screening and initial state effect studies can be explored at RHIC with measurements like heavy flavor tagged jets, hadron(γ)-heavy flavor correlation and identification of different quarkonia states in p+p, p+A and A+A collisions. This presentation will detail the conceptual design, the staging and the physics which can be addressed in this concept.

HF particle correlations / 66

Heavy Flavor Dileptons at PHENIX

Author: Jason Adrian Kamin¹

¹ *Johann-Wolfgang-Goethe Univ. (DE)*

Corresponding Author: jason.kamin@cern.ch

Charm and bottom quarks are very effective probes of the Quark Gluon Plasma (QGP) created in heavy ion collisions as they are created in the initial hard scattering and therefore fully traverse the hot nuclear medium before fragmentation into the vacuum. Along with a well-established baseline measurement, any observed modification to heavy-ion heavy flavor spectra can yield insight into the interaction between the heavy quarks and the QGP.

Dilepton correlations provide an extremely useful window to open heavy flavor. Semi-leptonic decays of B and D mesons can be studied through both the unlike sign and like sign correlated dilepton pairs. They yield a clean signal which is largely free of contamination and provide reasonable separation of charm and bottom. Moreover, they access a region of phase space where pQCD calculations are more reliable than for single lepton observables.

The PHENIX experiment has extrapolated the charm and bottom cross sections through both leading order and next-to-leading order

calculations for $\sqrt{s} = 200\text{-GeV}$ and $\sqrt{s} = 500\text{-GeV}$. These heavy flavor results from two particle correlations have been explored in both $p+p$ and $d+Au$ collisions at PHENIX. Heavy flavor production determined in both the dielectron and dimuon channels will be shown.

HF particle correlations / 67

From the RAA to azimuthal correlations: what can we learn from heavy-quark observables?

Author: Marlene Nahrgang^{None}

Corresponding Author: nahrgang@th.physik.uni-frankfurt.de

The observation of strong jet quenching and the suppression of high- p_t hadrons in relativistic heavy-ion collisions are striking experimental signatures for the formation of a deconfined QCD plasma in which partons suffer from medium-induced energy loss. In particular, heavy quarks are considered as suitable probes for revealing the properties of the produced matter as they are created at very early stages in hard scattering processes and assumed not to thermalize completely within the medium. Typical observables for the interaction of heavy quarks with the medium constituents are the nuclear modification factor R_{AA} and the elliptic flow v_2 , which we present within a combined Monte-Carlo approach and realistic fluid dynamic description of the expanding plasma for both RHIC and LHC energies. In the main part of this talk we will investigate the potential of correlations between heavy quarks and anti-quarks to reveal basic principles of energy loss scenarios at LHC. At low p_t any correlation of the initially heavy quark-antiquark pair is lost due to thermalization, at larger p_t , however, these correlations in p_t and azimuthal angle ϕ survive and show distinctive features for purely elastic and elastic plus radiative energy loss mechanisms. We discuss these results in view of the core-corona effect and the centrality dependence.

Quarkonia / 68

Upsilon measurements in STAR

Author: Manuel Calderon De La Barca¹

¹ UC Davis

Corresponding Author: mcalderon@ucdavis.edu

We report on measurements of Upsilon mesons in the dielectron channel with the STAR detector at RHIC.

Upsilon mesons are expected to be sensitive to the deconfinement transition in the Quark Gluon Plasma, and

to be useful in determining the medium temperature. We compare the yield obtained in Au+Au data, separated into three centrality classes, to the measured cross section from $p+p$ collisions. Our data are compared

to models of bottomonium dissociation.