

Measurement of Di-electron Continuum in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV by PHENIX using the Hadron Blind Detector

Sky D. Rolnick for the PHENIX Collaboration

UC Riverside

E-mail: skymeson@gmail.com

Abstract. Dielectrons play an important role in heavy ion collisions since they are produced during all stages of the collision and interact only electromagnetically. The PHENIX experiment was upgraded with the Hadron Blind Detector which provides excellent background rejection by removing pairs from both partially reconstructed Dalitz decays and photon conversions. In these proceedings we will report on the results obtained from 2009 data in $p+p$ collisions using the HBD as well as the Au+Au results obtained in 2010.

1. Introduction


Dileptons are very useful probes for studying many interesting puzzles in Heavy Ion Physics. They are produced during all stages of the collision and do not interact strongly with the medium. The dielectron continuum offers a rich and diverse region of physics to explore. It allows us to study such effects as chiral symmetry restoration or possible medium modifications of vector mesons. There are several known sources which contribute to the dielectron continuum including thermal radiation, direct photons, vector meson decays, and open charm and bottom yields. The cleanness of the dielectron measurements, compared to hadronic measurements, provides probes into multi-component flow and energy loss mechanisms and will provide checks on a variety of production mechanisms.

Previous measurements made by PHENIX in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV have shown strong enhancement of electron pairs compared to the expected hadronic cocktail in the low mass region $m_{ee} = 0.15 - 0.75$ GeV/ c^2 by a factor of $4.7 \pm 0.4(stat.) \pm 1.5(syst.)$ for minimum bias collisions. Such a large enhancement was not expected and has posed a mystery to many theorists [1]. The very low signal to background ratio of $\sim 1/200$ generates large systematics in the region of interest. To improve on this, a new measurement has been made by PHENIX utilizing the Hadron Blind Detector, which reduces the backgrounds and systematic uncertainty. In these proceedings, we will discuss the use of the Hadron Blind Detector as well as dielectron results for $p+p$ collisions and Au+Au collisions for three centrality bins.

2. Hadron Blind Detector

The Hadron Blind Detector is a novel type of Cherenkov detector utilizing a triple GEM configuration with a CSI photo-cathode layer operated in pure CF₄. The HBD has achieved the highest figure of merit of any gas Cherenkov detector to date $No = 322$ cm⁻¹ and designed with

a very low material budget of just 2.4% total radiation lengths [7]. It was successfully operated in 2009 $p+p$ data and 2010 Au+Au data sets.



- * Windowless Cherenkov detector
- * Tripple GEM
- * CSI photo-cathode
- * Pure CF₄: N₀ = 322 cm⁻¹
- * 2.4% total radiation length.

Figure 1. The HBD concept design.

The HBD was designed to work by preserving the opening angle of charged particles using the field free region of the PHENIX central arms. Electrons are tagged by the amount of Cherenkov radiation they produce and since heavier meson decays have large opening angles compared to Dalitz decays and conversions, it is possible to separate these different sources. Cherenkov light is continuously radiated along the path of the electron such that a small unfocused blob of Cherenkov light is formed on the surface of the detector. The typical photoelectron (p.e.) response is measured to be ~ 20 p.e. for a single electron and ~ 40 p.e for a double electron. In principle the difference in response should distinguish between single and double electrons.[3]

Several reconstruction algorithms have been developed and studied with a single electron tracking efficiency of $\sim 90\%$. [4] The main source of background arises from partially reconstructed π^0 Dalitz decays or gamma conversion in which one electron is lost either due to acceptance or tracking inefficiencies. The HBD allows for the tagging and removal of these uncorrelated electron pairs thereby reducing the background.

3. Hadronic Cocktail

The hadronic cocktail consists of dielectron decays from known hadronic sources including Dalitz decays of π^0 , η , and η' as well as direct decays from vector mesons ρ , ω , ϕ and J/ψ . The method for obtaining the cocktail has been outlined in [2] and uses a combined fit from neutral and charged pion spectra to a modified Hagedorn function. The p_T spectra of the other mesons is derived using m_T scaling of the π^0 spectra and normalized using the meson-to-pion ratio at high p_T . The momentum resolution of PHENIX tracking is taken into account using a smearing technique. The additional material of the HBD was taken into account yielding higher rates of conversion electrons which were modeled using a full Geant simulation of the PHENIX detector. The line shape of the J/ψ was also modified to account for Bremsstrahlung radiation produced by the additional material. The difference in magnetic field configurations are reflected in the acceptance of the electron pairs which is also accounted for. The heavy flavor contributions from charm and bottom are estimated using MC@NLO [5] as opposed to PYTHIA [6] since it is found to reproduce dielectron yields, especially at high p_T . The cocktail used matches very well to $p+p$ data over the whole mass range.




Figure 2. The left panel shows the PHENIX dielectron mass spectrum for Run9 $p+p$ overlaid with the cocktail shown by a solid line. These results show excellent agreement between data and cocktail (lower panel). The signal to background comparison for 2005 $p+p$ data (without HBD) to 2009 $p+p$ data (with HBD) is shown in the right panel.

4. Results for $p+p$ collisions

The measurements in $p+p$ collisions establish the baseline spectra without any medium effects and can be used in comparison to Au+Au results. They also provide a testing ground for understanding the HBD. The S/B ratio is improved by a factor of 5 in the low mass region by using the HBD which demonstrates the effectiveness of the HBD in background rejection.

Combinatorial background is subtracted using a like-sign subtraction technique which uses electron pairs with the same sign to estimate the correlated and uncorrelated background contributions. Examples of correlated like-sign pairs include cross pairs i.e. $\pi^0 \rightarrow \gamma\gamma \rightarrow e^+e^-e^+e^-$ and jet pairs where both electrons are correlated through the jet. Differences in acceptance between like-sign and unlike-sign are corrected with a relative acceptance correction. The main advantage in using like-sign subtraction technique is that combinatorial pairs are produced at the same rate as unlike-sign foreground and the distributions will be absolutely normalized. One drawback of this technique is that the statistical precision is limited compared to mixed event techniques.

5. Results for Au+Au collisions

In Au+Au collisions, the large occupancy rate in the HBD detector presents a challenge especially in central events. The Cherenkov signal begins to get overwhelmed by other sources of radiation primarily from scintillation but also heavy ionized particles or low momentum electrons looping back into the HBD. In central events, the mean scintillation per pad in the HBD approaches ~ 7 p.e. and when distributed over the average cluster size of 3 pads per cluster, the response from scintillation becomes comparable to the single electron response. Two techniques have been developed to reduce the background scintillation light including a mean pad subtraction based on centrality and a local background subtraction by looking at nearby pads around the clusters. Using the mean pad subtraction technique, occupancy rates can be reduced to $\sim 15\%$ for central collisions.




Figure 3. Dielectron mass spectra from Au+Au collisions for three centrality bins (a) 60-92%, (b) 40-60%, (c) 20-40%. The ratios to the cocktail are also shown in the lower panels.

Two neural networks were utilized to improve the electron ID and single/double electron separation. The neural networks were trained using detailed GEANT simulations. For the first stage of the analysis, a neural net selects tracks with a high probability for being an electron where studies have shown that the purity of the electron sample approaches 90%. A second neural net is used to reject contributions from double electrons in the HBD. This has been shown to remove up to 70% of double electrons reconstructed as singles by the central arm.[9]

The background subtraction technique used for Au+Au is slightly more complicated than that used for $p+p$ due to the larger combinatorial background in central events. There are two steps to remove the two types of background. First, a mixed event subtraction is used to remove the combinatorial background and then a like-sign subtraction is used to account for the correlated background. To isolate the correlated spectra from the like-sign pairs, background is subtracted using mixed events and the remaining correlated like-sign pairs are then corrected for relative acceptance, where FG12 is the unlike-sign foreground, BG12 is the unlike sign mixed

event background, and FG(LS) and BG(LS) are the foreground and background for the like-sign respectively.

$$S = FG12 - N \times BG12 = \alpha \times (FG(LS) - N \times BG(LS))$$

The normalization is done after a pair cut efficiency is applied and a normalization factor is calculated such that the relationship $\langle BG12 \rangle = 2\sqrt{\langle BG11 \rangle \langle BG22 \rangle}$ holds. The results are shown in Fig. 3 for 3 different centrality bins. In Fig. 4 these results are compared to the Au+Au obtained in Run 4 before the HBD was installed.[10]




Figure 4. Comparison of Au+Au results to previous published results. We see that the latest results using the HBD are fully consistent with previous results within the systematic errors. The ratios are plotted as a function of N_{part} integrated over two different mass regions.

6. Summary

The dielectron mass spectrum has been presented for $p+p$ and for three centrality bins of Au+Au. These results are consistent with previous published results at PHENIX. There is a hint of enhancement in the 20-40% centrality bin in the low- and intermediate-mass region. Currently this is not yet conclusive given present uncertainties and work is in progress to improve the systematics.

References

- [1] C. Gale, “Electromagnetic radiation in heavy ion collisions.” [arXiv:1208.2289[hep-ph]]
- [2] A. Adare *et al.* [PHENIX Collaboration], Phys. Rev. C 81, 034911 (2010) [arXiv:0912.0244 [nucl-ex]].
- [3] Woody *et al.* “Initial performance of the PHENIX Hadron Blind Detector at RHIC,” (NSS/MIC), 2009 IEEE , pp.1002-1008, Oct. 24 2009-Nov. 1 2009
- [4] QM 2011 poster by Yosuke Watanabe.
- [5] S. Frixione and, B.R. Webber, JHEP 0206,029 (2002). [hep-ph/0204244]
- [6] T. Sjostrand, S. Mrenna and P.Z. Skands, JHEP 0605 (2006) 026. [hep-ph/0603175]
- [7] W. Anderson, *et al.* Nucl. Instr. and Meth. A (2011). [doi:10.1016/j.nima.2011.04.015]
- [8] S. Afanasiev *et al.* (PHENIX), Enhancement of the dielectron continuum in $\sqrt{s_{NN}} = 200$ GeV Au+Au collisions (2007), submitted to Phys. Rev. Lett., Preprint [arXiv:0706.3034 [nucl-ex]]
- [9] “Dielectron Analysis in Au+Au Collisions at $\sqrt{s_{NN}} = 200$ GeV with the Hadron Blind Detector.” PHENIX Internal Analysis note 1040.
- [10] I. Tserruya, Quark Matter 2012 proceedings. [arXiv:1211.6002 [nucl-ex]]