

Re-evaluation of the Parton Distributions of Strange Quarks in the Nucleon

Uses the final HERMES kaon Multiplicities from SIDIS

[A. Airapetian et al., Phys. Rev. D (in press)]



H. E. Jackson - DIS 2013



Measurement of parton distributions of strange quarks in the nucleon from charged-kaon production in deep-inelastic scattering on the deuteron

HERMES Collaboration

HERMES Collaboration / Physics Letters B 666 (2008) 446–450

447

^{a,a} Yerevan Physics Institute, 375036 Yerevan, Armenia

ARTICLE INFO

Article history:

Received 3 June 2008

Received in revised form 18 July 2008

Accepted 23 July 2008

Available online 6 August 2008

Editor: L. Rolandi

PACS:

13.60.-r

13.88.+e

14.20.Dh

14.65.-q

ABSTRACT

The momentum and helicity density distributions of the strange quark sea in the nucleon are obtained in leading order from charged-kaon production in deep-inelastic scattering on the deuteron. The distributions are extracted from spin-averaged K^\pm multiplicities, and from K^\pm and inclusive double-spin asymmetries for scattering of polarized positrons by a polarized deuterium target. The shape of the momentum distribution is softer than that of the average of the \bar{u} and \bar{d} quarks. In the region of measurement $0.02 < x < 0.6$ and $Q^2 > 1.0 \text{ GeV}^2$, the helicity distribution is zero within experimental uncertainties.


© 2008 Elsevier B.V. All rights reserved.




The strange sea: $S(x)$ from K^\pm multiplicities

$$\frac{dN^{K^\pm}}{dN^{\text{DIS}}} = \frac{Q(x) \int D_Q^K(z) dz + S(x) \int D_S^K(z) dz}{5 Q(x) + 2 S(x)} \xrightarrow{x > 0.3} \frac{\int D_Q^K(z) dz}{5}$$

P.L. B666 (2008) 466



- $S(x)$ from CTEQ6L with $\int D_Q^K(z) dz$ & $\int D_S^K(z) dz$ as free parameters (dotted) does not fit the data



- $S(x)$ much softer than assumed by current PDFs (mainly based on $\nu N \rightarrow \mu^+ \mu^- X$)

Take $\int D_S^K(z) dz = 1.27 \pm 0.13$ from de Florian et al.

Signal for $S(x)=0$ at LO


In leading order:

$$M^{K^\pm}(x) \equiv \frac{d^2N^K(x)}{d^2NDIS(x)} = \frac{[f(x)]}{[f(x) + S(x)]}$$

where $f(x) = S(x)/Q(x) \ll 1$.

$$5M^{K^\pm}(x) = \left[\int \mathcal{D}_Q^K(z) dz + f(x) \right]$$

$$\frac{d5M^{K^\pm}(x)}{dx} = \frac{df(x)}{dx} \left[\left(1 - \frac{4}{5}f(x) \right)$$




$dM(x)/dx < 0$ if $S(x) \neq 0$


Signature for $S(x)=0 \rightarrow dM(x)/dx \geq 0$

Fit of Q(x) component to $M(x)=p[0]+p[1]*x_{bj}$


kaon multiplicities, plb66 vs 2012




Revised fit to kaon charged multiplicity



Fit to Component arising from


$$S(x) \int D^K(z) dz = x^{-a} e^{-x/b} (1-x)$$


Strange parton distribution $S(x)$ - revised



$$\text{Fit}(x) = x^{-0.867 \pm 0.019} e^{-0.331 \pm 0.014}(1-x)$$

Comparison with prediction of the NNPDF Collaboration



Summary

- $S(x) \approx 0$ with the measurement error for $x \geq 0.15$, as reported in PLB666, 446 (2008).
- $S(x)$ is similar in shape but ≈ 0.6 in magnitude of the data reported in 2008.
- In magnitude, but not detailed shape , $S(x)$ as extracted here is close to the recent predictions of the NNPDF collaboration (**NPB 855, 153 (2012)**).
- The shape of $S(x)$ suggests the possibility that the strange quark pdf may be a surrogate for a sea dominated by the gluon splitting component (see **Chang & Peng, PLB 704, 197 (2011)**).

Extra's

Comparison PLB666 with HERMES (2013)

