

Fundamental Concepts in Particle Physics

Lecture 3 :
Towards the Standard Model

Géraldine SERVANT
CERN-Th



Abelian versus non-abelian gauge theories


The (Yang-Mills) action $\mathcal{L}_{YM} = \bar{\Psi}(i\gamma^\mu D_\mu - m)\Psi - \frac{1}{2}F_{\mu\nu}F^{\mu\nu}$ is invariant under

$$\Psi(x) \rightarrow U(x)\Psi(x)$$

Abelian U(1) symmetry

$$U(x) = e^{iq\theta(x)}$$

$$A_\mu(x) \rightarrow A_\mu + \frac{i}{e}(\partial_\mu U)U^\dagger$$



Non-abelian SU(N)

$$U(x) = e^{ig\theta^a(x)T^a}$$

T^a : N^2-1 generators ($N \times N$ matrices) acting on

$$A_\mu(x) = A_\mu^a T^a$$

$$A_\mu(x) \rightarrow U A_\mu U^\dagger - \frac{i}{g}(\partial_\mu U)U^\dagger$$

infinitesimal transformation

$$U(x) = 1 + ig\theta^a(x)T^a + \mathcal{O}(\theta^2)$$

$$A_\mu^a(x) \rightarrow A_\mu^a + \partial_\mu\theta^a - g f^{abc}\theta^b A_\mu^c$$

$$D_\mu\Psi = (\partial_\mu + iqA_\mu)\Psi$$

$$D_\mu\Psi = (\partial_\mu - igA_\mu^a T^a)\Psi$$

The gauge symmetries of the Standard Model

Gauge Group $U(1)_Y$ (abelian)

$$\psi' = e^{+iY\alpha_Y} \psi,$$

$$B'_\mu = B_\mu - \frac{1}{g'} \partial_\mu \alpha_Y$$

$$B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$$

$$D_\mu \psi_R = (\partial_\mu + i g' Y B_\mu) \psi_R$$

Gauge Group $SU(2)_L$ acts on the two components of a doublet $\Psi_L = (u_L, d_L)$ or (ν_L, e_L)

$$\Psi_L \rightarrow e^{-iT^a \alpha^a} \psi_L \quad U = e^{-iT^a \alpha^a} \quad T^a = \sigma^a/2 \quad \text{Pauli matrices}$$

$$W_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g \epsilon^{abc} W_\mu^b W_\nu^c, \quad a = 1, \dots, 3$$

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \sigma_2 = -i \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$D_\mu \psi_L = (\partial_\mu - i g W_\mu^a T^a) \psi_L$$

Gauge Group $SU(3)_c$ $q=(q_1, q_2, q_3)$ (the three color degrees of freedom)

$$q \rightarrow e^{-iT^a \alpha^a} q \quad U = e^{-iT^a \alpha^a} \quad [T^a, T^b] = if^{abc} T^c \quad (3 \times 3) \text{ Gell-Mann matrices}$$

$$G_\mu^a T^a \rightarrow U G_\mu^a T^a U^{-1} - \frac{i}{g} \partial_\mu U U^{-1}$$

$$G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g f^{abc} G_\mu^b G_\nu^c, \quad a = 1, \dots, 8$$

$$D_\mu q = (\partial_\mu - i g G_\mu^a T^a) q$$

$$\lambda_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \lambda_2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \lambda_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\lambda_4 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \quad \lambda_5 = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix} \quad \lambda_6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

$$\lambda_7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix} \quad \lambda_8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}$$

The gauge symmetries of the Standard Model

Gauge Group $U(1)_Y$ (abelian)

$$\psi' = e^{+iY\alpha_Y} \psi,$$

$$B'_\mu = B_\mu - \frac{1}{g'} \partial_\mu \alpha_Y$$

$$B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$$

$$D_\mu \psi_R = (\partial_\mu + i g' Y B_\mu) \psi_R$$

Gauge Group $SU(2)_L$

$$\Psi_L \rightarrow e^{-iT^a \alpha^a} \psi_L \quad U = e^{-iT^a \alpha^a}$$

$$W_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g \epsilon^{abc} W_\mu^b W_\nu^c, \quad a = 1, \dots, 3$$

$$D_\mu \psi_L = (\partial_\mu - i g W_\mu^a T^a) \psi_L$$

Gauge Group $SU(3)_c$

$$q \rightarrow e^{-iT^a \alpha^a} q \quad U = e^{-iT^a \alpha^a}$$

$$G_\mu^a T^a \rightarrow U G_\mu^a T^a U^{-1} - \frac{i}{g} \partial_\mu U U^{-1}$$

$$G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g f^{abc} G_\mu^b G_\nu^c, \quad a = 1, \dots, 8$$

$$D_\mu q = (\partial_\mu - i g G_\mu^a T^a) q$$

$$\mathcal{L}_{YM} = \bar{\Psi} (i \gamma^\mu D_\mu - m) \Psi - \frac{1}{2} F_{\mu\nu} F^{\mu\nu}$$

all Standard Model fermions
carry $U(1)$ charge

$\Psi_L = (u_L, d_L) \text{ or } (\nu_L, e_L)$

only left-handed fermions charged
under it \rightarrow chiral interactions

$q = (q_1, q_2, q_3)$

all quarks transform under it
 \rightarrow vector-like interactions

The lagrangian of the Standard Model

$$\mathcal{L}_{\text{gauge}} = -\frac{1}{4}G_{\mu\nu}^a G^{a\mu\nu} - \frac{1}{4}W_{\mu\nu}^a W^{a\mu\nu} - \frac{1}{4}B_{\mu\nu} B^{\mu\nu} \quad \text{describe massless gauge bosons}$$

$$\mathcal{L}_{\text{Fermion}} = \sum_{\text{quarks}} i\bar{q}\gamma^\mu D_\mu q + \sum_{\psi_L} i\bar{\psi}_L \gamma^\mu D_\mu \psi_L + \sum_{\psi_R} i\bar{\psi}_R \gamma^\mu D_\mu \psi_R \quad \text{describe massless fermions and their interactions with gauge bosons}$$

only left-handed fermions all fermions carrying a $U(1)_Y$ charge
 i.e. all Standard Model fermions

$$D_\mu \psi_R = [\partial_\mu + ig' Y B_\mu] \psi_R$$

$$\mathcal{L}_{\text{Higgs}} = (D_\mu \Phi)^\dagger D_\mu \Phi + \mu^2 \Phi^\dagger \Phi - \lambda (\Phi^\dagger \Phi)^2 \quad \xrightarrow{\text{gives mass to EW gauge bosons}} \quad \frac{1}{2} M_Z^2 Z_\mu Z^\mu + M_W^2 W_\mu^+ W^{-\mu}$$

$$D_\mu \Phi = \left[\partial_\mu - i \frac{g}{\sqrt{2}} (\tau^+ W_\mu^+ + \tau^- W_\mu^-) - i \frac{g}{2} \tau_3 W_\mu^3 + i \frac{g'}{2} B_\mu \right] \Phi \quad : \text{covariant derivative of the Higgs}$$

H charged under $SU(2) \times U(1)_Y$

responsible for electroweak symmetry breaking!

$$\mathcal{L}_{\text{Yukawa}} = -Y_l \bar{L} \Phi \ell_R - Y_d \bar{Q} \Phi d_R - Y_u \bar{Q} \tilde{\Phi} u_R + \text{h.c.} \quad \xrightarrow{\text{gives mass to fermions}}$$

$$SU(3) \times SU(2)_L \times U(1)_Y \longrightarrow SU(3) \times U(1)_{em}$$

8 massless gluons

3 massive gauge bosons
 $W^+ W^- Z_0$

8 massless gluons
 1 massless photon γ
 remaining unbroken symmetry

The W and Z bosons interact with the Higgs medium, the γ doesn't.

$$\mathcal{L}_{\text{gauge}} = -\frac{1}{4}G_{\mu\nu}^a G^{a\mu\nu} - \frac{1}{4}W_{\mu\nu}^a W^{a\mu\nu} - \frac{1}{4}B_{\mu\nu} B^{\mu\nu}$$

$SU(3)_c$

$$G_{\mu\nu}^a = \partial_\mu G_\nu^a - \partial_\nu G_\mu^a + g f^{abc} G_\mu^b G_\nu^c$$

$$W_{\mu\nu}^a = \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g \epsilon^{abc} W_\mu^b W_\nu^c,$$

$$B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$$

$SU(2)_L$

in mass eigen state basis


$$W_\mu^\pm = \frac{W_\mu^1 \mp i W_\mu^2}{\sqrt{2}}$$

$$\cos \theta_W = g / \sqrt{g^2 + g'^2}$$


$$Z_\mu = W_\mu^3 \cos \theta_W + B_\mu \sin \theta_W$$

$$A_\mu = -W_\mu^3 \sin \theta_W + B_\mu \cos \theta_W$$

$$\sin \theta_W = g' / \sqrt{g^2 + g'^2}$$







three gauge
boson vertex



four gauge
boson vertex



no such
interactions
for photon!



The Standard Model of Particle Physics

- one century to develop it
- tested with impressive precision
- accounts for all data in experimental particle physics


Forces



The Higgs was the only remaining unobserved piece and is a portal to new physics hidden sectors

The (adhoc) Higgs Mechanism (a model without dynamics)


EW symmetry breaking is described by the condensation of a scalar field



$$\Phi = \begin{bmatrix} \phi^+ \\ v + H + i\varphi_Z \end{bmatrix} \sqrt{2}$$

Background value, Higgs medium
 Higgs boson: excitation of the higgs medium

The Higgs selects a vacuum state by developing a non zero background value. When it does so, it gives mass to SM particles it couples to.



$$V(\Phi) = \frac{\mu^2}{2} \Phi^\dagger \Phi + \frac{\lambda}{4} \Phi^\dagger \Phi$$

Why is μ^2 negative?

the puzzle:

We do not know what makes the Higgs condensate.

We ARRANGE the Higgs potential so that the Higgs condensates but this is just a parametrization that we are unable to explain dynamically.

Historically

Fermi Theory


$$n \xrightarrow{W^\pm} p + e^- + \bar{\nu}_e$$

$$\mathcal{L} = G_F (\bar{n}p)(\bar{\nu}_e e)$$

$$\mathcal{A} \propto G_F E^2$$

- no continuous limit
- inconsistent above 300 GeV

(paper rejected by Nature: declared too speculative !)



Gauge theory


microscopic theory


(exchange of a massive spin 1 particle)

$$G_F = \frac{\sqrt{2}g^2}{8m_W^2}$$


exp: $m_W = 80.4$ GeV

- ➡ $g \approx 0.6$, ie, same order as $e=0.3$
- unification EM & weak interactions





muon decay



beta decay

- ◆ We have quantized free fields
- ◆ We have introduced interactions
(particle creation and annihilation can only take place in theory with interactions)

We now would like to compute probability of processes like for instance a two-body decay $a \rightarrow c+d$ or a two-body reaction $a+b \rightarrow c+d$

"S-matrix approach" -> calculate probability of transition between two asymptotic states

The S-matrix

We consider a state $|a\rangle(t)$ which at an initial time t_i is labelled $|a\rangle$. Similarly we consider a state $|b\rangle(t)$ which at a final time t_f is labelled $|b\rangle$

At t_f the state $|a\rangle(t)$ is evolved as $e^{-iH(t_f-t_i)}|a\rangle$

where H is the hamiltonian of the theory

The amplitude for the process in which the initial state $|a\rangle$ evolves into the final state $|b\rangle$ is given by

$$\mathcal{M} = \langle b | e^{-iH(t_f-t_i)} | a \rangle$$

the final state is a set of well-separated particles long after the interaction

evolution operator
"S-matrix"

the initial state is either a one-particle (decay) or two well-separated particles (scattering), long before interaction happens

$|a\rangle$ and $|b\rangle$ are both described by free fields

The probability of the process is given by $|\mathcal{M}|^2$

and that can be linked to a transition rate per volume unit as measured by an experiment

Link to observables

- ◆ cross section: reaction rate per target particle per unit incident flux

$$\frac{[1/\text{time}]}{[1/(\text{time length}^2)]}$$

--> has units of a surface
measured in multiples of 1 barn= 10^{-24} cm^2

typical relevant LHC cross sections ~ in pb


$$1 \text{ picobarn} = 1 \text{ pb} = 10^{-36} \text{ cm}^2$$

- ◆ Decay width (inverse of lifetime of a particle) = transition rate
has dimension [1/time]

Example: decay width of EW gauge bosons

$$\Gamma \propto |\mathcal{M}|^2$$



scales as the square of the coupling constant



Z couplings to fermions

The coupling of Z to any fermion is proportional to $I_3 - \sin^2 \theta_W Q$
 where $I_3 = \pm \frac{1}{2}$ is z-component of weak isospin and Q is electric charge

$$\sin^2 \theta_W = 0.231$$



for the quarks:

u_L	$I_3 = +1/2$	$Q = +2/3$
u_R	$I_3 = 0$	$Q = +2/3$
d_L	$I_3 = -1/2$	$Q = -1/3$
d_R	$I_3 = 0$	$Q = -1/3$

and similarly for c, s , and b (t is too heavy for the Z to decay into it)

for the leptons:

e_L	$I_3 = -1/2$	$Q = -1$
e_R	$I_3 = 0$	$Q = -1$
ν_{e_L}	$I_3 = +1/2$	$Q = 0$

and similarly for $\nu, \tau, \nu_\mu, \nu_\tau$

Branching fractions for Z decay

for the quarks:

$$u_L \quad I_3 = +1/2 \quad Q = +2/3$$

$$u_R \quad I_3 = 0 \quad Q = +2/3$$

$$d_L \quad I_3 = -1/2 \quad Q = -1/3$$

$$d_R \quad I_3 = 0 \quad Q = -1/3$$

and similarly for c,s, and b (t is too heavy for the Z to decay into it)

for the leptons:

$$e_L \quad I_3 = -1/2 \quad Q = -1$$

$$e_R \quad I_3 = 0 \quad Q = -1$$

$$\nu_{e_L} \quad I_3 = +1/2 \quad Q = 0$$

and similarly for $\nu, \tau, \nu_\mu, \nu_\tau$

The decay rate is proportional to the square of the coupling constant $I_3 - \sin^2 \theta_W Q$

Also, for quarks, there is an additional factor $(1 + \frac{\alpha_s}{2\pi})$ where $\alpha_s = g_s^2/4\pi = 0.118$ due to the additional gluon emission


$$B(Z \rightarrow e^+ e^-) = B(Z \rightarrow e_L^+ e_L^-) + B(Z \rightarrow e_R^+ e_R^-)$$

$$B(Z \rightarrow e_L^+ e_L^-) = \frac{\Gamma(Z \rightarrow e_L^+ e_L^-)}{\sum_{all \ particles} \Gamma(Z \rightarrow particle, antiparticle)}$$

$$B(Z \rightarrow \nu \bar{\nu}) = B(Z \rightarrow \nu_e \bar{\nu}_e) + B(Z \rightarrow \nu_\mu \bar{\nu}_\mu) + B(Z \rightarrow \nu_\tau \bar{\nu}_\tau) \\ = 3B(Z \rightarrow \nu_e \bar{\nu}_e) = 20\%$$

$$B(Z \rightarrow e^+ e^-) = B(Z \rightarrow \mu^+ \mu^-) = B(Z \rightarrow \tau^+ \tau^-) = 3.33\%$$

$$B(Z \rightarrow all \ hadrons) = 3 \times [B(Z \rightarrow u\bar{u}) + B(Z \rightarrow d\bar{d}) + B(Z \rightarrow s\bar{s}) \\ + B(Z \rightarrow c\bar{c}) + B(Z \rightarrow b\bar{b})] = 69.9\%$$



Branching fractions for W decay

$$W^- \rightarrow e^- \bar{\nu}_e, \mu^- \bar{\nu}_\mu, \tau^- \bar{\nu}_\tau, d' \bar{u}, s' \bar{c}.$$

$$\begin{aligned} BR(W^- \rightarrow e^- \bar{\nu}_e) &= BR(W^- \rightarrow \mu^- \bar{\nu}_\mu) = BR(W^- \rightarrow \tau^- \bar{\nu}_\tau) \\ &= \frac{1}{3 + 6(1 + \alpha_s/\pi)} = 0.108, \end{aligned}$$

$$BR(W^- \rightarrow \text{hadrons}) = \frac{6(1 + \alpha_s/\pi)}{3 + 6(1 + \alpha_s/\pi)} = 0.675.$$

The Standard Model

	Q	d	u	L	e	B	W	g	H
$SU(3)_C$	3	3	3	1	1	1	1	8	1
$SU(2)_L$	2	1	1	2	1	1	3	1	2
$U(1)_Y$	+1/6	-1/3	+2/3	-1/2	+1	0	0	0	-1/2
spin	-1/2	+1/2	+1/2	-1/2	+1/2	1	1	1	0
flavor	3	3	3	3	3	1	1	1	1