

Detectors for Particle Physics


An Introduction

D. Bortoletto
Purdue University

INSTRUMENTATION

- “New directions in science are launched by new tools much more often than by new concepts. The effect of a concept-driven revolution is to explain old things in new ways. The effect of a tool-driven revolution is to discover new things that have to be explained”

Freeman Dyson



- “Instrumentation for the 21st century. No one does it better than physicists when it come to innovation for instrumentation, and thus the future of all scientific fields rests on our hands.”

Michael S. Turner


Why do we develop new instruments

Discoverie

MARK-I o

e



Ψ' (e



ATLAS
EXPERIMENT

Run Number: 201289, Event Number: 24151616
Date: 2012-04-16 16:53:58 CEST

nd D0



PARTICLE DETECTION


■ What is a particle


- Irreducible representation of the Poincaré group (Wigner)

■ What is a detector

- an instrument to measure the all properties (E, p, m, lifetime, quantum numbers,...) of a particle and identifying it


Elementary particles









π^+


Mesons and Baryons



Proton




Particle Physics Timeline



Today > 200 particles listed in PDG:

- 27 have $c\tau > 1 \mu\text{m}$
 - they can be seen as tracks in a detectors
- 13 have $c\tau < 500 \mu\text{m}$
 - displaced vertices



NOBEL PRIZES FOR INSTRUMENTATION

[http://www.lhc-closer.es/
php/index.php?
i=1&s=9&p=2&e=0](http://www.lhc-closer.es/php/index.php?i=1&s=9&p=2&e=0)



1927: C.T.R. Wilson, Cloud Chamber



1939: E. O. Lawrence, Cyclotron



1948: P.M.S. Blacket, Cloud Chamber



1950: C. Powell, Photographic Method




1954: Walter Bothe, Coincidence method



1960: Donald Glaser, Bubble Chamber



1968: L. Alvarez, Hydrogen Bubble Chamber



1992: Georges Charpak, Multi Wire Proportional Chamber

Particle discoveries

By 1959: 20 particles

e⁻ : fluorescent screen

n : ionization chamber

7 Cloud Chamber:

e⁺

μ^+ , μ^-

K⁰

Λ^0

Ξ^-

Σ^-

6 Nuclear Emulsion:

π^+ , π^-

anti- Λ^0

Σ^+

K⁺, K⁻

2 Bubble Chamber:

Ξ^0

Σ^0

3 with Electronic techniques:

anti-n

anti-p

π^0


Imaging Detectors: The Cloud Chamber

- The **cloud chamber** contains a supersaturated vapor of water or alcohol.
 - A charged particle interacting with the mixture, creates ions.
 - The ions act as condensation nuclei around which a mist will form
- High energy alpha and beta particles leave a track due to the ions they produce along their path
- If a magnetic field is applied positively and negatively charged particles will curve in opposite directions.




The positron

- Positron discovery, Carl Anderson 1933 [Nobel price 1936]
- Magnetic field 15000 Gauss, chamber diameter 15cm.
- A 63 MeV positron passes through a 6mm lead leaving the plate with energy 23MeV.
- The ionization of this particle, and its behavior in passing through the foil was the same as those of an electron but with positive charge




The muon

- Muons were discovered by Carl D. Anderson and Seth Neddermeyer at Caltech in 1936 with a cloud chamber while studying cosmic radiation




"The other double trace of the same type (figure 5) shows closely together the thin trace of an electron of 37 MeV, and a much more strongly ionizing positive particle with a much larger bending radius. The nature of this particle is unknown; for a proton it does not ionize enough and for a positive electron the ionization is too strong. The present double trace is probably a segment from a "shower" of particles as they have been observed by Blackett and Occhialini, i.e. the result of a nuclear explosion".

Kunze, P., Z. Phys. 83, (1933) 1




The muon and relativity



- Muons produced in the upper atmosphere ≈ 10 km
- Lifetime $\tau_{\mu} = 2.2 \times 10^{-6}$ s
 - $s = v\tau \approx 600$ m
 - Therefore no muon should reach Earth!
- But we see them
 - Relativity $\gamma = E/m_{\mu}c^2$
 - $s = v\gamma\tau = 12.5$ km
- Pions: $\tau_{\pi} = 2.6 \times 10^{-8}$ s,
 - $s = v\gamma\tau = 115$ m

The Pion

- The pion was discovered in Nuclear emulsion techniques, Powell 1947; Nobel Prize 1950
- Discovered in 1947 in nuclear emulsions exposed to cosmic rays, and they showed that it decay to a muon and an unseen partner.
- The constant range of the decay muon from the pion decay indicate that this is a two body decay





$$\pi^- \rightarrow \mu^- \bar{\nu}_\mu$$

$$\mu^- \rightarrow e^- \bar{\nu}_e \nu_\mu$$

The kinematics of two body decays

- Relativistic Kinematics



$$P_\pi = (m_\pi, 0) \quad P_\mu = (E_\mu, \vec{p}_\mu) \quad P_\nu = (E_\nu, \vec{p}_\nu)$$

$$P_\pi = P_\mu + P_\nu$$

$$(P_\pi - P_\nu)^2 = (P_\mu)^2$$

$$m_\nu \approx 0$$

$$P_\pi^2 + P_\nu^2 - P_\pi \cdot P_\nu = m_\mu^2$$

$$m_\pi \approx 140 \text{ MeV}$$

$$m_\pi^2 + 0 - 2m_\pi \cdot E_\nu = m_\mu^2$$


$$m_\mu \approx 106 \text{ MeV}$$

$$E_\nu = \frac{m_\pi^2 - m_\mu^2}{2m_\pi} \approx 30 \text{ MeV}$$

$$|\vec{p}_\mu| \approx 30 \text{ MeV} \quad E = \sqrt{\vec{p}_\mu^2 + m_\mu^2} = 110 \text{ MeV}$$

In the pion rest frame, the muon has
KE = 110 MeV - 106 MeV = 4 MeV



3-body kinematics: the neutrino



- Neutrinos do not carry any charge → take part in the electroweak and gravitational interaction → neutrinos hardly interact with matter
- Existence of neutrinos was inferred from studies of the lepton spectrum in beta decays.




"I have done a terrible thing. I invented a particle that cannot be detected."
W. Pauli



Single beta decay energy spectrum. The observed spectrum is continuous and not at a constant energy as was initially expected. [D. Stewart]




Imaging techniques: Nuclear Emulsions

- A nuclear emulsion plate is a photographic plate with a thick emulsion layer and uniform grain size.
 - It records the tracks of charged particles passing through
 - It produce a cumulative record
 - The plates must be developed before the tracks can be observed.
- In 1937, Marietta Blau and Hertha Wambacher discovered nuclear *disintegration stars* due to spallation in nuclear emulsions exposed to cosmic radiation at a height of 2,300 meters above sea level.



The discovery of the Kaon

- First evidence of the decay of the Kaon into 3 Pions was found in 1949 in Nuclear emulsion




Imaging Detectors: the Bubble chamber

Invented in 1952 by Glaser
(1960 Nobel Prize in Physics)


- A **bubble chamber** is a vessel filled with a superheated transparent liquid (Hydrogen at $T=30\text{K}$). A charge particle initiate boiling.
- The size of the chambers grew quickly:
 - 1954: 2.5' (6.4cm)
 - 1954: 4' (10cm)
 - 1956: 10' (25cm)
 - 1959: 72' (183cm)
 - 1963: 80' (203cm)
 - 1973: 370cm

- Some disadvantages:
 - It cannot be triggered
 - Low rate capability
 - The photographic readout: for data analysis one had to look through millions of photos





- Urban history: Glaser was inspired by the bubbles in a glass of beer
- In a 2006 talk he did experiments using beer to fill early prototypes.

Discovery of Omega




The 80-inch Bubble Chamber



BNL, First Pictures 1963, 0.03s cycle



$\Omega^- = sss$ Confirmation of the quark model

The discovery of neutral currents




- Gargamelle, a very large heavy-liquid (freon) chamber constructed at Ecole Polytechnique in Paris, came to CERN in 1970.
- 2 m in diameter, 4 m long and filled with Freon at 20 atm, in 2 T B field
- Gargamelle in 1973 was the tool that permitted the discovery of neutral currents.

Can be seen outside the
Microcosm Exhibition



Electronics detectors

- In the 70ies the logic (electronic) detectors took over
 - Geiger counters
 - Scintillator + photomultipliers
 - Spark counters
- The particle is not “seen” but its nature and existence “deduced” via a logic experiment (coincidences, triggers, detection of decay products ...)





Scintillating Screen:
Rutherford Experiment 1911, Zinc Sulfide screen was used as detector. If an alpha particle hits the screen, a flash can be seen through the microscope.

Spark Chamber

The Spark Chamber was developed in the early 60ies.


Schwartz, Steinberger and Lederman used it in discovery of the muon neutrino




- A charged particle traverses the detector and leaves an ionization trail.
- The scintillators trigger an HV pulse between the metal plates and sparks form in the place where the ionization took place.

Multi-wire proportional chambers


Tube, Geiger- Müller, 1928



Multi Wire Geometry, in H. Friedmann 1949





G. Charpak 1968, Multi Wire Proportional Chamber, readout of individual wires and proportional mode working point.



A charged particle traversing the detector leaves a trail of electrons and ions. The wires are on positive HV. The electrons drift to the wires in the electric field and start to form an avalanche in the high electric field close to the wire. This induces a signal on the wire which can be read out by an amplifier.

The merge of electronic Images

Discover of the W and Z (1983)
Rubbia & Van der Meer, Nobel Prize 1984



This computer reconstruction shows the tracks of charged particles from the proton-antiproton collision. The two white tracks reveal the Z's decay. They are the tracks of a high-energy electron and positron.

Particle Detection and stable particles

- The most frequent “stable” particles ($c\tau > 500 \mu\text{m}$) are: Electrons (e), muons (μ), photons (γ), pions (π), kaons (K), protons (p) and neutrons (n)
- A particle detector must be able to identify and measure at least the energy & momentum of these 8 particles.


EM Interactions: Electrons ($m_e=0.5 \text{ MeV}$), muons ($m_\mu=105.7 \text{ MeV}$), photons ($m_\gamma=0$)

EM & Strong interaction: pions ($m_\pi=139.6 \text{ MeV}$), Charged kaons ($m_K+=493.7 \text{ MeV}$), protons ($m(p)=938.3 \text{ MeV}$)

Strong interaction: Neutral K^0 ($m_K = 497.7 \text{ MeV}$) and n ($m_n=939.6 \text{ MeV}$)

The difference in Mass, Charge, Interaction is the key to particle identification

Unstable Particles





$$m_{inv,x}^2 = (E_1 + E_2)^2 - (p_1 + p_2)^2$$




$$E_x = E_1 + E_2$$

$$\vec{p}_x = \vec{p}_1 + \vec{p}_2$$


$$J/\psi \rightarrow \mu^+ \mu^-$$



Quark confinement and Jets



Particle Identification



- Electrons ionize and show bremsstrahlung due to the small mass
- Photons do not ionize but undergo pair production in high Z material
- Charge hadrons ionize and produce hadron shower in dense material
- Neutral hadrons do not ionize but produce hadron shower in dense material

The advent of silicon

The measurement of the lifetime of c and b quark

Event: 7030744 Run: 142820 EventType: DATA | Unpresc: 0,1,33,34,35,4,36,7,8,9,10,42,11,44,13,45,14,15,17,49,20,21,23,24

Missing Et
Et=85.7 phi=4.5




List of tracks

Id	pt	phi	eta
Cdf tracks: first 5			
468	23.5	-0.7	-0.3
439	25.9	2.9	-0.4
489	21.2	1.6	-0.3
469	16.4	0.9	0.5
500	8.7	0.7	-0.1


To select track type
SelectCdfTrack(Id)

Svt	pt	phi	eta
Svt tracks: first 5			
0	180.8	1.2	
2	45.2	3.5	
1	-15.1	3.2	
5	9.5	2.4	
3	1.4	4.6	

To select track type
SelectSvtTrack(Id)



Tracking and vertex detectors were crucial to the discovery of top




Detector Optimization

- Which kind of “particle” we have to detect?
- What is the required dimension of the detector?
- Which “property” of the particle we have to know?
 - Position
 - Lifetime
 - Quantum numbers
 - Energy
 - Charge
- What is the maximum count rate?
- What is the “time distribution” of the events?
- What is the required resolution ?
- What is the dead time?

Measurement Resolution

- Resolution generally defined as 1 standard deviation (1σ) for a Gaussian distribution, or the FWHM (Δz)
 - For a Gaussian $\sigma = \text{FWHM}/2.36$
- If the measurement is dominated by Poisson fluctuations



$$\frac{\sigma_z}{\langle z \rangle} = \frac{\sqrt{N}}{N} = \frac{1}{\sqrt{N}}$$




What if the distribution is not Gaussian ?

- Box distribution: $\sigma_z = \Delta z / \sqrt{12}$
- Other distributions: RMS

$$\sigma^2 = \frac{\int_{-\frac{p}{2}}^{\frac{p}{2}} (x_r - x_m)^2 D(x_r) dx_r}{\int_{-\frac{p}{2}}^{\frac{p}{2}} D(x_r) dx_r} = \frac{p^2}{12}$$


$D(x) = 1$ uniform distribution of tracks
 $X_m = 0$ pixel centre

Configuration of HEP Detectors




Multipurpose Detectors



Particle flow reconstruction



Particle Flow and Missing E_T





Particle Flow: reconstruct stable particles in the event (e , μ , γ & charged and neutral hadrons) from the information from all sub-detectors, to optimize the determination of particle types, directions and energies



- At the LHC an unknown proportion of the energy of the colliding protons escapes down the beam-pipe.
- If invisible particles (neutrinos, neutralinos ?) are created their momentum can be constrained in the plane transverse to the beam direction

$$E_T^{\text{miss}} = - \sum_i p_T(i)$$

Application of HEP Detectors: PET



Useful material & acknowledgments

- I have taken part of the content of these lecture from Werner Riegler's summer student lectures in 2011 and Erika Garutti's DESY lecture notes
- Useful books
 - Detector for particle radiation, Konrad Kleinknecht
 - Techniques for Nuclear and Particle Physics Experiments, W. R. Leo
 - Particle Detectors, Claus Grupen
 - Introduction to Experimental Particle Physics, R. Fernow
 - The Physics of Particle Detectors, D. Green
 - Review in data particle book on Passage of particles through matter
 - Review in data particle book on Particle Detectors at accelerators

Historical legacies

- CDF pioneered the silicon vertex detector in the hadron collider environment and pioneered the silicon vertex trigger separating b -hadrons
- CDF and D0 developed multi-level triggering with fast microprocessor farms to select interesting events.
- Tracker information at L2 triggering with associative memories



CDF's first Silicon Vertex Detector at the Smithsonian Museum, Washington

Many of these advances have been adopted and improved for the LHC experiments