

# CTEQ NNLO PDFs, heavy flavors, and LHC data

Pavel Nadolsky

Southern Methodist University  
Dallas, TX, U.S.A.

*in collaboration with*  
*M. Guzzi, J. Gao, J. Huston, H.-L. Lai, Z. Li,*  
*Z. Liang, J. Pumplin, D. Soper, D. Stump, C.-P. Yuan*

May 23, 2012

# Two sets of CT NNLO error PDFs

## 1. CT10 NNLO eigenvector set

Available at [http://hep.pa.msu.edu/cteq/public/ct10\\_2012.html](http://hep.pa.msu.edu/cteq/public/ct10_2012.html); is being submitted to LHAPDF; main focus of this talk

Complements the CT10/CT10W NLO PDF sets (*Lai et al., PRD82, 074024 (2010)*)

- Includes only “pre-LHC” CT10 data. Can be used to predict LHC cross sections based on pre-LHC experimental inputs
- Same input parameters, functional forms for input PDFs as in the CT10 NLO PDFs
  - ▶  $\alpha_s(M_Z) = 0.118 \pm 0.002$ ,  $m_c^{pole} = 1.3$  GeV,  $m_b^{pole} = 4.75$  GeV
  - ▶ Simpler assumptions about the PDF flavor composition at  $\mu_0 = m_c^{pole} = 1.3$  GeV, e.g.,  $\bar{u}(x)/\bar{d}(x) \rightarrow 1$  as  $x \rightarrow 0$
- Updated  $N_f = 3$  and 4 NLO sets. The world-average  $\alpha_s(Q, N_f = 5)$  is converted to  $\alpha_s(Q, N_f \neq 5)$  at  $Q = m_\tau$  or  $M_Z$  and evolved with  $N_f \neq 5$  to other  $Q$  values




# Two sets of CT NNLO error PDFs

## 2. CT12 NLO and NNLO eigenvector sets

### To be released within a few months

- Include LHC W and Z rapidity data, ATLAS and CMS jet data, HERA'2011  $F_L$  data
- Updated  $\alpha_s, m_c, m_b$  values
- Flexible  $\bar{d}/\bar{u}$  ratio at  $x \rightarrow 1$ , updated  $(s + \bar{s})/(\bar{u} + \bar{d})$  at  $x \lesssim 10^{-2}$ 
  - ▶ Constrained by the LHC  $W/Z$  rapidity distributions

# CT10 NNLO error PDFs (compared to CT10W NLO)



## CT10 PDF sets: the naming conventions

- **Two NLO PDF sets**, without/with Tevatron Run-2 data on  $W$  charge asymmetry  $A_\ell$

CT10 NLO does not include  
CT10W NLO includes 4  $p_{T\ell}$  bins of D0 Run-2  $A_\ell$  data


⇒ CT10 and CT10W sets differ mainly in the behavior of  $d(x, Q)/u(x, Q)$  at  $x > 0.1$

- **One NNLO PDF set:** only 2 inclusive  $p_{T\ell}$  bins of D0 Run-2  $A_\ell$  data are included that have smallest theory uncertainties
- **The NNLO set is a counterpart of both CT10 NLO and CT10W NLO.** It uses only a part of the  $A_\ell$  data sample that distinguishes between CT10 NLO and CT10W NLO.

## Striving for NNLO accuracy in the PDFs


- So far, only “partial NNLO” global fits exist. For some fitted processes (inclusive jet production, CC DIS with  $m_q \neq 0$ ), QCD contributions are known only to NLO. NLO EW contributions, power corrections, other systematic errors may be comparable to NNLO QCD effects.
- CT10 “NNLO” PDFs underwent validation studies for about one year. We identified several types of uncertainties that compete with NNLO QCD contributions.
- CT10 NNLO and NLO PDFs produce about the same  $\chi^2/N_{pt} \approx 1.05 - 1.10$  for  $N_{pt} = 2700$  data points
- Shapes of the NNLO PDFs have noticeably evolved compared to NLO as a result of  $\mathcal{O}(\alpha_s^2)$  contributions, updated electroweak contributions, revised statistical procedures

## CT10 NNLO central PDFs, as ratios to NLO, $Q=2$ GeV



1. At  $x < 10^{-2}$ ,  $\mathcal{O}(\alpha_s^2)$  evolution suppresses  $g(x, Q)$ , increases  $q(x, Q)$
2.  $c(x, Q)$  and  $b(x, Q)$  change as a result of the  $\mathcal{O}(\alpha_s^2)$  GM VFN scheme
3. At  $x > 0.1$ ,  $g(x, Q)$  and  $d(x, Q)$  are reduced by revised EW couplings, alternative treatment of correlated systematic errors, scale choices




# CT10 NNLO central PDFs, as ratios to NLO, $Q=85$ GeV






# CT12 NLO predictions for LHC jet production

ATLAS single-inclusive jet production (*arXiv:1112.6297*); FastNLO 2;  $R=0.6$ ;  
 $\chi^2/N_{d.o.f} = 0.72 (0.98)$  for CT12 NLO (CT10 NLO)



CT10 NNLO and CT12 PDFs (black lines) predict smaller jet cross sections at large  $p_T$ , as a result of reduced  $g(x, Q)$  at  $x > 0.1$

# CT10 NNLO PDFs compared to MSTW NNLO




1. CT10 gluon and quarks are harder at  $x \rightarrow 0$ ;

$$g(x, Q_0) > 0 \text{ at } 10^{-5} \leq x \leq 1$$

2. The CT10 strange PDF is larger at  $x \sim 10^{-3}$

# Predictions for production of electroweak bosons

## NNLO cross sections



## Details of the CT10 NNLO computation


- **NNLO hard-scattering contributions in DIS** (in the S-ACOT- $\chi$  mass scheme) and **vector boson production** (NNLO  $K$  factors from FEWZ for  $d\sigma/dy$ ; NNLL/NLO+K from ResBos for  $W$  charge asymmetry)
- NNLO evolution for  $\alpha_s$  and PDFs (HOPPET)
  - ▶ matching coefficients relating the PDFs in  $N_f$  and  $N_{f+1}$  schemes (*Smith, van Neerven, et al.*)
- **Pole quark masses or  $\overline{MS}$  quark masses as an input**
  - ▶ CT10 NNLO: pole masses  $m_c = 1.3$  GeV,  $m_b = 4.75$  GeV

# Neutral-current DIS in S-ACOT- $\chi$ scheme at NNLO

M. Guzzi, P.N., H.-L. Lai, C.-P. Yuan, arXiv:1108.5112 (hep-ph)

## Objectives


- elucidate fundamental principles that a viable GM scheme must satisfy
- modify the QCD factorization theorem for DIS with massive quarks (*Collins, 1998*) to satisfy momentum conservation in all heavy-quark scattering channels
- provide algorithmic implementation of NNLO massive contributions, in close analogy to the ZM scheme



NNLO scattering contributions

# S-ACOT- $\chi$ scheme: merging FFN and ZM

$x=0.01$



Les Houches toy PDFs, evolved at NNLO with threshold matching terms

$\mathcal{O}(\alpha_s^2)$  flavor-creation contributions with  $m_c \neq 0$  are included exactly (based on the calculation by Riemersma, Smith, van Neerven, PL B347, 143 (1995))

The implementation will be made available in HERA FITTER

A complementary calculation (a “hybrid mass scheme”; exact  $\mathcal{O}(\alpha_s)$  massive ACOT terms + approximate  $\mathcal{O}(\alpha_s^2)$  and  $\mathcal{O}(\alpha_s^3)$  massive terms) has been published by Stavreva, Olness, Schienbein, et al., arXiv:1203.0282

# Efforts to reduce differences between NNLO fits

## 1. Benchmarking comparisons of the fitting codes

1.1 Benchmarks for inclusive DIS cross sections  
(with *S. Alekhin, A. Glazov, A. Guffanti, J. Rojo*)

⇒ importance of subleading/NLO EW effects

1.2. Benchmarks for NLO jet production calculations  
(*J. Gao, Z. Liang at SMU + J. Rojo representing ApplGrid*)

⇒ effect on the large- $x$  gluon PDF

2. Re-examination of correlated systematic uncertainties from both theory and experiment (*cf. also Jun Gao's talk*)

PDF4LHC is an ideal venue to complete these efforts, extend to NuTeV dimuon cross sections, etc.

## 1.1. Benchmarking of $\gamma^*Z$ interference of NC DIS cross sections

CTEQ implementation of the  $\gamma^*Z$  interference terms in NC DIS is based on the helicity formalism (*Aivazis, Olness, Tung, 1994*). It shared definitions of EW couplings with vector boson production.


During the benchmarking, we modified the  $Z$  boson couplings to comply with the formulas for NC DIS in the combined HERA analysis (*arXiv:0911.0884*).

The revised CTEQ code was compared at ZM LO against another code (*A. Guffanti*). Excellent agreement was reached.

**This modification reduced  $\bar{q}(x, Q)$ ,  $g(x, Q)$  at  $x > 0.1$  in CT10 NNLO. Subleading EW contributions are relevant in this case.**



## 1.1. Benchmarking of NC DIS cross sections




Point-by-point contributions to  $\chi^2$  for the combined HERA NC DIS set at ZM LO in the CTEQ code and alternative code (A. Guffanti). Some differences between two codes were observed at large  $Q$ , in the  $(x, Q)$  region where experimental errors also increase. Differences become practically zero after the benchmarking.

## 1.2 Benchmark comparison of NLO jet cross sections

J. Gao, Z. Liang, P. N., in 2011 Les Houches Proceedings, arXiv:1203.6803;  
in collaboration with D. E. Soper, H.-L. Lai, C.-P. Yuan

Benchmarking of NLO cross sections for inclusive jet and dijet production is needed to understand theory uncertainties at the LHC (including significant scale dependence at largest  $p_T^{jet}$ )

It is also a preparation step toward the NNLO calculation for jet production



## Modified EKS program *(publicly available)*

CTEQ fits have been using the EKS program, which realizes an NLO calculation for single-incl. jet and dijet production by S. D. Ellis, Z. Kunszt, and D. E. Soper *(PRL 69, 1496 (1992))*



We modified the EKS code to calculate NLO jet cross sections more efficiently, with flexible input and output formats, and provide them in “almost differential” finely binned tables.

Other available programs include NLOJET++ *(Z. Nagy, PRL 88, 122003 (2002))* , FastNLO *(Kluge et. al., hep-ph/0609285)*, POWHEG *(Alioli et. al., JHEP 04081 (2011))*, ApplGrid *(EPJC66, 503 (2010))*



We identified specific conditions needed to reconcile MEKS and FastNLO outputs. For very specific settings of the jet algorithm, recombination scheme, jet trigger, QCD scale choices, MEKS and FastNLO show excellent agreement at most  $y^{jet}$ ,  $p_T^{jet}$ , and  $M^{jj}$ .

# Comparison of MEKS and FastNLO 1.0

**Dijet production:** excellent agreement at both the Tevatron and LHC



**Single-inclusive jet production:** discrepancies of 3-10% exist at large  $p_T$ , possibly due to different definitions of the "jet  $p_T$ " used as the QCD scale.




## The choice of scale in jet production cross sections

The “jet  $p_T$ ” may refer to the “ $p_T$  of the leading jet in an event”, “ $p_T$  of each jet in each  $p_T$  bin”, “average  $p_T$  in each  $p_T$  bin (FastNLO 1)”, or “ $p_T$  of the leading jet in each bin (ATLAS)”. Differences resulting from these definitions are comparable to NNLO/PDF uncertainties.

CT10 NNLO/CT12 PDFs use  $\mu_F = \mu_R = \langle p_T \rangle_{bin}$  and FastNLO 2 (implemented as an alternative to the  $K$ -factor lookup tables based on the MEKS calculation).

⇒ Softer gluon than in CT10 NLO



## 2. Computation of correlated systematic errors


$$\chi^2 = \sum_{\{\text{exp.}\}} \left[ \sum_{k=1}^{N_{pts}} \frac{1}{S_k^2} \left( D_k - T_k(\{a\}) - \sum_{\alpha=1}^{N_\lambda} \lambda_\alpha \beta_{k\alpha} \right)^2 + \sum_{\alpha=1}^{K_e} \lambda_\alpha^2 \right]$$

The experimental correlated systematic errors  $\beta_{k\alpha}$  are often published as percentages. It can be taken to be a percentage of the theoretical prediction  $T_k$  ("truth") or the experimental datum  $D_k$ .


- 1. Percentage of  $T_k$ :** results in smooth  $\beta_{k\alpha}$  :-); may depend on the theoretical model :-(-
- 2. Percentage of  $D_k$ :**  $\beta_{k\alpha}$  is deduced from the measured data :-), but may not be smooth due to statistical fluctuations :-(-

The methods are equivalent if  $T_k$  is close to  $D_k$ . In the actual CTXX fits to the Tevatron Run-2 jet data, **method 1** (used in pre-2012 CTEQ fits) results in a harder gluon at  $x > 0.1$  than **in method 2**. We use **method 2** in the latest NNLO fits.

## 2.2. Impact on the best fit NLO PDFs




**pc317l:** CT12 NLO candidate obtained with method 1



**pc317n:** CT12 NLO candidate obtained with method 2


Notice changes in  $u(x, Q)$ ,  $d(x, Q)$ ,  $g(x, Q)$

# Strangeness in CT12 PDFs and LHC $W/Z$ cross sections



In 2008, our CTEQ6.6 PDF correlation analysis pointed out the sensitivity of ratios  $\sigma_W/\sigma_Z$  at the LHC to the strangeness PDF, with implications to EW precision measurements (P.N., Lai, Cao, Huston, Pumplin, Tung, Yuan, PRD, 78 (2008) 013004).



The ATLAS analysis (arXiv:1203.4051) of  $W$  and  $Z$  production suggests that  $\bar{s}(x, Q)/\bar{d}(x, Q) = 1.00^{+0.25}_{-0.28}$  at  $x = 0.023$  and  $Q^2 = 1.9 \text{ GeV}^2$



What is the impact of the new LHC  $W$  and  $Z$  data on the CT12 PDFs that will include them?



# Small- $x$ limits of $\bar{d}(x, Q)/\bar{u}(x, Q)$ and $\bar{s}(x, Q)/\bar{u}(x, Q)$ in the CT12 analysis (PRELIMINARY)



The CT12 analysis explores the possibility of  $\lim_{x \rightarrow 0} \bar{d}/\bar{u} \neq 1$ . Some “unbiased” CT12 candidate fits have  $\bar{s}(x, Q)/\bar{u}(x, Q) > 1$  at  $x < 10^{-3}$ .

We would like to better understand the flavor decomposition at small  $x$  before releasing the CT12 PDFs.

## Conclusions

- The CT10 NNLO PDF analysis (based on pre-LHC data only) is released. It is based on a new streamlined implementation of heavy-quark DIS contributions at two loops (*Guzzi et al.*, *arXiv:1108.5112*).
- The CT12 NLO and NNLO analysis (in progress) will include latest LHC data on  $W$ ,  $Z$ , and jet production. Possible impact on  $SU(3)$  properties of quark sea at  $x < 10^{-3}$ .
- Several factors that are comparable to NNLO contributions (treatment of percentage corr. syst. errors, choices of scales, electroweak radiative contributions, ...) have been thoroughly examined in this analysis
- We use a specific choice to evaluate these factors in the CT12 (N)NLO fits. The uncertainty associated with this choice need to be examined in the future

# Backup slides


# Neutral-current DIS in a general-mass scheme at NNLO

M. Guzzi, P.N., H.-L. Lai, C.-P. Yuan, arXiv:1108.5112 (hep-ph)

## Objectives

■ The CT10 fit computes  $c$ ,  $b$  quark contributions to NC DIS in the S-ACOT- $\chi$  general-mass factorization scheme (Aivasis, Collins, Olness, Tung, 1994; Collins, 1998; Kramer, Olness, Soper, Tung, Kretzer, Schmidt)

■ We have realized this scheme at NNLO. We have also demonstrated how to derive this scheme (including kinematic rescaling of heavy-quark scattering terms at the mass threshold) from the QCD factorization theorem by Collins




NNLO scattering contributions

## Massive quark contributions to neutral-current DIS...

...affect predictions for the LHC  $W$  and  $Z$  cross sections (Tung et al., hep-ph/0611254)

$W^\pm$  &  $Z$  cross sections at the LHC (14 TeV)



### Extensive recent work

Tung et al., hep-ph/0611254; Thorne, hep-ph/0601245; Tung, Thorne, arXiv:0809.0714; PN., Tung, arXiv:0903.2667; Forte et al., arXiv:1001.2312; J. Rojo et al., arXiv:1003.1241; Alekhin, Moch, arXiv:1011.5790;...

### Several heavy-quark factorization schemes

FFN, ACOT, BMSN, CSN, FONLL, TR'...

The NNLO realization of the S-ACOT- $\chi$  factorization scheme combines benefits of several approaches

## Main features of the S-ACOT- $\chi$ scheme

- It is proved to all orders by the QCD factorization theorem for DIS (*Collins, 1998*)
- **Universal PDFs**
- It is relatively simple
  - ▶ One value of  $N_f$  (and one PDF set) in each  $Q$  range
  - ▶ sets  $m_h = 0$  in ME with incoming  $h = c$  or  $b$
  - ▶ matching to FFN is **implemented at the level of the QCD factorization theorem**
- It reduces to the ZM  $\overline{MS}$  scheme at  $Q^2 \gg m_Q^2$ , without additional renormalization
- It reduces to the FFN scheme at  $Q^2 \approx m_Q^2$ 
  - ▶ has reduced dependence on tunable parameters at NNLO



## Components of inclusive $F_{2,L}(x, Q)$

S-ACOT- $\chi$  NNLO expressions are reminiscent of the ZM scheme (e.g., in Moch, Vermaseren, Vogt, 2005), with all components available from literature

Components of inclusive  $F_{2,L}(x, Q^2)$  are classified according to the quark couplings to the photon

$$F = \sum_{l=1}^{N_l} F_l + F_h \quad (1)$$

$$F_l = e_l^2 \sum_a [C_{l,a} \otimes f_{a/p}] (x, Q), \quad F_h = e_h^2 \sum_a [C_{h,a} \otimes f_{a/p}] (x, Q). \quad (2)$$




At

$$F_h^{(2)} = e_h^2 \left\{ c_{h,h}^{NS,(2)} \otimes (f_{h/p} + f_{\bar{h}/p}) + C_{h,l}^{(2)} \otimes \Sigma + C_{h,g}^{(2)} \otimes f_{g/p} \right\}$$

$\mathcal{O}(\alpha_s^2)$ :

$$F_l^{(2)} = e_l^2 \left\{ C_{l,l}^{NS,(2)} \otimes (f_{l/p} + f_{\bar{l}/p}) + c^{PS,(2)} \otimes \Sigma + c_{l,g}^{(2)} \otimes f_{g/p} \right\}. \quad (3)$$


## Rescaling to all orders of $\alpha_s$ and the factorization theorem



We show that a minor modification of the QCD factorization theorem (Collins, 1998)...

- enables suppression of charm production at  $Q^2 \rightarrow m_{c,b}^2$  in all channels and at each  $\alpha_s$  order without extra smoothness conditions or damping factors


- preserves universality of heavy-quark PDFs






## CT12 predictions for ATLAS jet production (2)

ATLAS single-inclusive jet production (*arXiv:1112.6297*); FastNLO 2;  $R=0.4$ ;  
 $\chi^2/N_{d.o.f} = 0.76 (0.95)$  for CT12 NLO (CT10 NLO)



# CT12 predictions for ATLAS jet production (3)



## Factorization scale in NLO jet cross sections

NLO jet cross sections depend significantly on renorm. and fact. scales,  $\mu_F$  and  $\mu_R$

CT10 fit assumed the default scale  $\mu_F^0 = \mu_R^0 = p_T/2$ ; other groups and experimentalists often use  $\mu_F^0 = \mu_R^0 = p_T$





Trade-offs between the scale choices (see the next slide)

- $\mu_{F,R} = p_T/2$ :  $K \approx 1$  at small  $y_{jet}$ , large scale dependence at large  $y_{jet}$
- $\mu_{F,R} = p_T$ :  $K \approx 1.4$  at small  $y_{jet}$ , smaller scale dependence at large  $y_{jet}$

CT10 NNLO will provide a PDF set for  $\mu_{F,R} = p_T$

# NLO corrections for $\mu_{F,R}^0 = p_T/2$ (left) and $p_T$ (right)

Jun Gao, 2011



Scale dependence (green) corresponds to variations  $1/2 \leq \mu_{F,R}/\mu_{F,R}^0 \leq 2$ . Red bands reflect the PDF uncertainty in the lookup tables for the NLO K-factors.