



Aspects of saturation and final states

Krzysztof Kutak






Based on:

JHEP 1202 (2012) 117 K.K, K. Golec-Biernat, S. Jadach, M. Skrzypek

arXiv:1205.5035 K.K, Sebastian Sapeta

LHC as a scanner of gluon

$$S = 2P_1 \cdot P_2$$



$$x_1 = \frac{1}{\sqrt{S}} (p_{t1} e^{y_1} + p_{t2} e^{y_2}) \quad \xrightarrow{y_1 \sim 0, y_2 \gg 0} \quad \sim 1$$

$$x_2 = \frac{1}{\sqrt{S}} (p_{t1} e^{-y_1} + p_{t2} e^{-y_2}) \quad \ll 1$$

High energy limit of QCD


$$\frac{d\sigma}{dy_1 dy_2 d^2p_{1t} d^2p_{2t}} = \sum_{a,b,c,d} \int \frac{d^2k_{1t}}{\pi} \frac{d^2k_{2t}}{\pi} \frac{1}{16\pi^2(x_1 x_2 S)^2} |\overline{\mathcal{M}}_{ab \rightarrow cd}|^2 \delta^2(\vec{k}_{1t} + \vec{k}_{2t} - \vec{p}_{1t} - \vec{p}_{2t})$$

$$\times \phi_{a/A}(x_1, k_{1t}^2, \mu^2) \phi_{b/B}(x_2, k_{2t}^2, \mu^2) \frac{1}{1 + \delta_{cd}}$$

Gribov, Levin, Ryskin' 81
Ciafaloni, Catani, Hautman '93

Implemented in Monte Carlo generator CASCADE
H. Jung

- Gluon density depends on k_t
 - Off shell initial state partons with shellness $\sim k_t$




High energy prescription and forward-central dijets

Deak, Jung, Hautmann Kutak
JHEP 0909:121,2009

$$\frac{d\sigma}{dy_1 dy_2 dp_{1t} dp_{2t} d\phi} = \sum_{a,c,d} \frac{p_{t1} p_{t2}}{8\pi^2 (x_1 x_2 S)^2} |\overline{\mathcal{M}}_{ag \rightarrow cd}|^2 x_1 f_{a/A}(x_1, \mu^2) \phi_{g/B}(x_2, k_t^2, \mu^2) \frac{1}{1 + \delta_{cd}}$$

$$S = 2P_1 \cdot P_2$$



- Resummation of logs of x and logs of hard scale
- Knowing well parton densities at large x one can get information about low x physics
- Framework goes recently under name “ k_T framework”






$$x_1 = \frac{1}{\sqrt{S}} (p_{t1} e^{y_1} + p_{t2} e^{y_2}) \quad \xrightarrow{y_1 \sim 0, y_2 \gg 0} \sim 1$$

$$x_2 = \frac{1}{\sqrt{S}} (p_{t1} e^{-y_1} + p_{t2} e^{-y_2}) \quad \ll 1$$

$$k_1^\mu = x_1 P_1^\mu$$

$$k_2^\mu = x_2 P_2^\mu + k_t^\mu$$

Forward central – jet production



- HEJ and Cascade based on unordered in k_t emissions but use different parton densities
- Herwig and PYTHIA use k_t ordered shower but differ in approximations in ME and ordering conditions in shower

Deak, Jung, Hautmann, Kutak, '10

Observable sensitive to **saturation**
 Khazeev, Levin, McLerran, '05; Marquet 2007

High energy factorization and saturation


Saturation – state where number of gluons stops growing due to high occupation number.

More generally saturation is an example of **percolation** which has to happen since partons have size $1/k_t$ and hadron has finite size

Cross sections change their behavior from power like to **logarithmic like**.


On microscopic level it means that gluon apart splitting recombine

splitting




recombination

Nonlinear evolution equations
BK, JIMWLK
CGC framework
DIPSY



Linear evolution equation




Bartels, Wusthoff

Z.Phys. C66 (1995)
157-180

Chirilli, Szymanowski, Wallon '10

Forward physics as the way to constrain gluon both at large and small p_t

- Too flat behaviour of σ at large k_t
- Lack of saturation in CCFM small k_t



Needed framework which unifies both correct behaviors

Gluon density and saturation

Go to momentum space

dipole density



$$\Phi(x, k^2) = \int \frac{d^2\mathbf{b}d^2\mathbf{r}}{2\pi} \exp(-i\mathbf{k} \cdot \mathbf{r}) \frac{N(x, r, b)}{r^2}$$

$$\Phi(x, k^2) = \Phi_0(x, k^2) + \bar{\alpha}_s \int_{x/x_0}^1 \frac{dz}{z} \int_0^\infty \frac{dl^2}{l^2} \left[\frac{l^2\Phi(x/z, l^2) - k^2\Phi(x/z, k^2)}{|k^2 - l^2|} + \frac{k^2\Phi(x/z, k^2)}{\sqrt{(4l^4 + k^4)}} \right] - \frac{\bar{\alpha}_s}{\pi R^2} \int_{x/x_0}^1 \frac{dz}{z} \Phi^2(x/z, k^2)$$

Balitsky '95, Kovchegov 98

Unintegrated
gluon density

UGD



$$\frac{\partial \mathcal{F}_{BK}(x, k^2)}{\partial \ln 1/x} = \frac{N_c \alpha_s}{\pi} \int_0^\infty \frac{dl^2}{l^2} \left[\frac{l^2 \mathcal{F}_{BK}(x, l^2) - k^2 \mathcal{F}_{BK}(x, k^2)}{|k^2 - l^2|} + \frac{k^2 \mathcal{F}_{BK}(x, k^2)}{\sqrt{(4l^4 + k^4)}} \right] - \frac{\alpha_s^2}{R^2} \left\{ \left[\int_{k^2}^\infty \frac{dl^2}{l^2} \mathcal{F}_{BK}(x, l^2) \right]^2 + \mathcal{F}_{BK}(x, k^2) \int_{k^2}^\infty \frac{dl^2}{l^2} \ln \left(\frac{l^2}{k^2} \right) \mathcal{F}_{BK}(x, l^2) \right\}$$

Leading order equation
not applicable at large k_t

Not obvious probabilistic interpretation

Might be difficult for MC

Towards final states - resummed form of the BK

The strategy:

- Use the equation for dipole density. Simple nonlinear term
- Split linear kernel into resolved and unresolved parts
- Resumm the virtual contribution and unresolved ones in the linear part
- Use analogy to postulate nonlinear CCFM

The starting point:

$$\Phi(x, k^2) = \Phi_0(x, k^2) + \bar{\alpha}_s \int_{x/x_0}^1 \frac{dz}{z} \int_0^\infty \frac{dl^2}{l^2} \left[\frac{l^2 \Phi(x/z, l^2) - k^2 \Phi(x/z, k^2)}{|k^2 - l^2|} + \frac{k^2 \Phi(x/z, k^2)}{\sqrt{(4l^4 + k^4)}} \right] - \frac{\bar{\alpha}_s}{\pi R^2} \int_{x/x_0}^1 \frac{dz}{z} \Phi^2(x/z, k^2)$$

R is chosen such

That $\pi R^2 = 1$

Towards final states - resummed form of the BK

K. Kutak, K. Golec-Biernat, S. Jadach, M. Skrzypek

JHEP 1202 (2012) 117

$$\begin{aligned}\Phi(x, k^2) &= \Phi^0(x, k^2) \\ &+ \bar{\alpha}_s \int_x^1 \frac{dz}{z} \int \frac{d^2\mathbf{q}}{\pi q^2} [\Phi(x/z, |\mathbf{k} + \mathbf{q}|^2) - \theta(k^2 - q^2)\Phi(x/z, k)] \\ &- \bar{\alpha}_s \int_x^1 \frac{dz}{z} \Phi^2(x/z, k)\end{aligned}$$

Write in exclusive form

$$\begin{aligned}\Phi(x, k^2) &= \Phi^0(x, k^2) \\ &+ \bar{\alpha}_s \int_x^1 \frac{dz}{z} \int \frac{d^2\mathbf{q}}{\pi q^2} \Phi(x/z, |\mathbf{k} + \mathbf{q}|^2) \theta(q^2 - \mu^2) \\ &+ \bar{\alpha}_s \int_x^1 \frac{dz}{z} \int \frac{d^2\mathbf{q}}{\pi q^2} [\Phi(x/z, |\mathbf{k} + \mathbf{q}|^2) \theta(\mu^2 - q^2) - \theta(k^2 - q^2)\Phi(x/z, k)] \\ &- \bar{\alpha}_s \int_x^1 \frac{dz}{z} \Phi^2(x/z, k).\end{aligned}$$

Resolution scale introduced

Perform Mellin transform w.r.t x to get rid of "z" integral

$$\bar{\Phi}(\omega, k^2) = \int_0^1 dx x^{\omega-1} \Phi(x, k^2)$$

$$\Phi(x, k^2) = \int_{c-i\infty}^{c+i\infty} d\omega x^{-\omega} \bar{\Phi}(\omega, k^2)$$

BK equation in the resummed exclusive form

K. Kutak, K. Golec-Biernat, S. Jadach, M. Skrzypek


JHEP 1202 (2012) 117

$$\Phi(x, k^2) = \tilde{\Phi}^0(x, k^2) + \bar{\alpha}_s \int_x^1 dz \int \frac{d^2\mathbf{q}}{\pi q^2} \theta(q^2 - \mu^2) \frac{\Delta_R(z, k, \mu)}{z} \left[\Phi\left(\frac{x}{z}, |\mathbf{k} + \mathbf{q}|^2\right) - q^2 \delta(q^2 - k^2) \Phi^2\left(\frac{x}{z}, q^2\right) \right]$$

$$\Delta_R(z, k, \mu) \equiv \exp\left(-\bar{\alpha}_s \ln \frac{1}{z} \ln \frac{k^2}{\mu^2}\right)$$

- **The same** resummed piece for **linear and nonlinear**
- Initial distribution also gets multiplied by the Regge form factor
- **New scale introduced** to equation. One has to check dependence of the solution on it
- **Suggestive form to promote the CCFM equation to nonlinear equation** which is more suitable for description of final states

CCFM evolution equation - evolution with observer



- p - incoming proton, $p = (1, 0, 0, 1)P$
- q_i - emitted gluons, $q_i = y_i p + \bar{y}_i \bar{p} + q_{i\perp}$
- axial gauge with the gauge vector $\bar{p} = (1, 0, 0, -1)P$
- gluon polarization vector purely transverse $\varepsilon_\mu^{(\lambda)}(q) = g_\mu^{(\lambda)} - \frac{q_\mu \bar{p}^{(\lambda)}}{q\bar{p}}$

Extension of CCFM to non linear equation

- The second argument should be motivated by analogy to BK
- The third argument should reflect locally the angular ordering

$$\Phi(x, k^2) = \tilde{\Phi}^0(x, k^2) + \bar{\alpha}_s \int_x^1 dz \int \frac{d^2 \mathbf{q}}{\pi q^2} \theta(q^2 - \mu^2) \frac{\Delta_R(z, k, \mu)}{z} \left[\Phi\left(\frac{x}{z}, |\mathbf{k} + \mathbf{q}|^2\right) - q^2 \delta(q^2 - k^2) \Phi^2\left(\frac{x}{z}, q^2\right) \right]$$

$$\mathcal{E}(x, k^2, p) = \mathcal{E}_0(x, k^2, p) + \bar{\alpha}_s \int_x^1 dz \int \frac{d^2 \bar{\mathbf{q}}}{\pi \bar{q}^2} \theta(p - z\bar{q}) \Delta_s(p, z\bar{q}) \left(\frac{\Delta_{ns}(z, k, q)}{z} - \frac{1}{1-z} \right) \left[\mathcal{E}\left(\frac{x}{z}, k'^2, \bar{q}\right) - \bar{q}^2 \delta(q^2 - k^2) \mathcal{E}^2\left(\frac{x}{z}, \bar{q}^2, \bar{q}\right) \right]$$

Extension of CCFM to nonlinear equation



The unintegrated gluon density is obtained from

$$\mathcal{A}_{non-linear}(x, k^2, p) = \frac{N_c}{\alpha_s \pi^2} k^2 \nabla_k^2 \mathcal{E}(x, k^2, p)$$

The **nonlinear** term can be understood as a way to introduce the **decoherence** in emission of gluons which build unintegrated gluon density.

CCFM with saturation – consequences

Jung, Kutak '09
Avsar, Iancu '09



Avsar, Stasto '10

introduce line which will introduce effectively saturation effects in evolution. trajectories which enter the saturation region are rejected.

saturation scale saturates itself because of limited phase space due to existence of hard scale

Consequences for entropy production



K.Kutak '11

$$S = \frac{6C_F A_\perp}{\pi\alpha_s} Q_s^2(x) + S_0$$

Kiritsis, Tsaliotis '11

Jets and saturation another atrategy at present

S.Sapeta. KK
arxiv:1205.5035



$$\mathcal{F}_p(x, k^2) = \mathcal{F}_p^{(0)}(x, k^2)$$

$$+ \frac{\alpha_s(k^2) N_c}{\pi} \int_x^1 \frac{dz}{z} \int_{k_0^2}^{\infty} \frac{dl^2}{l^2} \left\{ \frac{l^2 \mathcal{F}_p\left(\frac{x}{z}, l^2\right) \theta\left(\frac{k^2}{z} - l^2\right) - k^2 \mathcal{F}_p\left(\frac{x}{z}, k^2\right)}{|l^2 - k^2|} + \frac{k^2 \mathcal{F}_p\left(\frac{x}{z}, k^2\right)}{|4l^4 + k^4|^{\frac{1}{2}}}\right\}$$

$$+ \frac{\alpha_s(k^2)}{2\pi k^2} \int_x^1 dz \left(P_{gg}(z) - \frac{2N_c}{z} \right) \int_{k_0^2}^{k^2} dl^2 \mathcal{F}_p\left(\frac{x}{z}, l^2\right)$$

$$- \frac{2\alpha_s^2(k^2)}{R^2} \left[\left(\int_{k^2}^{\infty} \frac{dl^2}{l^2} \mathcal{F}_p(x, l^2) \right)^2 + \mathcal{F}_p(x, k^2) \int_{k^2}^{\infty} \frac{dl^2}{l^2} \ln\left(\frac{l^2}{k^2}\right) \mathcal{F}_p(x, l^2) \right]$$


Corrections
of higher orders
Included.
Kin. Constr
DGLAP spf

Kwiecinski, Kutak '03

Andersson, Gustafson, Samuelsson '96
Kwiecinski, Martin, Sutton '96, Kwiecinski, Martin, Stasto '97

Further hints for saturation in F₂ data

S.Sapeta. KK
arxiv:1205.5035



Fit of BK-DGLAP
and BFKL-DGLAP
to combined H1-ZEUS
data

Very good description
with BK-DGLAP in range
 $Q^2 > 4.5 \text{ GeV}^2$



$$\chi^2 = 1.73$$

Very good description
with BFKL-DGLAP in
range
 $Q^2 > 4.5 \text{ GeV}^2$


$$\chi^2 = 1.5$$

Signatures of saturation in p-p and p-Pb

S.Sapeta. KK
arxiv:1205.5035



Observable suggested to study BFKL effects
Sabio-Vera, Schwensen '06



Reflects $\sim k^2$
behavior of gluon at small k^2

Conclusions and outlook

- There comes opportunity to test parton densities both when the parton density is probed at low x and at high kt .
- Used so far equations did not allow for this
- New representation for BK equation allowed for ansatz for well motivated equation which incorporates both **saturation** effects and **coherence**
- In the future it will be interesting to check whether this equation predicts **saturation of the saturation** scale as in other frameworks
- Results based on BK/DGLAP approach support hints for saturation in F_2
- Results based on BK/DGLAP approach **predict saturation** in p-Pb and suggest its presence in p-p
- Comparison to more general framework provided by *F. Dominguez, F. Huan, C. Marquet, B. Xiao, Phys.Rev.D83:105005,2011*