





Central Exclusive Production
(Selected New Results) 
(KRYSTHAL Collaboration)



V.A. Khoze (IPPP, Durham & PNPI)



Based on work by V.A. Khoze, M.G. Ryskin, W.J. Stirling and L.A. Harland-Lang. (KHYSTHAL collaboration)
For more details see [arXiv:1005.0695](https://arxiv.org/abs/1005.0695), [arXiv:1011.0680](https://arxiv.org/abs/1011.0680) and [arXiv:1105.1626](https://arxiv.org/abs/1105.1626)
 [arXiv:1204.4803](https://arxiv.org/abs/1204.4803)

- **Introduction (why we are interested in CEP processes?)**
- **'Diffractive Higgs' revisited.**
- **Standard Candle CEP processes.**
- **CEP as a way to study old and new heavy resonances.**
- **CEP through the KRYSTHAL eyes (new results, selected topics).**
 - ▶ **Diphoton CEP.**
 - ▶ **Dimeson CEP.**
- **SuperCHIC MC.**
- **Summary and Outlook.**

Introduction (why we are interested in CEP ?)

Why are we interested in central exclusive χ_c (χ_b , $\gamma\gamma$, jj) production?

- Driven by same mechanism as Higgs (or other new object) CEP at the LHC.

DO jj - results. LHCb χ_c
CMS, RHIC data expected

- χ_c , jj and $\gamma\gamma$ CEP has been observed by CDF.

→ Can serve as 'Standard Candle' processes, which allow us to check the theoretical predictions for central exclusive new physics signals at the LHC.

- $\chi_{c,b}$ production is of special interest:

- Heavy quarkonium production can shed light on the physics of bound states (lattice, NRQCD. . .).
- Potential to produce different J^P states, which exhibit characteristic features (e.g. angular distributions of forward protons).
- Possibility to shed light on the various 'exotic' charmonium states observed recently (X,Y,Z) charmonium-like states.


Spin-Parity Analyzer

(KMR-00, KKMR-2003)

- Detailed tests of dynamics of soft diffraction (KMR-02)

'Diffractive Higgs' revisited

(Marek's talk)



- Prospects for high accuracy ($\sim 1\%$) mass measurements (irrespective of the decay mode).

Higgs width (some BSM scenarios) ! ?

- Quantum number **filter/analyser**.
(0^{++} dominance ; **C,P-even**)

- H \rightarrow bb opens up (Hbb- coupl.)

(gg)_{CE}D bb in LO ; NLO, NNLO, b- mass effects - controllable.

- For some areas of the BSM param. space **CEP may** become **a discovery channel !**
- H \rightarrow WW*/WW - an added value (less challenging experimentally + small bgds., better PU cond.)



$\tau\tau$

- New leverage** -proton momentum correlations (probes of QCD dynamics , CP- violation effects...)

★ LHC : 'after discovery stage', Higgs **ID**..... How do we know what we've found?



Higgs or technipion or ???



mass, spin, couplings to fermions and Gauge Bosons, invisible modes...
 \rightarrow for all these purposes the **CEP** will be particularly handy !

Higgs Boson: cross section predictions

(Halfway)



$$\langle S_{\text{tot}}^2 \rangle = 0.02$$

- Cross section \sim fbs, i.e. roughly 4 orders of mag. lower than inclusive case (price paid for exclusivity).
- CEP of a CP-odd Higgs suppressed by $\sigma(0^-)/\sigma(0^+) \sim 1/100 \rightarrow$ with just a few signal events, the Higgs quantum numbers can be determined (does not rely on coupling to weak bosons).



THINGS TO DO !



- Account for the b-dependence of the survival factors $S_{\text{enh}}^2, S_{\text{eik}}^2$
- NLO effects in the unintegrated parton densities
- A systematic account of self-energy insertions in the propagator of the screening gluon
- The dependence on the gluon PDF is amplified by the fact that the CEP cross section is essentially proportional to $(xg(x))^4$.

(GLM-new results)

(Asher, Uri)



KKMR-04; CLP-07

CDF $\gamma\gamma$ data *may* suggest more 'LO-type' PDFs (\rightarrow more optimistic Higgs cross sections) are appropriate.

Available MSSM CEP x-sections

(Marek's talk)

2012 results M.Tasevsky+ HKW



M_h contours stay constant with M_A

Available MSSM CEP x-section stay constant with M_A (because $R = \text{MSSM}/\text{SM} = 1$ in this region) reaching maximum of 1.8 fb

x-section of 1.5 fb reachable but in a tiny allowed phase-space region. Outside this region the x-section is very small

Max -factor 5-10 enhancement.

a) LHC MSSM exclusion regions (red area) [HiggsBounds: P. Bechtle et al., Comput. Phys. Commun. 181 (2010) 138]

b) One possible region of interest (green area): SM Higgs at $M = 125 \text{ GeV} \pm 1 \text{ GeV}$ (exper.).

If theory uncertainties added: $122 < M < 128 \text{ GeV}$

[S. Heinemeyer et al., arXiv:1112.3026[hep-ph]]

CEP as a way to study old and new heavy resonances.

- Heavy Quarkonia

- Zoo of charmonium -like XYZ states



X(3872)

first and most puzzling state
(observed in 2003 at Belle)



- Discovered by BELLE in 2003, confirmed by BaBar, CDF, D0, CMS, LHCb.
- Possible spin-parity assignment: 1^{++} or 2^{-+}
- May well be of exotic nature : loosely bound molecule, diquark-antidiquark, hybrid,..... but a conventional 2 P-wave charmonium interpretation is still on the table (recent renewal of interest).
- BaBar (2010) seems to favour 2^{-+} though various theory groups find this assignment highly problematic.
- According to PDG $\Gamma(\pi^+ \pi^- J/\psi(1S))/\Gamma_{\text{total}} > 2.6\%$; $\Gamma(\gamma\psi(2S))/\Gamma_{\text{total}} > 3.0\%$, $\Gamma < 1.2$ MeV
(maybe two different states X(3872), X(3875))

CEP as a spin-parity analyzer could help to resolve the X(3872) puzzle.



Z(3930) $\equiv \chi_{c2}(2P)$

■ Above DD threshold .

■ Vertex detection at LHCb & RHIC→


■ **exclusive open charm: D^+D^- , $D^0\bar{D}^0$,**

■ Roughly the same expectations for CEP as for χ_{c2}

Triggering on J/ψ: $M \rightarrow J/\psi + \gamma$, $J/\psi + \rho$

CEP through the eyes of the KRYSTHAL (2008-2012)

- Colliding protons interact via a colour singlet exchange and remain intact: can be measured by adding detectors far down the beam-pipe. (or LRGs)
- A system X of mass M_X is produced at the collision point, and *only* its decay products are present in the central detector.
- The generic process $pp \rightarrow p + X + p$ is modeled perturbatively by the exchange of two t-channel gluons, with the use of pQCD justified by the presence of a hard scale $\sim M_X$.
- ‘ $J_z = 0$ selection rule’: production of states with non- $J_z^P = 0^+$ quantum numbers is strongly suppressed by ~ 2 orders of magnitude.



• $\chi_c, \gamma\gamma$ CEP already observed by CDF and jj CEP observed by CDF & D0. (Christina’s talk)



χ_{cJ} CEP is reported by LHCb (DIS-11)

(CMS-2012-first studies)



new CDF $\gamma\gamma$ CEP results (PRL-2012)

(Wenbo’s talk)



All measurements in agreement with Durham group (pre)dictions.



A MC event generator including⁸:

- Simulation of different CEP processes, including all spin correlations:
 - $\chi_{c(0,1,2)}$ CEP via the $\chi_c \rightarrow J/\psi\gamma \rightarrow \mu^+\mu^-\gamma$ decay chain.
 - $\chi_{b(0,1,2)}$ CEP via the equivalent $\chi_b \rightarrow \Upsilon\gamma \rightarrow \mu^+\mu^-\gamma$ decay chain.
 - $\chi_{(b,c)J}$ and $\eta_{(b,c)}$ CEP via general two body decay channels
 - Physical proton kinematics + survival effects for quarkonium CEP at RHIC.
 - Exclusive J/ψ and Υ photoproduction.
 - $\gamma\gamma$ CEP.
 - Meson pair ($\pi\pi$, KK , $\eta\eta\dots$) CEP.
 - More to come (dijets, open heavy quark, **Higgs**...?).
- Via close collaboration with CDF, STAR and LHC collaborations, in both proposals for new measurements and applications of SuperCHIC, it is becoming an important tool for current and future CEP studies.

⁸The SuperCHIC code and documentation are available at
<http://projects.hepforge.org/superchic/>


Standard Candle Processes

- CEP is a promising way to study new physics at the LHC, but we can also consider the CEP of lighter, established objects : χ_c , $\gamma\gamma$ and jj CEP already observed at the Tevatron.




- Can serve as ‘Standard Candle’ processes, which allow us to check the theoretical predictions for central exclusive new physics signals at the LHC, as well as being of interest in their own right¹.
- This talk will focus on the CEP of $\gamma\gamma$ and light meson pairs, $M\bar{M}$, at sufficiently high invariant mass for perturbative formalism to be applicable:
 - ▶ Provides novel application/test of hard exclusive formalism, complementary to more standard photon-induced processes ($\gamma\gamma \rightarrow M\bar{M}$, $\gamma\gamma^{(*)} \rightarrow M$ etc²).
 - ▶ Demonstrates application of MHV formalism to simplify/check calculations.
 - ▶ $\pi^0\pi^0$ CEP a possible background to $\gamma\gamma$ CEP.
 - ▶ Could probe the $q\bar{q}$ and gg content of η , η' mesons?
 - ▶ An interesting potential observable @ RHIC, Tevatron and LHC: meson pair CEP data (at lower p_{\perp}) already being taken by ALICE and CDF.

¹See LHL, V.A. Khoze, M.G. Ryskin, W.J. Stirling, [arXiv:1005.0695](https://arxiv.org/abs/1005.0695) and [arXiv:1011.0680](https://arxiv.org/abs/1011.0680) .

²For a recent review, see for example V. L. Chernyak, [arXiv:0912.0623](https://arxiv.org/abs/0912.0623). 

Standard Candle processes: χ_c CEP

- In [arXiv:0902.1271](#) CDF reported 65 ± 10 signal χ_c events observed via the $\chi_c \rightarrow J/\psi\gamma \rightarrow \mu^+\mu^-\gamma$ decay channel. This corresponds to $d\sigma(\chi_c)/dy_\chi|_{y=0} = (76 \pm 14)$ nb, in good agreement with Durham prediction of ~ 60 nb. (Krakow group)
- Recent LHCb data⁵: select ‘exclusive’ $\chi_c \rightarrow J/\psi\gamma$ events by vetoing on additional activity in given η range.
- LHCb see:



	$\sigma(pp \rightarrow pp(\chi_c \rightarrow \mu^+\mu^-\gamma))$ LHCb (pb)	SuperCHIC (pb)
χ_{c0}	9.3 ± 4.5	14
χ_{c1}	16.4 ± 7.1	10
χ_{c2}	28 ± 12.3	3



→ Encouraging agreement between data and (Durham) theory, accounting for (large) theory uncertainties for χ_c case, and potential inclusive contamination at LHCb.

Proton dissociation contribution

⁵LHCb-CONF-2011-022.

Dimeson CEP, motivation: $\gamma\gamma$ production

- 3 candidate events observed by CDF ([arXiv:0707.237](https://arxiv.org/abs/0707.237)), **Now 43 events**
- Similar uncertainties to χ_c case for low $E_{\perp\gamma} < E_{\text{cut}}$ scale, but this decreases for higher scales.
- More CDF events allow us to probe scaling of σ with cut on photon E_{\perp} ($\lesssim M_{\gamma\gamma}/2$): strong predicted fall-off with $M_{\gamma\gamma}$ driven by Sudakov factor (already seen in dijet data).



(KMRS-04)
(HKRS-10)




- **However:** $\pi^0\pi^0(\eta\eta)$ production, with one photon from each decay either undetected or two photons merging, is a potentially important background (pure QCD process).



(now proved to be very small (CDF) in agreement with our expectations)

Standard Candle processes: $\gamma\gamma$ CEP

- $\gamma\gamma$ CEP: represents clean signal, with less of the theory issues related to, e.g. χ_c CEP. \rightarrow ideal 'standard candle'.
- **New** CDF $\gamma\gamma$ data⁶ for $E_{\perp}(\gamma) > 2.5$ GeV, $|\eta(\gamma)| < 1$. They find $\sigma_{\gamma\gamma} = 2.48^{+0.40}_{-0.35}$ (stat) $^{+0.40}_{-0.51}$ (syst) pb,
- Theory predictions: 1.42 pb (MSTW08LO) and 0.35 pb (MRST99), with approx. uncertainties (additional to PDFs) $\sim \times 2$.
- $\pi^0\pi^0$ BG observed to be small, in agreement with non-trivial Durham prediction (follows from $J_z = 0$ selection rule).
- **New** CMS $\gamma\gamma$ search for $E_{\perp}(\gamma) > 5$ GeV⁷: $\sigma_{\gamma\gamma} < 1.3$ pb @ 95% confidence. (Wenbo's talk)




⁶CDF Collaboration, T. Aaltonen et al., Phys. Rev. Lett. 108, 081801 (2012) [arXiv:1112.0858](https://arxiv.org/abs/1112.0858).

⁷CMS PAS FWD-11-005.

New Exclusive $\gamma\gamma$: Conclusions

Exclusive Photon-Pair Production	
Theoretical	$\sigma_{\text{SuperCHIC}}^{ \eta < 1, E_T > 2.5 \text{ GeV}} = 0.35 \times 3 \text{ pb (MRST99)}$
	$\sigma_{\text{SuperCHIC}}^{ \eta < 1, E_T > 2.5 \text{ GeV}} = 1.42 \times 3 \text{ pb (MSTW08LO)}$
Measured	$\sigma_{\gamma\gamma \text{ excl.}}^{ \eta < 1, E_T > 2.5 \text{ GeV}} = 2.48^{+0.40}_{-0.35} (\text{stat})^{+0.40}_{-0.51} (\text{syst}) \text{ pb}$



- **First observation** of exclusive $\gamma\gamma$ in hadron-hadron collisions.
- Measurement of the cross section of the exclusive production of two high- E_T photons in hadron hadron collisions.
- This corresponds to 1 in 25 billion inelastic collisions.
- Constraint on central exclusive Higgs if existing (produced by same mechanism).
- Paper recently published:
Phys. Rev. Lett. 108, 081801 (2012).

NEW! NLO effects-factor of 1.55

Currently theor. uncertainties are under further revision.

Modeling meson pair CEP perturbatively


- Simpler exclusive process $\gamma\gamma \rightarrow M\bar{M}$ ($= \pi^0\pi^0, \pi^+\pi^-, K^+K^- \dots$) at large angles was calculated ~ 30 years ago³.
- Total amplitude given by convolution of parton level $\gamma(\lambda_1)\gamma(\lambda_2) \rightarrow q\bar{q}q\bar{q}$ amplitude with non-perturbative pion wavefunction $\phi(x)$

$$\mathcal{M}_{\lambda_1\lambda_2}(s, t) = \int_0^1 dx dy \phi(x)\phi(y) T_{\lambda_1\lambda_2}(x, y; s, t)$$


where helicity amplitudes $T_{\lambda_1\lambda_2}$ can be calculated perturbatively.

- With suitable choice of $\phi(x)$ shape, $\gamma\gamma \rightarrow M\bar{M}$ data are described quite well (see plot⁴).

40 diagrams (4 basic)



★ $T_{++} = T_{--} = 0$



³S. J. Brodsky and G. P. Lepage, Phys. Rev. D 24 (1981) 1808.

(M. Benayoun, V. Chernyak, -1990)

⁴Data taken from Belle Collaboration, Phys. Lett. B615 (2005) 39

- Simplest case: production of flavour non-singlet scalar mesons (e.g. $\pi^0\pi^0, \pi^+\pi^- \dots$).
- Can calculate the LO $gg \rightarrow M\bar{M} (= q\bar{q}q\bar{q})$ amplitudes to give

$$T_{++} = T_{--} = 0,$$

is this easy to understand?



$$T_{-+} = T_{+-} \propto \frac{\alpha_S^2}{a^2 - b^2 \cos^2 \theta} \left(\frac{N_c}{2} \cos^2 \theta - C_F a \right),$$


where $a, b = (1 - x)(1 - y) \pm xy$.

- $J_Z = 0$ amplitudes vanish, as in $\gamma\gamma \rightarrow M\bar{M}$ for neutral mesons. We therefore expect a strong suppression in flavour non-singlet $M\bar{M}$ CEP due to $J_Z = 0$ selection rule.
- $J_Z = 2$ amplitudes contain 'radiation zero', vanishing for a physical value of $\cos^2 \theta$. Well known effect in all gauge theories (e.g. $u\bar{d} \rightarrow W^+\gamma$), but usually washed out in QCD by colour averaging. Here, position of zero depends on the choice of $\phi(x)$, and we find that there is always a zero in the physical region for any choice of $\phi(x)$ and general N_c .

$gg \rightarrow M\bar{M}$ amplitude: Feynman diagrams

Vanishing of T_{++}, T_{--} follows after calculating:

is this easy to understand ?



currently popular (among the **more** formal community) **MHV-technique**



$gg \rightarrow M\bar{M}$ amplitude: MHV calculation (1)

- $g(+)\bar{g}(+) \rightarrow q(\pm)\bar{q}(\mp)q(\pm)\bar{q}(\mp)$ amplitude is MHV: maximum $(n - 2)$ number of particles have same helicity.
- Such amplitudes known to have remarkably simple forms, and corresponding 'spinor helicity' formalism can greatly simplify calculation.
- T_{++}, T_{--} can be calculated from known Parke-Taylor amplitude⁵

$$M_n \propto \sum_{\sigma} \frac{\langle k_p k_{\bar{q}} \rangle}{\langle k_p a_1 \rangle \cdots \langle a_l k_{\bar{q}} \rangle} \frac{\langle k_q k_{\bar{p}} \rangle}{\langle k_q b_1 \rangle \cdots \langle b_{l'} k_{\bar{p}} \rangle} (\lambda^{a_1} \cdots \lambda^{a_l})_{i_1 j_2} (\lambda^{b_1} \cdots \lambda^{b_{l'}})_{i_2 j_1}$$

$$- \frac{1}{N_c} \frac{\langle k_p k_{\bar{p}} \rangle}{\langle k_p a_1 \rangle \cdots \langle a_l k_{\bar{p}} \rangle} \frac{\langle k_q k_{\bar{q}} \rangle}{\langle k_q b_1 \rangle \cdots \langle b_{l'} k_{\bar{q}} \rangle} (\lambda^{a_1} \cdots \lambda^{a_l})_{i_1 j_1} (\lambda^{b_1} \cdots \lambda^{b_{l'}})_{i_2 j_2} .$$

- Making colour singlet identification ($i_1 = j_2, i_2 = j_1$) and identifying $q\bar{q}, p\bar{p}$ with collinear quarks within mesons


$$k_q = xk_3 \quad k_{\bar{q}} = (1 - y)k_4 \quad k_p = yk_4 \quad k_{\bar{p}} = (1 - x)k_3 ,$$

then amplitude reduces to

$$M \propto \langle k_3 k_2 \rangle \langle k_1 k_4 \rangle + \langle k_1 k_3 \rangle \langle k_2 k_4 \rangle - \langle k_3 k_4 \rangle \langle k_1 k_2 \rangle = 0 ,$$

which vanishes from the Schouten identity.


⁵M. L. Mangano, S. J. Parke, Phys. Rept. 200 (1991) 301-367



Here the indices $r(\bar{r})$ and $s(\bar{s})$ refer to the quarks (antiquarks) with colour indices $i_1(j_1)$ and $i_2(j_2)$, respectively, and the labels a_i, b_i refer to the gluons, while the standard spinor contraction ' $\langle k, l \rangle$ '

$gg \rightarrow M\bar{M}$ amplitude: MHV calculation (2)


- The vanishing of the $gg \rightarrow M\bar{M}$ $J_Z = 0$ amplitudes follow directly from the corresponding 6-particle MHV amplitude. This result depends crucially on the colour singlet projection and collinearity of the $q\bar{q}$ pairs, and only occurs for non-flavour singlet mesons
- The MHV formalism can be extended to include the non-MHV $|J_Z| = 2$ amplitude: contributing amplitudes given by tree graphs in which the vertices are the usual tree-level MHV scattering amplitudes continued off-shell⁶.
F. Cachazo, P. Svrcek, E. Witten, JHEP 0409 (2004) 006 [hep-th/0403047].
- More complicated than $J_Z = 0$ case, but an explicit calculation within this framework confirms our result.



⁶see e.g. G. Georgiou, V. V. Khoze, JHEP 0405 (2004) 070.

Flavour singlet meson production

- A second set of diagrams can in general contribute, where the $q\bar{q}$ forming the mesons connected by a quark line (no equivalent diagram in $\gamma\gamma \rightarrow M\bar{M}$ process).
- Only relevant for flavour singlet states (e.g. for $gg \rightarrow \pi^0\pi^0$, $|u\bar{u}\rangle$ and $|d\bar{d}\rangle$ Fock components interfere destructively).
- In this case the $J_z = 0$ amplitude does not vanish \rightarrow Expect strong enhancement in $\eta'\eta'$ CEP rate⁷ and (through η - η' mixing), some enhancement to $\eta\eta$ rate. $\eta\eta'$ CEP can also occur via this mechanism.
- Also: any sizable gg component to flavour singlet states, contributing through $gg \rightarrow 4g$ and $gg \rightarrow q\bar{q}gg$ processes, may in principle strongly enhance the CEP cross section (again $J_z = 0$ amplitudes do not vanish). A significant 'excess' in future CEP data could be evidence for this.




⁷Recall quark content of $|\eta'\rangle$ is dominantly $\sim |u\bar{u} + d\bar{d} + s\bar{s}\rangle$



Numerical Results

(our new results will be available soon)

- Strong enhancement in flavour singlet states clear, with precise η'/η hierarchy given by choice of $\eta - \eta'$ mixing angle.
- CEP cross sections for vector mesons ($\rho\rho, \omega\omega, \phi\phi$) can be calculated in the same way.



- $\pi^0\pi^0$ CEP can in principle be an important background to $\gamma\gamma$ CEP, but we find this not to be the case. This depends crucially on vanishing of the $gg \rightarrow \pi^0\pi^0$ amplitude for $J_Z = 0$ initial-state gluons.
- However: possible $J_Z = 0$ contribution from higher twist effects, NNLO corrections... could increase flavour non-singlet rate by a factor 'a few'. Also, possible non-perturbative contribution at lower p_{\perp} ? (K-factor,..)

New CDF data nicely confirm this !



Prospects of further measurements

Data/Durham theory comparison

- χ_c , $\gamma\gamma$ and jj CEP already observed at the Tevatron, χ_c at the LHC, with more to come...
- So far, data has been in overall good agreement with the Durham theory, giving confidence in the Higgs predictions, up to a $\sim \frac{\times 2}{\div 2}$ uncertainty.
- CDF $\gamma\gamma$ data *may* suggest more 'LO-type' PDFs (\rightarrow more optimistic Higgs cross sections) are appropriate.
- Studies are ongoing, and other observables (e.g. CEP of meson pairs, $\pi^+\pi^-$, $\eta^{(\prime)}\eta^{(\prime)}$...) are being explored theoretically and experimentally.

Towards a Full Acceptance Detector (bj- 1992)



CMS (& ATLAS) currently blind between $\eta = 6.4$ (CASTOR) and beam rapidity ($y_p = 8.9$ @ 7 TeV) except ZDC (neutrals).

(Totem:T1,T2)

IS THERE A WAY OUT ?

Yes, an addition of **Forward Shower Counters** around beam pipes at CMS!

(8 FSC per side see showers from particles with $|\eta| = 7-9$)

(Alice is installing such counters, discussions at the LHCb)

(FSC → at least a good foot in the door)



CMS+TOTEM- common data
taking foreseen in 2012
90m expected in July
Coverage in η : -6.5-6.5



The Compact Muon Solenoid Experiment

CMS Note

Mailing address: CMS CERN, CH-1211 GENEVA 23, Switzerland



July 19, 2010

Physics and Beam Monitoring with Forward Shower Counters (FSC) in CMS

Alan J. Bell, David d'Enterria, Richard Hall-Wilton ^{a)}, Gabor Veres

CERN, Geneva, Switzerland

Valery Khoze

Institute for Particle Physics Phenomenology, Durham University, U.K.

Michael Albrow ^{a)}, Nikolai Mokhov, Igor Rakhno

Fermi National Accelerator Laboratory, USA

Erik Brücken, Jerry Lamsa ^{b)}, Rauno Lauhakangas, Risto Orava

Dept. of Physical Sciences, University of Helsinki and Helsinki Institute of Physics, Finland

Paul Debbins, Edwin Norbeck, Yasar Onel, Ionos Schmidt

University of Iowa, USA

Oleg Grachov, Michael Murray

Kansas University, USA

Jeff Gronberg

Lawrence Livermore National Laboratory, USA

Jonathan Hollar

U.C. Louvain, Belgium

Greg Snow

2009 JINST 4 P10001

RECEIVED: August 4,
ACCEPTED: September 20,
PUBLISHED: October 2,

Forward physics with rapidity gaps at the LHC

M.G. Albrow,^{a,1} A. De Roeck,^b V.A. Khoze,^c J. Lämsä,^{d,d'} E. Norbeck,^f Y. Onel,^f
R. Orava,^e A. Penzo^g and M.G. Ryskin^h

**Approved by CMS MB
for Jan-Feb 2011 installation.**

“Limited approval” :
Go ahead without detracting from
necessary shutdown work.

Most value is 2011 running
& when $\langle n/x \rangle < \sim 5$

**Station 3 (114m) Installed on both
sides.**

March Technical Stop (28-31.03.11).

The FSCs- these are for real !

Main concern- lumi per bunch crossing might be too high. 🤔

What about a precise measurement of SD? This certainly needs all the counters and some low lumi run, or at least bunches. 🤔

Both CMS & ATLAS requested special low-PU runs in 2012

**GOOD
LUCK**



Summary and Outlook

- CEP processes observed at the Tevatron, RHIC and early LHC can serve as 'standard candles' for new physics CEP at the LHC.
- New LHCb $\chi_c \rightarrow J/\psi$ data, support: previous suggestion that $\chi_{c(1,2)}$ contribute to CDF χ_c data.
- First estimates of dissociative background given.
- χ_{c0} CEP via two-body decays ($\pi^+\pi^+$, K^+K^- ...) interesting and realistic channels, with continuum background expected to be low. Other decay channels (e.g. $p\bar{p}$, $\Lambda\bar{\Lambda}$, $2(\pi^+\pi^-)$...) also possible.
- The CEP of mesons pairs at high invariant masses ($/k_\perp$) is an interesting process, representing a novel application of pQCD framework for describing exclusive processes.
- Measurement of $\pi\pi$ (KK ...) CEP at lower mass/ k_\perp values would help constrain non-perturbative models.
- CEP could help probe the gluonic structure of η , η' mesons.
- Perturbative calculation predicts that $\pi^0\pi^0$ BG to $\gamma\gamma$ CEP is suppressed.
- New CDF $\gamma\gamma$ data gives encouraging results! Could shed light on the gluon density...
- More CEP results to come from RHIC, the Tevatron and LHC in the future.



BACKUP




Central Exclusive $\gamma\gamma$ and e^+e^- Production

95% confidence level upper limit

$$\sigma_{\text{exclusive } \gamma\gamma \text{ production}}^{E_T(\gamma) > 5.5 \text{ GeV}, |\eta(\gamma)| < 2.5} < 1.30 \text{ pb}$$

This upper limit on the cross section is actually on the sum of

- ▶ exclusive production
- ▶ semi-exclusive production with no particles from the proton dissociation within $|\eta| < 5.2$ (much less controlled theoretically, \sim similar magnitude)



Theoretical predictions


- ▶ exclusive only
- ▶ predictions would be higher by a factor of ~ 2 if contributions from semi-exclusive included

Diffraction with Forward Shower Counters FSC

Mike Albrow,

BSC very important as rap gap detectors.
All LHC experiments should have them!

What: We propose to install a set of scintillating
outgoing beam pipes at CMS, ~



Why:


- (a) As veto in Level 1 diff. triggers to reduce useless pile-up events.
- (b) To detect rapidity gaps in diffractive events (p or no-p).
- (c) Measure “low” mass diffraction and double pomeron exchange.
- (d) Measure σ_{INEL} (if luminosity known, e.g. by Van der Meer)
- (e) Help establish exclusivity in central exclusive channels
- (f) To monitor beam conditions on incoming and outgoing beams.
- (g) To test forward flux simulations (MARS etc.)
- (h) Additional Luminosity monitor.

Also: They may provide valuable tests of radiation environment
to be expected for HPS = High Precision Spectrometers

FSC & others


FSC alone

ZDC alone



>4 hits in FSC or > 1 track in HF
or CASTOR or ZDC(min)

M. Albrow et al, JINST 4:P10001,2009.



10 GeV


Generated diffractive mass (PYTHIA/PHOJET) as $\log(M_X)$, M_X in GeV/c^2 , cf to calculated from rapidity gap edge:
(a) full η coverage
(b) $\eta < 4.7$ (no FSC)

Below 10 GeV/c^2 FSC contain most particles

$\gamma\gamma$ CEP: new results (2)

- Expect theory estimates to be somewhat conservative:
 - ▶ S_{enh}^2 effect somewhat overestimated– latest number $\approx 20\%$ bigger.
 - ▶ Small fraction of $\gamma\gamma$ events that are not truly exclusive ($\approx 10\%$).
 - ▶ NLO corrections could be numerically quite large (c.f. $\chi_{c0} \rightarrow gg$ and $H \rightarrow gg$, both receive infrared π^2 numerical enhancement). Including finite part of 1-loop corrections¹⁴ to $gg \rightarrow \gamma\gamma$ get $K_{\text{nlo}} \approx 1.6$, so a similar enhancement may be present. **However**: need full NLO calculation, divergences included in f_g 's now cancel virtual IR divergences, and will get new finite contributions specific to CEP.

- Must also bear in mind reasonable theory uncertainties, but nevertheless some tension between theory (MRST99) and new data exists...



$\gamma\gamma$ CEP: PDF comparison (2)

- The gluon density is not sufficiently well described by fixed order, twist = 2 DGLAP at low x and Q^2 .
- There is some indication from, e.g. diffractive J/ψ production that the $g(x, Q^2)$ is larger than the current NLO PDFs¹⁵.
- Can also use, e.g. $\gamma\gamma$ CEP to shed light on the gluon density, with the LO and NLO gluons giving approx. upper and lower bounds on the CEP cross section due to the (large) PDF uncertainty.
- Use an updated model¹⁶ for S_{eik}^2 , which includes the new TOTEM elastic data (requires $\Omega(b_t) \uparrow$ in particular at lower b_t , and therefore $S_{\text{eik}}^2 \downarrow$), and for S_{enh}^2 (somewhat higher than previously), gives factor ~ 2 decrease in σ @ 7 TeV. The $\gamma\gamma$ CEP cross sections (in pb) are predicted to be (for $E_{\perp} > 2.5$ GeV):

	MSTW08LO	CTEQ6L	MRST99	CT10	NNPDF2.1
$\sqrt{s} = 1.96$ TeV ($ \eta < 1$)	1.4	2.2	0.35	0.47	0.29
$\sqrt{s} = 7$ TeV ($ \eta < 1$)	2.1	2.0	0.32	0.29	0.16
$\sqrt{s} = 7$ TeV ($ \eta < 2.4$)	6.2	6.2	0.94	0.91	0.50


¹⁵A. Martin, C. Nockles, M. G. Ryskin, and T. Teubner, Phys.Lett. B662, 252 (2008), 0709.4406.

¹⁶M. Ryskin, A. Martin, and V. Khoze, (2012), 1201.6298.

What we expect within the framework of the Perturbative Durham formalism

(KMR-01, KKMR-03, KMRS-04, HKRS-10)

Example, O++ -case



$$T = A\pi^2 \int \frac{d^2 Q_{\perp} P(\chi(0^+))}{Q_{\perp}^2 (\vec{Q}_{\perp} - \vec{p}_{1\perp})^2 (\vec{Q}_{\perp} + \vec{p}_{2\perp})^2} f_g(x_1, x'_1, Q_1^2, \mu^2; t_1) f_g(x_2, x'_2, Q_2^2, \mu^2; t_2),$$

$$A^2 = 8\pi\Gamma(\chi \rightarrow gg)/M_{\chi}^3 \quad *K_{\text{NLO}} \quad P(\chi(0^+)) = (\vec{Q}_{\perp} - \vec{p}_{1\perp}) \cdot (\vec{Q}_{\perp} + \vec{p}_{2\perp}).$$

- The $gg \rightarrow \chi_{cJ}$ vertex can be calculated by a simple extension of the previous $\gamma\gamma \rightarrow \chi_c$ potential model results. These give the Lorentz structure of the vertices, while the normalisation is set by the derivative of the P-wave meson wavefunction at the origin $\phi'_P(0)$.

- Strong sensitivity to the polarization structure of the vertex in the bare amplitude.

KMR-01

Absorption is sizeably distorted by the polarization structure (affects the b-space distr.)

- χ_c, χ_b -production is especially sensitive to the effects of enhanced absorption
 - larger available rapidity interval

KMR-02, KKMR-03,
HKRS 09-10

- lower scale \rightarrow larger dipole size \rightarrow larger absorption
(Gap size for χ_c at the Tevatron is expected to exceed that for the Higgs at the LHC)

KMR-02

- Forward proton distributions & correlations- possibility to test diffraction dynamics


χ_c CEP @ LHCb (1)

- Select 'exclusive' events by vetoing on additional activity in given η range— $\chi_c \rightarrow J/\psi\gamma$ events seen by LHCb.
- Expect $\sigma_{\chi_0} \approx \sigma_{(\chi_1+\chi_2)}$ → recalling $\text{Br}(\chi_{c0} \rightarrow J/\psi\gamma)$ suppression, observation of χ_{c0} events strongly favours exclusivity.
- LHCb see¹:

Proton dissociation contribution

	$\sigma(pp \rightarrow pp(J/\psi + \gamma))$ LHCb (pb)	SuperChic prediction (pb)
χ_{c0}	9.3 ± 4.5	14
χ_{c1}	16.4 ± 7.1	10
χ_{c2}	28 ± 12.3	3


→ Good agreement for $\chi_{c(0,1)}$ states (recall large theory uncertainty), but a significant excess of χ_{c2} events above theory prediction.



¹Preliminary data— LHCb-CONF-2011-022

J/ψ + Photon Mass

- $\chi_{c0}:\chi_{c1}:\chi_{c2}$ ratio determined from fit to mass spectrum
- $\Psi(2S)$ background



25 χ_{c0} Candidates


56 χ_{c1} Candidates

99 χ_{c2} Candidates

Future Plans



- In 2011 LHCb collected $\sim 1 \text{ fb}^{-1}$
- Average $\mu \sim 1.5$
- x 30 Increase in Stats!!



- Odderon?
Low x gluon behavior?
- χ_b ? More perturbative
better test of CEP
- Luminosity measurement
with $\Upsilon\Upsilon$ Dimuons

- Low Mult DiHadron Trigger



- Exclusive $\pi^+\pi^-, K^+K^-$
- $\chi_c \rightarrow \pi^+\pi^-, K^+K^-$ (0,1,2 Separation)
- MisId determination for $\Upsilon\Upsilon$ Dimuons

- Low Mult DiPhoton Trigger



- Exclusive $\Upsilon\Upsilon$

Dermot Moran (University of Manchester)
On behalf of the LHCb Collaboration