



# Supernova Neutrinos

Georg G. Raffelt

Max-Planck-Institut für Physik, München, Germany

# Sanduleak -69 202



Tarantula Nebula

Large Magellanic Cloud  
Distance 50 kpc  
(160.000 light years)



**Sanduleak –69 202**




**Supernova 1987A**

23 February 1987



凡十一日没三年三月乙巳出東南方大中祥符四年正月丁丑見南斗魁前天禧五年四月丙辰出軒轅前星西北大如桃速行經軒轅太星入太微垣掩右執法犯次將歷屏星西北凡七十五日入濁没明道元年六月乙巳出東北方近濁有芒彗至丁巳凡十三日没至和元年五月己丑出天關東南可數寸歲餘稍没熙寧二年六月丙辰出箕度中至七月丁卯犯箕乃散三年十一月丁未出天因元祐六年十一月辛亥出參度中犯掩側星壬子犯九游星十二月癸酉入奎至七年三月辛亥乃散紹興八年五月守婁

# The Crab Pulsar




Chandra x-ray images



# Supernova Remnant in Cas A (SN 1667?)

Chandra  
x-ray image



Non-pulsar  
compact remnant



Walter Baade (1893–1960)




Fritz Zwicky (1898–1974)

Baade and Zwicky were the first to speculate about a connection between supernova explosions and neutron-star formation


[Phys. Rev. 45 (1934) 138]

# Stellar Collapse and Supernova Explosion

Main-sequence star



Helium-burning star






# Stellar Collapse and Supernova Explosion


Onion structure

Collapse (implosion)



# Stellar Collapse and Supernova Explosion

**Newborn Neutron Star**




**Proto-Neutron Star**


$$\rho \sim \rho_{\text{nuc}} = 3 \times 10^{14} \text{ g cm}^{-3}$$

$$T \sim 30 \text{ MeV}$$

**Explosion**



# Stellar Collapse and Supernova Explosion



Gravitational binding energy

$$E_b \approx 3 \times 10^{53} \text{ erg} \approx 17\% M_{\text{SUN}} c^2$$

This shows up as

99% Neutrinos

1% Kinetic energy of explosion

0.01% Photons, outshine host galaxy


Neutrino luminosity

$$L_\nu \sim 3 \times 10^{53} \text{ erg} / 3 \text{ sec}$$

$$\sim 3 \times 10^{19} L_{\text{SUN}}$$

While it lasts, outshines the entire visible universe

# Neutrino Signal of Supernova 1987A




Kamiokande-II (Japan)  
Water Cherenkov detector  
2140 tons  
Clock uncertainty  $\pm 1$  min

Irvine-Michigan-Brookhaven (US)  
Water Cherenkov detector  
6800 tons  
Clock uncertainty  $\pm 50$  ms

Baksan Scintillator Telescope  
(Soviet Union), 200 tons  
Random event cluster  $\sim 0.7/\text{day}$   
Clock uncertainty  $+2/-54$  s

**Within clock uncertainties,  
all signals are contemporaneous**

# Interpreting SN 1987A Neutrinos



Assume

- Thermal spectra
- Equipartition of energy between  $\nu_e, \bar{\nu}_e, \nu_\mu, \bar{\nu}_\mu, \nu_\tau$  and  $\bar{\nu}_\tau$

Jegerlehner,  
Neubig & Raffelt,  
PRD 54 (1996) 1194

# Predicting Neutrinos from Core Collapse

## The Possible Role of Neutrinos in Stellar Evolution

It can be considered at present as definitely established that the energy production in stars is caused by various types of thermonuclear reactions taking place in their interior. Since these reaction chains usually contain the processes of  $\beta$ -disintegration accompanied by the emission of high speed neutrinos, and since the neutrinos can pass almost without difficulty through the body of the star, we must assume that a certain part of the total energy produced escapes into interstellar space without being noticed as the actual thermal radiation of the star. Thus, for example, in the case of the carbon-nitrogen cycle in the sun, about 7 percent of the energy produced is lost in the form of neutrino radiation. However, since, in such reaction chains, the energy taken away by neutrinos represents a definite fraction of the total energy liberation, these losses are of but secondary importance for the problem of stellar equilibrium and evolution.

More detailed calculations on this collapse process are now in progress.

The George Washington University,  
Washington, D. C.,

University of São Paulo,  
São Paulo, Brazil,  
November 23, 1940.

\* Fellow of the Guggenheim Memorial Foundation. Now in Washington, D. C.


Phys. Rev. 58:1117 (1940)





# Explosion Mechanism

# Why No Prompt Explosion?




- $0.1 M_{\text{sun}}$  of iron has a nuclear binding energy  $\approx 1.7 \times 10^{51}$  erg
- Comparable to explosion energy

- **Shock wave forms within the iron core**
- **Dissipates its energy by dissociating the remaining layer of iron**




# Neutrinos Rejuvenating Stalled Shock





Picture adapted from Janka, astro-ph/0008432

# Delayed Explosion




Wilson, Proc. Univ. Illinois Meeting on Num. Astrophys. (1982)  
Bethe & Wilson, ApJ 295 (1985) 14

# Exploding Models (8–10 Solar Masses)




Kitaura, Janka & Hillebrandt: “Explosions of O-Ne-Mg cores, the Crab supernova, and subluminous type II-P supernovae”, astro-ph/0512065


# 3D Simulation (Garching group)



# Standing Accretion Shock Instability (SASI)



# Gravitational Waves from Core-Collapse Supernovae




Müller, Rampp, Buras, Janka, & Shoemaker, astro-ph/0309833

“Towards gravitational wave signals from realistic core collapse supernova models”


# Three Phases of Neutrino Emission

## Prompt $\nu_e$ burst




- Shock breakout
- De-leptonization of outer core layers

## Accretion



- Shock stalls  $\sim 150$  km
- Neutrinos powered by infalling matter

## Cooling



Cooling on neutrino diffusion time scale

- Spherically symmetric model ( $10.8 M_{\odot}$ ) with Boltzmann neutrino transport
    - Explosion manually triggered by enhanced CC interaction rate
- Fischer et al. (Basel group), A&A 517:A80, 2010 [arxiv:0908.1871]



**Neutrinos from Next Nearby SN**



# Operational Detectors for Supernova Neutrinos

MiniBooNE  
(200)

HALO  
(tens)

LVD (400)  
Borexino (100)


Baksan  
(100)

Super-K ( $10^4$ )  
KamLAND (400)


Daya Bay  
(100)

IceCube ( $10^6$ )


In brackets events  
for a “fiducial SN”  
at distance 10 kpc



# Super-Kamiokande Neutrino Detector




# Simulated Supernova Burst in Super-Kamiokande




Movie by C. Little, including work by S. Farrell & B. Reed,  
(Kate Scholberg's group at Duke University)

<http://snews.bnl.gov/snmovie.html>

# IceCube as a Supernova Neutrino Detector



- Each optical module (OM) picks up Cherenkov light from its neighborhood
- $\sim 300$  Cherenkov photons per OM from SN at 10 kpc
- Bkgd rate in one OM  $< 300$  Hz
- SN appears as “correlated noise” in  $\sim 5000$  OMs




SN signal at 10 kpc  
10.8  $M_{\text{sun}}$  simulation  
of Basel group  
[arXiv:0908.1871]


Pryor, Roos & Webster (ApJ 329:355, 1988), Halzen, Jacobsen & Zas (astro-ph/9512080)

# Variability seen in Neutrinos

## Luminosity




## Detection rate in IceCube



Probably smaller in realistic 3D models

Lund, Marek, Lunardini, Janka & Raffelt, arXiv:1006.1889  
Using 2-D model of Marek, Janka & Müller, arXiv:0808.4136


# Quark-Matter Phase Transition Signature in IceCube



Dasgupta, Fischer, Horiuchi, Liebendörfer, Mirizzi, Sagert & Schaffner-Bielich  
arXiv:0912.2568

# Next Generation Large-Scale Detector Concepts


**DUSEL  
LBNE**





**Hyper-K**

**Memphys**

**Megaton-scale  
water Cherenkov**



**5-100 kton  
liquid Argon**



## DETECTOR LAYOUT

**Cavern**  
height: 115 m, diameter: 50 m  
shielding from cosmic rays: ~4,000 m.w

**Muon Veto**  
plastic scintillator panels (on top)  
Water Cherenkov Detector  
1,500 phototubes  
100 kt of water  
reduction of fast  
neutron background


**Steel Cylinder**  
height: 100 m, diameter: 30 m  
70 kt of organic liquid  
13,500 phototubes

**Buffer**  
thickness: 2 m  
non-scintillating organic liquid  
shielding external radioactivity

**Nylon Vessel**  
parting buffer liquid  
from liquid scintillator

**Target Volume**  
height: 100 m, diameter: 26 m  
50 kt of liquid scintillator

vertical design is favourable in terms of rock pressure and buoyancy forces




**100 kton scale  
scintillator**

**LENA  
HanoHano**

# SuperNova Early Warning System (SNEWS)


Early light curve of SN 1987A



- Neutrinos arrive several hours before photons
- Can alert astronomers several hours in advance



<http://snews.bnl.gov>









# Supernova Rate

# Local Group of Galaxies


With megatonne class (30 x SK)  
60 events from Andromeda



Current best neutrino detectors  
sensitive out to few 100 kpc




# Core-Collapse SN Rate in the Milky Way



References: van den Bergh & McClure, *ApJ* 425 (1994) 205. Cappellaro & Turatto, *astro-ph/0012455*. Diehl et al., *Nature* 439 (2006) 45. Strom, *Astron. Astrophys.* 288 (1994) L1. Tammann et al., *ApJ* 92 (1994) 487. Alekseev et al., *JETP* 77 (1993) 339 and my update.


# High and Low Supernova Rates in Nearby Galaxies

M31 (Andromeda)      $D = 780 \text{ kpc}$




Last Observed Supernova: 1885A

NGC 6946      $D = (5.5 \pm 1) \text{ Mpc}$



Observed Supernovae:  
1917A, 1939C, 1948B, 1968D, 1969P,  
1980K, 2002hh, 2004et, 2008S


# The Red Supergiant Betelgeuse (Alpha Orionis)



Size of Star

Size of Earth's Orbit

Size of Jupiter's Orbit



First resolved image of a star other than Sun

Distance  
(Hipparcos)  
130 pc (425 lyr)

If Betelgeuse goes Supernova:


- $6 \times 10^7$  neutrino events in Super-Kamiokande
- $2.4 \times 10^3$  neutrons /day from Si burning phase (few days warning!), need neutron tagging  
[Odrzywolek, Misiaszek & Kutschera, astro-ph/0311012]



**Diffuse SN Neutrino Background**


# Diffuse Supernova Neutrino Background (DSNB)

- Approx. 10 core collapses/sec in the visible universe
- Emitted  $\nu$  energy density  
~ extra galactic background light  
~ 10% of CMB density
- Detectable  $\bar{\nu}_e$  flux at Earth  
 $\sim 10 \text{ cm}^{-2} \text{ s}^{-1}$   
mostly from redshift  $z \sim 1$
- Confirm star-formation rate
- Nu emission from average core collapse & black-hole formation
- Pushing frontiers of neutrino astronomy to cosmic distances!



Window of opportunity between  
reactor  $\bar{\nu}_e$  and atmospheric  $\nu$  bkg

# Supernova vs. Star Formation Rate in the Universe



Measured SN rate about half the prediction from star formation rate

Many “dark SNe” ?

Horiuchi, Beacom, Kochanek, Prieto, Stanek & Thompson  
arXiv:1102.1977




# Neutron Tagging in Super-K with Gadolinium

Background suppression: Neutron tagging in  $\bar{\nu}_e + p \rightarrow n + e^+$

- Scintillator detectors: Low threshold for  $\gamma(2.2 \text{ MeV})$
- Water Cherenkov: Dissolve Gd as neutron trap (8 MeV  $\gamma$  cascade)
- Need 100 tons Gd for Super-K (50 kt water)

EGADS test facility at Kamioka

- Construction 2009–11
- Experimental program 2011–2013




Mark Vagins  
Neutrino 2010

Selective water & Gd  
filtration system

200 ton  
water tank      Transparency  
measurement

# Average spectral properties from DSNB



90% CL sensitivity to average SN spectrum from DSNB after 5 years of Gd enhanced Super-K


Adapted from Yüksel, Ando & Beacom, astro-ph/0509297



# Particle-Physics Constraints

# Do Neutrinos Gravitrate?

## Early light curve of SN 1987A



- Neutrinos arrived several hours before photons as expected
- Transit time for  $\nu$  and  $\gamma$  same (160.000 yr) within a few hours

Shapiro time delay for particles moving in a gravitational potential

$$\Delta t = -2 \int_A^B dt \Phi[r(t)]$$

For trip from LMC to us, depending on galactic model,

$$\Delta t \approx 1-5 \text{ months}$$

Neutrinos and photons respond to gravity the same to within

$$1-4 \times 10^{-3}$$

Longo, PRL 60:173, 1988

Krauss & Tremaine, PRL 60:176, 1988

# Millisecond Bounce Time Reconstruction

## Super-Kamiokande

- Emission model adapted to measured SN 1987A data
- “Pessimistic distance” 20 kpc
- Determine bounce time to a few tens of milliseconds

Pagliaroli, Vissani, Coccia & Fulgione  
arXiv:0903.1191

## IceCube




FIG. 1: Typical Monte Carlo realization (red histogram) and reconstructed fit (blue line) for the benchmark case discussed in the text for a SN at 10 kpc.

Halzen & Raffelt, arXiv:0908.2317

# Neutrino Limits by Intrinsic Signal Dispersion

## Time of flight delay by neutrino mass

G. Zatsepin, JETP Lett. 8:205, 1968

$$\Delta t = 2.57s \frac{D}{50 \text{ kpc}} \left( \frac{10 \text{ MeV}}{E_\nu} \right)^2 \left( \frac{m_\nu}{10 \text{ eV}} \right)^2$$

SN 1987A signal duration implies

$$m_{\nu_e} \lesssim 20 \text{ eV}$$

Loredo & Lamb

Ann N.Y. Acad. Sci. 571 (1989) 601

find 23 eV (95% CL limit) from detailed maximum-likelihood analysis

- At the time of SN 1987A competitive with tritium end-point
- Today  $m_\nu < 2.2 \text{ eV}$  from tritium
- Cosmological limit today  $m_\nu \lesssim 0.2 \text{ eV}$

## “Milli charged” neutrinos

Path bent by galactic magnetic field, inducing a time delay

$$\frac{\Delta t}{t} = \frac{e_\nu^2 (B_\perp d_B)^2}{6E_\nu^2} < 3 \times 10^{-12}$$

SN 1987A signal duration implies



$$\frac{e_\nu}{e} < 3 \times 10^{-17} \frac{1\mu\text{G}}{B_\perp} \frac{1 \text{ kpc}}{d_B}$$

- Barbiellini & Cocconi, Nature 329 (1987) 21
- Bahcall, Neutrino Astrophysics (1989)

Assuming charge conservation in neutron decay yields a more restrictive limit of about  $3 \times 10^{-21} e$

# Supernova 1987A Energy-Loss Argument


## SN 1987A neutrino signal



Emission of very weakly interacting particles would “steal” energy from the neutrino burst and shorten it.  
(Early neutrino burst powered by accretion, not sensitive to volume energy loss.)

**Late-time signal most sensitive observable**


# Axion Bounds and Searches





# Diffuse Supernova Axion Background (DSAB)

- Neutrinos from all core-collapse SNe comparable to photons from all stars
- Diffuse Supernova Neutrino Background (DSNB) similar energy density as extra-galactic background light (EBL), approx 10% of CMB energy density
- DSNB probably next astro neutrinos to be measured



- Axions with  $m_a \sim 10$  meV near SN 1987A energy-loss limit
- Provide DSAB with comparable energy density as DSNB and EBL
- No obvious detection channel


Raffelt, Redondo & Viaux  
work in progress (2011)

# Dirac Neutrino Constraints by SN 1987A

- If neutrinos are Dirac particles, right-handed states exist that are “sterile” (non-interacting)
- Couplings are constrained by SN 1987A energy-loss


Right-handed currents		$G_R \lesssim 10^{-5} G_F$
Dirac mass		$m_D \lesssim 30 \text{ keV}$
Dipole moments		$\mu_\nu \lesssim 10^{-12} \mu_B$
Milli charge		$e_\nu \lesssim 10^{-9} e$

# Degenerate Fermi Seas in a Supernova Core



In true thermal equilibrium with flavor mixing, only **one** chemical potential for charged leptons and **one** for neutrinos.  
No chemical potential for Majorana neutrinos (lepton number violation)

# Degenerate Fermi Seas in a Supernova Core



Equilibration by flavor lepton number violation, but flavor oscillations ineffective (matter effect)

Non-standard interactions could be effective, most sensitive environment

Consequences in core collapse should be studied numerically

Equilibration by lepton number violation, but Majorana masses too small

R-parity violating SUSY interactions?  
TeV-scale bi-leptons?

# TeV-scale bileptons, see-saw type II and lepton flavor violation in core-collapse supernova

Oleg Lychkovskiy<sup>1,a</sup>, Sergei Blinnikov<sup>1,2</sup>, Mikhail Vysotsky<sup>1</sup>


<sup>1</sup>Institute for Theoretical and Experimental Physics, B. Cheremushkinskaya 25, 117218 Moscow, Russia

<sup>2</sup>IPMU, University of Tokyo, 5-1-5 Kashiwanoha, Kashiwa, 277-8568, Japan

Received: 7 January 2010 / Published online: 31 March 2010

© Springer-Verlag / Società Italiana di Fisica 2010

**Abstract** Electrons and electron neutrinos in the inner core of the core-collapse supernova are highly degenerate and therefore numerous during a few seconds of explosion. In contrast, leptons of other flavors are non-degenerate and therefore relatively scarce. This is due to lepton flavor conservation. If this conservation law is broken by some non-standard interactions,  $\nu_e$  are converted to  $\nu_\mu$ ,  $\nu_\tau$ , and  $e$  are converted to  $\mu$ . This affects the supernova dynamics and the supernova neutrino signal. We consider lepton flavor violating interactions mediated by scalar bileptons, i.e. heavy scalars with lepton number 2. It is shown that in case of TeV-mass bileptons the electron Fermi gas is equilibrated with non-electron species inside the inner supernova core at a time scale  $\sim(1-100)$  ms. In particular, a scalar triplet which generates neutrino masses through the see-saw type II mechanism is considered. It is found that the supernova core is sensitive to yet unprobed values of masses and couplings of the triplet.




**Fig. 1**  $ee \rightarrow \mu\mu$  LFV transition mediated by the doubly charged bilepton  $\Delta^{--}$



# Neutrino Flavor Oscillations

# Flavor Oscillations in Core-Collapse Supernovae

Neutrino-neutrino refraction causes a flavor instability, flavor exchanged between different parts of spectrum





**More on Thursday ...**