




# Understanding the Spherically Symmetric Gamma-Ray Residual at the Galactic Center



Tim Linden  
UC - Santa Cruz

with Dan Hooper, Elizabeth Lovegrove,  
Stefano Profumo and Farhad Yusef-Zadeh

Closing in on Dark Matter - Aspen January 30, 2012



# The J-Factor of the Galactic Center

Ackermann et al. 2012

## Dwarfs

Name	l deg.	b deg.	d kpc	$\overline{\log_{10}(J)}$ $\log_{10}[\text{GeV}^2\text{cm}^{-5}]$	$\sigma$	ref.
Bootes I	358.08	69.62	60	17.7	0.34	[15]
Carina	260.11	-22.22	101	18.0	0.13	[16]
Coma Berenices	241.9	83.6	44	19.0	0.37	[17]
Draco	86.37	34.72	80	18.8	0.13	[16]
Fornax	237.1	-65.7	138	17.7	0.23	[16]
Sculptor	287.15	-83.16	80	18.4	0.13	[16]
Segue 1	220.48	50.42	23	19.6	0.53	[18]
Sextans	243.4	42.2	86	17.8	0.23	[16]
Ursa Major II	152.46	37.44	32	19.6	0.40	[17]
Ursa Minor	104.95	44.80	66	18.5	0.18	[16]

- Corresponds to the relative annihilation rate of the region compared to other astrophysical sources

$$\Phi_\gamma \propto J = \frac{1}{\Delta\Omega} \int d\Omega \int_{\text{l.o.s.}} \rho^2(l) dl(\psi)$$

- The J-factor of the galactic center is approximately:

$$\log_{10}(J) = 21.0$$


for a region within  $1^\circ$  of the Galactic center and an NFW profile

Ackermann et al. 2010

## Clusters

Cluster	RA	Dec.	z	J ( $10^{17} \text{ GeV}^2 \text{ cm}^{-5}$ )
AWM 7	43.6229	41.5781	0.0172	$1.4^{+0.1}_{-0.1}$
Fornax	54.6686	-35.3103	0.0046	$6.8^{+1.0}_{-0.9}$
M49	187.4437	7.9956	0.0033	$4.4^{+0.2}_{-0.1}$
NGC 4636	190.7084	2.6880	0.0031	$4.1^{+0.3}_{-0.3}$
Centaurus (A3526)	192.1995	-41.3087	0.0114	$2.7^{+0.1}_{-0.1}$
Coma	194.9468	27.9388	0.0231	$1.7^{+0.1}_{-0.1}$




# Subtracting the Astrophysical Background: Fermi



- We employ a model of the galactic gas density (Kalberla & Kerp 2009) to subtract the contributions from the galactic plane.
- This emission template provides a superb match to the total emission spectrum
- This large residual at the center of the galaxy is a factor of 10 brighter than anything else in the inner  $20^\circ \times 10^\circ$

# Is it a Point Source?

- Several efforts have been made to fit the GC point source, using both best-fitting point-source tools from the Fermi collaboration (Boyarsky et al. Chernyakova et. al), as well as independent software packages (Hooper & Goodenough)
- In all cases, the morphology of the observed emission cannot be fully accounted for by a single point source smeared out by the angular resolution of the Fermi-LAT




Hooper & Linden (2011)

# Independent Confirmation!

- Abazajian & Kaplinghat employed a more sophisticated template-based regression analysis
- This also found an extremely significant improvement in the overall fit with the addition of a spherical profile with similar characteristics to that of Hooper & Goodenough and Hooper & Linden

See talk by Kev  
Abazajian

Abazajian & Kaplinghat (2012)





# Independent Confirmation!

- Note: Two different, and independent methods find strong evidence for a **bright, spatially extended, spherically symmetric residual** at the position of the galactic center
- What can we learn from this?

# A Hadronic Scenario



- The HESS spectrum is well fit by the Fermi acceleration of protons and their subsequent interaction with galactic gas
- Can the combined Fermi + HESS spectrum be described in the same way?
- **Problem 1:** The spectrum at GeV energies is significantly softer than at TeV energies - some modification is needed to control this transition
- **Problem 2:** The H.E.S.S. spectrum is point-like, with a better angular resolution than Fermi-LAT



Acero et al. (2010)

# Controlling the Emission Spectrum with Diffusion

- We can imagine two scenarios for cosmic-ray transport from the central black hole: rectilinear or diffusive transportation
- In the regime where the diffusion stepsize exceeds the diffusion region, the emission intensity is energy independent, and an  $E^{-2}$  proton injection spectrum corresponds directly to an  $E^{-2}$  gamma-ray spectrum




- In the regime where the diffusion step is small, then the emission intensity depends linearly on the time the particle spends within the diffusion region



# Hadronic Emission Models for Fermi and HESS


- By setting allowing the diffusion constant to float to a set of best fit values - a single hadronic emission model can fit the entirety of the Fermi/HESS data



- Several model parameters can also be adjusted, such as the duration of particle injection, the occurrence of recent flares, the maximum radius for diffusion etc.
- Models are formed with a step-function gas density profile ( $1000 n_{\text{H}}/\text{cm}^{-3}$  within 3 pc of the galactic center, and  $0 n_{\text{H}}/\text{cm}^{-3}$  outside)


# Employing a Realistic Gas Model

- Detailed models of the galactic gas density exist in the literature
- We employ a spherically symmetric model for galactic gas, and use this to calculate the morphology of the gamma-ray emission as a function of energy
- By far the dominant feature is the Circumnuclear ring between 1-3 pc from the GC




# Employing a Realistic Gas Model

- The vast majority of emission stems from within 3 pc of the galactic center at all energies
- This lies below the PSF of all current gamma-ray instruments
- This effectively rules out hadronic interactions from Sgr A\* as the source of the Fermi-LAT excess




Linden et al. (2012)




# CTA and the Galactic Center


- By convolving our models of the gas and proton densities in the galactic center region with the PSF and effective area of each instrument, we can determine whether CTA can distinguish between these scenarios
- CTA will conclusively determine whether the galactic center source stems from a hadronic emission channel



# Understanding High Energy Emission from the Galactic Center: 2 Convincing Stories





Linden et al. (2012)



# Story 2: Low-Mass Dark Matter


- For a best fitting profile  $\gamma = 1.3$ , we find an available parameter space for dark matter models which match the observed GC excess
- These models are compatible with estimates for the relic density of dark matter



- The models combine with best fitting astrophysical backgrounds such as the GC point source and the galactic ridge, to fit the total GC excess



Hooper & Linden (2011)

# Dark Matter Limits in the Simplest Way Possible




Hooper & Linden (2011)



- After subtracting emission from known point sources, and an extrapolation of the line-of-sight gas density, the following "galactic center" emission is calculated
- This directly corresponds to a limit on the dark matter interaction cross-section which depends only on assumed dark matter density profile



# Comparison to Other Indirect Detection Regimes




- Hooper et al. (2012) further tweaked the methods used to derive these limits, deriving rigorous constraints under a wide variety of assumptions
- These are the strongest gamma-ray limits on the cross-section for dark matter annihilation






# Comparison to Other Indirect Detection Regimes



Hooper & Linden (2011)



- With some adiabatic contraction of the inner dark matter profile, these limits can become substantially stronger than any other indirect detection limit



Ackermann et al. (2011)

# Story 3: Milli-second Pulsars



- Populations of Millisecond pulsars have been observed in multiple globular clusters (Terzan 5, Omega Cen, NGC 6388, M 28)
- GC source is ~200 brighter than Omega Cen - which correlates nicely with the 1000x larger mass of the GC region
- Spectrum of MSP population is very similar to the observed gamma-ray excess



Hooper & Linden (2011) Spectral Index  $\Gamma$

# Story 3: Milli-second Pulsars

- The galactic center residual spectrum ( $\Gamma < \approx 1.0$ ) is somewhat harder than the population of observed pulsars - though uncertainties in the astrophysical spectrum which is subtracted are uncertain
- Must explain the high density of pulsars near the Galactic Center ( $\sim r^{-2.6}$ )
- Two body interactions in the densest clusters?
- Mass segregation?




Voss & Gilfanov (2008)

# Hope at Separating these Models?

- The expected gamma-ray signatures of MSPs and light dark matter annihilation in the galactic center are very similar
- Fair Statement: Dark matter provides a slightly better statistical fit than MSPs. However our bayesian prior (that MSPs emit in the GC) may be higher than for dark matter
- Need new techniques or observations to differentiate these signals


# The Multi-wavelength Galactic Center




VLA



Chandra



EGRET



HESS



Fermi-LAT



**Extinction > 10**  
Fritz et al. 2011

Regis & Ullio 2009

# Angular Scales of the Galactic Center

$\text{I} = \times 100 \text{ sr}$


BH  
VLA  
Chandra




CTA  
HESS

Fermi (100 GeV)  
Fermi (1 GeV)

# The Galactic Center "Zoo"




# WMAP Haze??






- The same dark matter model provides a reasonable explanation to the intensity and morphology of the WMAP haze
- The magnetic field must be slightly stronger above the galactic plane than usually assumed

Dobler et al. (2007)





# WMAP Haze??




- However, correlating the WMAP haze against the Fermi bubbles shows a clear edge
- Furthermore, multiple galaxies are underluminous in radio emission, compared to expectations from dark matter annihilation

# Filamentary Arcs

- Bright Radio Synchrotron sources near (<100 pc) the Galactic Center
- Polarization measurements imply the magnetic field is highly ordered

$$B_{\text{tot}} \sim 50-1000 \mu\text{G}$$
$$\frac{B_{\text{ord}}^2}{B_{\text{tot}}^2} > 0.6$$



- Mechanism of filament creation and emission is unknown



Yusef-Zadeh et al. (2004)

# Filamentary Arcs

- The same dark matter model also provides a fit to the spectrum and intensity of the filamentary arcs
- Light DM annihilation naturally provides the near delta-function electron spectrum necessary to explain the synchrotron spectrum of the filaments



- Dark matter injection also naturally predicts an  $r^{-2}$  trend for the flux of filaments which are farther from the GC, an implication not shared by local astrophysical sources

# Filamentary Arcs

$$E_M = 7 \text{ GeV} \left( \frac{V_A}{2000 \text{ km s}^{-1}} \right)^2 \left( \frac{B^2/8\pi}{8 \cdot 10^{-6} \text{ erg cm}^{-3}} \right)^{-1} \left( \frac{K_{\parallel}}{10^{24} \text{ cm}^2 \text{ sec}^{-1}} \right)^{-1} \quad (7a).$$

ASTRONOMY  
AND  
ASTROPHYSICS

1988ARA...1002...19

Astron. Astrophys. 200, L9-L12 (1988)

## Letter to the Editor

**Monoenergetic relativistic electrons in the galactic center**

H. Lesch\*, R. Schlickeiser, and A. Crusius

Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-5300 Bonn 1, Federal Republic of Germany

Received March 29, accepted May 27, 1988

### Summary

It is shown that the nonthermal radio spectra of the center, including Sgr A\* and the extended component (Arc) is neither due to self-absorbed nor due to thermal absorption. A function which propagates into the




$$\delta\theta_{\text{crit}} = 2.6 \cdot 10^9 S_M^{1/2} v_M^{-5/4} B^{1/4} \text{ arcsec}$$

where  $S_M$  is the observed flux density of the self-absorbed source at a frequency of 10 GHz (Reich et al., 1988) and  $v_M$  is the magnetic field. With the flux density of  $10^{-2}$  G (Sofue and Fujimoto, 1988) we

$$\delta\theta_{\text{crit}} = 4 \cdot 10^{-4} \text{ arcseconds}$$

The source is resolved with an angular size of several arcseconds (Reich et al., 1988) and probably small structures


# Arcade Excess?



- Arcade-2 Collaboration noted a hard synchrotron residual in low frequency (<10 GHz radio data)
- Emission is hard to account for with known astrophysical sources
- Can be accounted for with **light** dark matter annihilation

# Can the Distribution of GC MSPs be Determined?



- X-Ray observations find a total of 2347 point sources within 40 pc of the GC - this could include a large population of MSPs
- MSPs exist in a particular location on the luminosity-color diagram in 47 Tuc
- Can this information be used to determine the statistical distribution of MSPs?



Heinke et al. (2006)

# Diffuse Secondary Emission

- Another method for distinguishing between gamma-ray emission models is to investigate the production of electron and positron pairs
- These charged leptons will lose considerable energy to synchrotron radiation, producing a bright radio signal in the galactic center



Positive: The angular resolution of radio telescopes is significantly greater than gamma-ray observatories

Negative: The diffusion and energy loss time of charged electrons adds additional uncertainties to the model

# Conclusions

- There is **strong** evidence for an extended, spherically symmetric, excess in  $\sim 1$  GeV gamma-ray emission surrounding the galactic center
- This excess is not easily accounted for by any known astrophysical model - and the background subtraction models used indicate that it is not correlated with galactic gas
- Dark Matter Annihilation and Pulsars both provide plausible models for this excess
- New observations, and also novel models, are needed to separate these components





# Extra Slides

# Best fitting Models for Low-Mass Dark Matter


- Abazajian & Kaplinghat find a wider range of dark matter masses which provide improved fits to the data
- However, fits with low dark matter mass are much, much better

TABLE II. The best-fit TS, negative log likelihoods, and  $\Delta\mathcal{L}$  from the baseline, for specific dark matter channel models, using the  $\alpha\beta\gamma$  profile (Eq. 2.1) with  $\alpha = 1, \beta = 3, \gamma = 1.2$ .



channel, $m_\chi$	TS	$-\ln \mathcal{L}$	$\Delta \ln \mathcal{L}$
$b\bar{b}$ , 10 GeV	2385.7	139913.6	156.5
$b\bar{b}$ , 30 GeV	3460.3	139658.3	411.8
$b\bar{b}$ , 100 GeV	1303.1	139881.1	189.0
$b\bar{b}$ , 300 GeV	229.4	140056.6	13.5
$b\bar{b}$ , 1 TeV	25.5	140108.2	-38.0
$b\bar{b}$ , 2.5 TeV	7.6	140114.2	-44.0
$\tau^+\tau^-$ , 10 GeV	1628.7	139787.7	282.5
$\tau^+\tau^-$ , 30 GeV	232.7	140055.9	14.2
$\tau^+\tau^-$ , 100 GeV	4.10	140113.4	-43.3



# Other Observations Fitting Light DM: Direct



- Light Dark Matter ( $\sim 10$  GeV) provides a compelling fit to the excesses currently observed by DAMA, CoGeNT and CRESST
- Light Dark Matter may also be compatible with observed signal/limits at CDMS
- However, a recent error found in CoGeNT analysis may affect some early dark matter interpretations



# Independent Confirmation!

- Abazajian & Kaplinghat employed a more sophisticated template-based regression analysis
- This also found an extremely significant improvement in the overall fit with the addition of a spherical profile with similar characteristics to that of Hooper & Goodenough and Hooper & Linden

Spatial Model	Spectrum	TS	$-\ln \mathcal{L}$	$\Delta \ln \mathcal{L}$
Baseline	—	—	140070.2	—
Density $\Gamma = 0.7$	LogPar	1725.5	139755.5	314.7
Density <sup>2</sup> $\gamma = 0.9$	LogPar	1212.8	139740.0	330.2
Density <sup>2</sup> $\gamma = 1.0$	LogPar	1441.8	139673.3	396.9
Density <sup>2</sup> $\gamma = 1.1$	LogPar	2060.5	139651.8	418.3
Density <sup>2</sup> $\gamma = 1.2$	LogPar	4044.9	139650.9	419.2
Density <sup>2</sup> $\gamma = 1.3$	LogPar	7614.2	139686.8	383.4
Density <sup>2</sup> Einasto	LogPar	1301.3	139695.7	374.4
Density <sup>2</sup> $\gamma = 1.2$	PLCut	3452.5	139663.2	407.0


TABLE II. The best-fit TS, negative log likelihoods, and  $\Delta \mathcal{L}$  from the baseline, for specific dark matter channel models, using the  $\alpha\beta\gamma$  profile (Eq. 2.1) with  $\alpha = 1, \beta = 3, \gamma = 1.2$ .

channel, $m_\chi$	TS	$-\ln \mathcal{L}$	$\Delta \ln \mathcal{L}$
$b\bar{b}$ , 10 GeV	2385.7	139913.6	156.5
$b\bar{b}$ , 30 GeV	3460.3	139658.3	411.8
$b\bar{b}$ , 100 GeV	1303.1	139881.1	189.0
$b\bar{b}$ , 300 GeV	229.4	140056.6	13.5
$b\bar{b}$ , 1 TeV	25.5	140108.2	-38.0
$b\bar{b}$ , 2.5 TeV	7.6	140114.2	-44.0
$\tau^+\tau^-$ , 10 GeV	1628.7	139787.7	282.5
$\tau^+\tau^-$ , 30 GeV	232.7	140055.9	14.2
$\tau^+\tau^-$ , 100 GeV	4.10	140113.4	-43.3


Abazajian & Kaplinghat (2012)

# HESS Limits on TeV Dark Matter


- HESS observations of the Galactic center, and Galactic Halo provide the strongest indirect limits on TeV dark matter
- Limits are strongly profile dependent -- background subtraction weakens bounds on isothermal dark matter models as well



Abazajian & Harding (2011)



Abramowski et al. (2011)




# Fermi Telescope (2008-Present)



- Fermi-LAT is a space based gamma-ray detector with an effective energy range of 20 MeV-300 GeV





- Effective Area  $\sim 0.8 \text{ m}^2$
- Field of View  $\sim 2.4 \text{ sr}$
- Energy Resolution  $\sim 10\%$
- Angular Resolution: Energy Dependent



- In analyses of the Galactic Center, we will constrict ourselves to Front converting events


# Understanding Astrophysical Backgrounds: HESS

- HESS spectrum well matched by flat  $E^{-2}$  spectrum, up to energies of  $\sim 10$  TeV, where an exponential cutoff is observed
- HESS source is localized to within  $13''$  of Galactic center (solid white curve) - the 68% and 95% confidence levels on the source extension are at  $\sim 1$  and 3 pc



# CTA and the Galactic Center

- By convolving our models of the gas and proton densities in the galactic center region with the PSF and effective area of each instrument, we can determine whether CTA can distinguish between these scenarios
- CTA will conclusively determine whether the galactic center source stems from a hadronic emission channel





Linden & Profumo  
(2012)



# Understanding Astrophysical Backgrounds: HESS

- However, HESS shows no variability, even during outbursts observed by Chandra
- This implies that the source of the emission is spatially distinct from lower energy sources



Aharonian et al. (2008)



Aharonian et al. (2009)


# HESS Telescope (2004-Present)

- HESS is an Atmospheric Cherenkov Telescope built in Namibia
- Effective over the energy range  $\sim 500$  GeV - 100 TeV with an effective area on the order of  $10^5$  m<sup>2</sup>.
- Energy Resolution  $\sim 10\%$
- Angular Resolution ( $>1$  TeV)  $\sim 0.075^\circ$ .
- Total Observation of the Galactic Center: 93h/112h





# And some surprises!

$I = x10^{20}$



BH  
VLA  
Chandra

Fermi Bubbles? Do they extend to the galactic center?




Non-thermal Radio Filaments - Bright, polarized synchrotron sources shaped like "thin threads" and lying perpendicular to galactic plane (Yusef-Zadeh et al. 1984)

CTA  
HESS



Dark Matter??

Fermi (100 GeV)  
Fermi (1 GeV)

# Comparison to Other Indirect Detection Regimes




- Under the assumption of an NFW profile, the 95% confidence limits are as good or better than those from dwarf-spheroidals
- They are especially stronger for leptophilic annihilation paths




# Employing a Realistic Gas Model

But CTA may be able to probe this emission profile directly!





Linden et al. (2012)





# CTA and the Galactic Center

- However, CTA may be able to distinguish between these models:
- The instrument specifications for CTA are not yet entirely known, so we employ the following:
  - An order of magnitude improvement in the effective area over HESS
  - A reduction in the PSF from 1-10 TeV from  $0.075^\circ$  to  $0.03^\circ$



# Variability at the Galactic Center

- Sgr A\* is highly variable (on multiple time scales) at both radio and X-Ray energies



## Motivating Question:

Why would the  
galactic center be an  
interesting place to  
look for Dark Matter?



Slides Courtesy of G. Zaharijas

Diemand et al. 2008





# Fitting the Residual: Hadronic Processes

- The lack of variability indicates that the emission may be stemming from a region farther away from the GC itself
- A recent model examined the possibility that protons emitted from the galactic center produce gamma-rays through their subsequent interaction with galactic gas
- This has the potential to produce the vast majority of emission from TeV scales all the way down to radio energies
- Normalization depends sensitively on diffusion (**stay tuned!**)




# Fitting the Residual: Hadronic Processes

- A recent model examined the possibility that protons injected from the galactic center encountered the circumnuclear ring
- This region of high density molecular gas would produce bright gamma-ray emission upon the interaction with energetic protons




Ferriere (2012)



Ballantyne et al. 2007



# Negative: The Profile Dependence



- Assumptions for the slope of the inner dark matter profile can make **orders of magnitude** differences in the expected dark matter annihilation rate
- Dark Matter is not a dominant gravitational source near the galactic center, so there are few observational handles on the dark matter density in the GC region

# Positive! Progress in Simulations

- Simulations including the effects of baryonic contraction show a steepening of the spectral slope from  $\gamma \approx 1.0$  to  $\gamma \approx 1.2-1.5$
- Much more work is required to understand the dark matter content of the GC region
- This is imperative for understanding the signals from indirect detection





as reported in Gnedin et al. 2011

# History of Galactic Center Observations (in 60 seconds)

Muno et al. 2007


- Sgr A\* Discovered via radio observations in 1974
- Measurements of stellar motion confirm the status of the central object as a black hole (Gillissen et al. 2009)
- Majority of radio emission thought to stem from accretion disk, rather than at BH event horizon (Doeleman et al. 2008)




Gillissen et al. 2009

# Dark Matter Indirect Detection

## Particle Physics




Slides Courtesy of G. Zaharijas



## Instrumental Response

## Astrophysics




Diemand et al. 2008

# What is the WMAP Haze?

Hooper & Goodenough (2011)



- To determine the best-fit dark matter annihilation profile, Hooper & Goodenough bin the residuals as a function of radius
- Then the residual as a function of radius can be compared with the dark matter injection profile convolved with the PSF of the Fermi-LAT



# What is the WMAP Haze?

- Discovered by Doug Finkbeiner in 2004
- Synchrotron origin determined by subsequent observations
- Hard spectrum difficult to fit with lepton injection spectra typical of astrophysical phenomena
- Well fit by dark matter models with typical annihilation cross-sections and spectra
- However, modifications are needed to magnetic fields in galactic halo



Dobler et al. (2008)





# The Radial Dependence of the Filamentary Arcs



- The intensity of multiple filamentary arcs show a strong dependence on their distance from the galactic center
- This is expected in dark matter models, but not in most astrophysical interpretations of the filaments



Linden et al. (2011)

# Dark Matter at the Galactic Center

- Can use a Kolmogorov-Smirnov test after finding the CDF for the radial profile of dark matter annihilation
- Since the CDFs for dark matter and the background point-source can be compared linearly, strong limits can quickly be set on dark matter annihilation
- Limits on photon counts can then be translated to a limit on annihilation cross-section
- Of course, large uncertainties exist, stemming from models in the gas density, and in the ratio of background emission stemming from point-source vs. gas




# Modeling Benefits of the Hadronic Scenario!



- Under the assumption that the proton source has a power-law spectrum and is in steady-state, then the slope of gamma-ray emission strongly constrains the diffusion constant in the galactic center region:

$$D_0 = 1.2 \times 10^{26} (E/1 \text{ GeV})^{0.91}$$

- This adds additional constraints to the an understanding of lepton diffusion and propagation in the galactic center region




# Models of the Galactic Center Magnetic Field




- This is particularly interesting in light of recent models which have set a minimum strength of  $50 \mu\text{G}$  on the magnetic fields in the galactic center (best fit range  $100\text{-}300 \mu\text{G}$ )
- This almost ensures that synchrotron is the dominant energy loss mechanism for high energy electrons
- In the hadronic scenario, the diffusion parameters are set by the fit to the gamma-ray data

**Note:** Models of light dark matter and millisecond pulsars seek only to explain the bump in the Fermi GeV spectrum.

In both cases, another mechanism (such as proton emission from the galactic center) must be responsible for the TeV emission



Hooper & Linden (2011)



# Conclusions - Galactic Center

- The galactic center is one of the most exciting places to search for a dark matter signal
- Present observatories are capable of both making exciting discoveries, and setting stringent limits on the properties of WIMP dark matter
- Upcoming instruments are likely to make exciting discoveries of both the astrophysical and dark matter properties of the galactic center region