



Subthreshold ϕ meson production in nucleus-nucleus collisions

Krzysztof Piasecki for the FOPI Collaboration

Institute of Experimental Physics, University of Warsaw, Poland

- ϕ mesons in nucleus-nucleus collisions
- Production of ϕ at 1.5 .. 2A GeV: yields and slopes
- Systematics of data
- Summary and outlook




ϕ mesons in nucleus-nucleus collisions

- $\phi(s\bar{s})$: $m = 1019$ MeV
- $E_{\text{thr NN}} = 2.6$ GeV (below threshold for the GSI experiments)
- Considered production channels for subthreshold energies :



$$\begin{aligned} BB &\rightarrow \phi BB, \quad B = [\Delta, N] \\ \pi B &\rightarrow \phi B, \quad \rho B \rightarrow \phi B \\ \pi N^* &\rightarrow \phi N, \quad \rho \pi \rightarrow \phi \end{aligned}$$

H.W. Barz et al. Nucl. Phys. A 705 (2002) 223


- $c\tau = 50$ fm
- $\phi \rightarrow K^+ K^-$ (BR = 49%)
- Yield of K^- is comparable to ϕ



FOPI experimental setup




- Nearly 4π coverage
- Drift chambers: CDC, Helitron
- ToF : Plastic Barrel, RPC
- Forward: Plastic Wall, Zero Degree
- Direct PID of π^\pm , K^\pm , p , d , t , ${}^{3,4}\text{He}$



Ni+Ni @ 1.93A GeV

- First measurement: FOPI, 1995
- Trigger: 12% most central events




A. Mangiarotti et al, NPA 714 (2003) 89

- Number of $\phi = 23 \pm 7 \pm 2$
- ϕ yield depends on Temperature, which was not found due to scarce statistics.
- But within $T \in [70, 130]$ MeV, $P(\phi) = (1.2 .. 4.5) \cdot 10^{-3}$ (first estimate)


Ni+Ni @ 1.93A GeV

- Trigger: 20% most central events



- Number of $\phi = 100 \pm 17 \pm 20$
- ϕ yield per triggered event:
 $(6 \pm 1 \pm 2) \cdot 10^{-4}$

- K^- from KaoS, central collisions




- K^- yield per triggered event:
 $(2.1 \pm 0.4) \cdot 10^{-3}$

$$\rightarrow \frac{\phi}{K^-} = 29 \pm 7 \pm 10\%$$


$\rightarrow 14 \pm 3\%$ of K^- originate from ϕ decays

Ni+Ni @ 1.93A GeV

- Trigger: 50% most central events



- Number of $\phi = 210$
Significance = 9.6
- ϕ yield per triggered event:
 $(4.4 \pm 0.3) \cdot 10^{-4}$




- K^+ yield per triggered event:
 $(3.87 \pm 0.10 \pm 0.12) \cdot 10^{-2}$
- K^- / K^+ ratio (in the measurable range)
 $(3.32 \pm 0.07 \pm 0.13) \cdot 10^{-2}$

$$\rightarrow \frac{\phi}{K^-} = 34 \pm 3 \pm 2.5 \%$$


$$\rightarrow 17 \pm 2 \% \text{ of } K^- \text{ originate from } \phi \text{ decays}$$

Al+Al @ 1.9A GeV

- Trigger: 9% most central events



- Number of $\phi = 108 \pm 15$
Significance = 7.4
- ϕ yield per triggered event:
 $(3.3 \pm 0.5 \pm 0.6) \cdot 10^{-4}$




- Number of $K^- = 3090 \pm 60$
- K^- yield per triggered event:
 $(1.1 \pm 0.3 \pm 0.2) \cdot 10^{-3}$



→ $\frac{\phi}{K^-} = 30 \pm 8 \pm 5 \%$

→ $15 \pm 4 \%$ of K^- originate from ϕ decays


Ar + KCl @ 1.756A GeV



- Trigger: 35% most central events



- Number of $\phi = 168 \pm 18$
- ϕ yield per triggered event:
 $(2.6 \pm 0.7 \pm 0.1) \cdot 10^{-4}$




- K^- yield per triggered event:
 $(7.1 \pm 1.5 \pm 0.3) \cdot 10^{-4}$

$$\Rightarrow \frac{\phi}{K^-} = 37 \pm 13 \pm 3 \%$$

$$\Rightarrow 18 \pm 6 \% \text{ of } K^- \text{ originate from } \phi \text{ decays}$$

Systematics of ϕ meson production




- Conclusion: regardless of system size and exact beam energy,

$$\frac{\phi}{K^-} \approx \frac{1}{3}$$



About 15 .. 20% of K^- are produced from ϕ decays

ϕ mesons: inverse slopes




Al+Al @ 1.9A GeV

$$T_B = 83 \pm 26 \text{ MeV}$$

Assuming:

$$T_B(y) = \frac{T_{\text{eff}}}{\cosh(y - y_{CM})}$$

$$T_{\text{eff}} = 91 \pm 20 \text{ MeV}$$




Ar+KCl @ 1.9A GeV

Assuming:

$$T_B(y) = \frac{T_{\text{eff}}}{\cosh(y - y_{CM})}$$


$$T_{\text{eff}} = 84 \pm 8 \text{ MeV}$$



Ni+Ni @ 1.9A GeV

50% most central σ

$$T = 92 \pm 12 \text{ MeV}$$



Ni+Ni @ 1.9A GeV

12% most central σ

$$T = 120 \pm 12 \pm 20 \text{ MeV}$$

Summary and outlook

- At $E_{beam} = 1.5 \dots 2A$ GeV, ϕ mesons from five data sets were measured:
 - FOPI: Al+Al and Ni+Ni (3 different centrality classes) @ 1.9A GeV
 - HADES: Ar+KCl @ 1.756A GeV
- ϕ meson yields: $(0.3 \dots 1.3) \cdot 10^{-3}$ per triggered event
 - $P(\phi)$ seems to scale with $\langle A_{\text{participants}} \rangle_b$
- $\frac{\phi}{K^-}$ ratio seems to be stable ($\frac{1}{3}$) and independent from E_b , $\langle A_{\text{part}} \rangle$.
 - About 15 .. 20% K^- originate from decays of ϕ
- First insights into the ϕ phase space:
 - Inverse slopes of ϕ : 80 .. 120 MeV
- To-do:
 - Final verification of ϕ data from Ni+Ni collisions
 - evaluation of systematic errors

*Thank
You!*

Backup slides

Strangeness production and absorption


	K⁺	K⁻	φ
<i>Production (primary)</i>	$BB \rightarrow BYK^+$ $T_{pp \rightarrow p\Lambda K^+} = 1.58 \text{ GeV}$	$BB \rightarrow BBK^+K^-$ $T_{pp \rightarrow ppK^+K^-} = 2.5 \text{ GeV}$	$BB \rightarrow BB\phi$ $T_{pp \rightarrow ppK^+K^-} = 2.6 \text{ GeV}$
<i>Production (secondary)</i>	$\pi B \rightarrow YK^+$	$\pi Y \rightarrow (\Sigma^* \rightarrow) BK^-$ $BY \rightarrow NK^-\Lambda$ $BY \rightarrow BBK^-$ $\pi B \rightarrow BK^+K^-$ $\phi \rightarrow K^+K^-$	$\pi B \rightarrow B\phi$ $\rho B \rightarrow B\phi$ $\pi N^* \rightarrow N\phi$ $\rho\pi \rightarrow \phi$ $K^+K^- \rightarrow \phi \text{ negligible}$
Absorption	$K^+Y \rightarrow \pi B$	$K^-B \rightarrow \pi Y$	$\phi N \rightarrow K\Lambda$
Elastic scat. (char. exch.)	$K^+B \leftrightarrow K^+ B$ $K^+n \leftrightarrow K^0 p$	$K^-B \leftrightarrow K^-B$ $K^-p \leftrightarrow \bar{K}^0 n$	$\phi N \rightarrow \phi N$

$$[B] = p, n, N, N^*, \Delta$$

$$[Y] = \Lambda, \Sigma$$


Yields from	Ni + Ni (1.93 GeV)
B + B	3.5×10^{-4}
$\pi + B$	2.9×10^{-4}
$\rho + B$	8.9×10^{-4}
$\pi + \rho$	1.6×10^{-4}
$\pi + N(1520)$	0.5×10^{-4}
Total yield	1.7×10^{-3}

H.W. Barz et al. (BUU),
Nucl. Phys. A 705 (2002) 223



C.B. Dover, G.E. Walker
Phys. Rep. **89** (1982) 1

ϕ/K^- within the statistical model approach




J. Cleymans et al. PLB **603**, 146 (2004)


G. Agakishiev et al., PRC **80**, 025209 (2009)

Examples of mixing of different sources of K-

- ϕ influences investigation of U_{KN} potential



- K- flow and ϕ flow mix together



KAOS: F. Uhlig et al., PRL 95, 012301(2005)


First ϕ 's from Ni+Ni : 1995 session

- Ni+Ni @ 1.93A GeV

(A. Mangiarotti)


→ Trig: 12% most centr. ($\langle A_{part} \rangle = 86$)

→ $N_\phi = 23 \pm 7 \pm 2$




→ $P_\phi / \text{collision}$: depends on T_ϕ

$\times 10^{-2}$



$$T_\phi = 70 \text{ MeV} \rightarrow (4.5 \pm 1.4 \pm 2.2) \cdot 10^{-3}$$

$$T_\phi = 130 \text{ MeV} \rightarrow (1.2 \pm 0.4 \pm 0.6) \cdot 10^{-3}$$




First (but only) ϕ data published:


A. Mangiarotti et al, NPA 714 (2003) 89

Phase space of ϕ mesons

Ni+Ni @ 1.9A GeV
50% most central σ




Al+Al @ 1.9A GeV




K-/K+ ratio for Ni+Ni from FOPI and KaoS

FOPI




KaoS



M. Menzel et al. (KaoS), Phys Lett B 495 (2000) 26

Azimuthal anisotropy of Kaons from Ni+Ni @ 1.93A GeV

$$R(\theta_{CM}) = \frac{\sigma_{measured}(\theta_{CM})}{\sigma_{Isotropic}(\theta_{CM})}$$



M. Menzel et al. (KaoS), Phys Lett B 495 (2000) 26