


Future Jets at RHIC with sPHENIX



Ali Hanks

for the PHENIX Collaboration

August 23rd, 2012


Jet Modification in the RHIC and LHC era



Two questions to answer

- Can we learn from full jet measurements at RHIC in the LHC era?
- Can we measure jets at RHIC?
 - untriggered and at high rates


Probing properties of the QGP




- Measurements integrate over time evolution of medium
 - all properties are varying: T , η/s , q -hat, e -hat
- How can we unambiguously see changes near T_c ?
 - run through same time history at different T_0

Investigating the QGP along multiple axes

What is the temperature dependence for the QGP?




What are the inner workings (quasi-particles, strong fields, ...)?




(length scale probed)

$Q^2(t)$
(virtuality of hard process)


How does the medium evolve?



Jet observables at RHIC




Models constrained by LHC data



Sensitivity to *effective* coupling

Sensitivity to interaction of jet with medium

Sensitivity to the effective constituents of the QGP?




sPHENIX stage I design

Make use of new technologies to get something that is:


compact (cheap)

uniform

fast



Rates at RHIC for 20 week Run



	Au+Au central 20%	p+p	d+Au
>20 GeV	10 ⁷ jets 10 ⁴ photons	10 ⁶ jets 10 ³ photons	10 ⁷ jets 10 ⁴ photons
>30 GeV	10 ⁶ jets 10 ³ photons	10 ⁵ jets 10 ² photons	10 ⁶ jets 10 ³ photons
>40 GeV	10 ⁵ jets	10 ⁴ jets	10 ⁵ jets
>50 GeV	10 ⁴ jets	10 ³ jets	10 ⁴ jets

Cu+Au ~ Au+Au/5

U+U (tip-tip) ~ Au+Au/500


Only stochastic cooling of Au beams assumed

Greater rate and p_T reach than singles



Huge rates allow differential measurements with geometry
(v_2 , v_3 , A+B, U+U, ...)
precise control measurements (d+Au & p+p).
Over 80% as dijets into $|\eta| < 1$

Simulating 200 GeV Au+Au with HIJING

Underlying event distribution



Underlying event distribution



Clear η dependence for larger acceptance motivated ATLAS/CMS-like method

HIJING seems to get overall multiplicities correct for RHIC energies

Large background/fluctuations motivate careful study of subtractions and fake rates


Iterative jet finding algorithm

Run jet reco (anti- k_T)
algorithm on 0.1×0.1
calorimeter cells

inspired by ATLAS

J.A. Hanks et al., Phys. Rev. C 86, 024908 (2012), arXiv:1203:1353


Iterative jet finding algorithm




inspired by ATLAS

J.A. Hanks et al., Phys. Rev. C 86, 024908 (2012), arXiv:1203:1353


Stepping through the subtraction procedure




Stepping through the subtraction procedure




Stepping through the subtraction procedure




Stepping through the subtraction procedure




Stepping through the subtraction procedure




Stepping through the subtraction procedure




Stepping through the subtraction procedure




Stepping through the subtraction procedure




Stepping through the subtraction procedure



Stepping through the subtraction procedure




Fake Jets at RHIC (R=0.2)




arXiv:1203:1353

Fake Jets at RHIC (R=0.2)



not matched jets:
no nearby HIJING jets
“fakes”


Fake Jets at RHIC (R=0.2)



**not matched jets:
no nearby HIJING jets
“fakes”**

**matched jets:
within $\Delta R < 0.25$ of a
HIJING truth jet ($>5\text{GeV}$)**

Fake Jets at RHIC (R=0.2)




**not matched jets:
no nearby HIJING jets
“fakes”**

**matched jets:
within $\Delta R < 0.25$ of a
HIJING truth jet ($> 5\text{GeV}$)**


**HIJING true jets:
reconstruct jets directly
from simulated
fragmentation**

Fake Jets at RHIC (R=0.2)




arXiv:1203:1353

Fake Jets at RHIC (R=0.2)





arXiv:1203:1353

Fake Jets at RHIC (R=0.2)



arXiv:1203:1353



Fake Jets at RHIC vs R



How well are we doing for real jets?


Embed PYTHIA jets in HIJING background

p+p shows effect of discretizing PYTHIA






How well are we doing for real jets?

Embed PYTHIA jets in HIJING background
 p+p shows effect of discretizing PYTHIA




Including detector effects



Parameterize single particle resolution and apply to cells used to measure jets in HIJING

- (check that effective resolution is comparable to GEANT4 in $p+p$)


Including detector effects



to measure jets in HIJING

- (check that effective resolution is comparable to GEANT4 in p+p)

Unfolding detector effects




PYTHIA jets plus
underlying
central Au+Au event
plus detector smearing

use ROOUNFOLD Iterative
Bayes' method

recovers truth spectrum


jet R_{AA} to high p_T possible

Sensitivity to inclusive jet-quenching





~10% systematic
from unfolding

Sensitivity to inclusive jet-quenching



~10% systematic
from unfolding



Sensitivity to di-jet asymmetry



Uncertainties shown correspond to statistics available from ~1yr (20 wks)
 - embedded in 0-10% HIJING

Approximate unfolding:
 - apply $\langle \Delta E_T \rangle$ shift to leading jet
 \approx recover underlying asymmetry
 \Rightarrow remain sensitive to modification!






Sensitivity to γ -jet asymmetry




Uncertainties shown correspond to statistics available from ~ 1 yr (20 wks)
 - embedded in 0-10% HIJING

Full unfolding with ROOUNFOLD
 \approx recover underlying asymmetry
 \Rightarrow more sensitive to modification!

Recovering 'lost' energy



A few key future additions





There is a future for jets at RHIC!



sPHENIX plan well suited for a variety of jet measurements!

Backup Slides

Direct photons at RHIC



- γ/π^0 very large at RHIC
- good S/B > 20 GeV
- substantial rate even > 30 GeV
- RHIC a very good place for γ -jet correlations

Identifying truth jets

deep within the HIJING Event Generation...

Identifying truth jets

deep within the HIJING Event Generation...

parton 1

Identifying truth jets

deep within the HIJING Event Generation...

parton 1



hadrons A

Identifying truth jets

deep within the HIJING Event Generation...

parton 1



hadrons A



FastJet

anti- k_T

Identifying truth jets

deep within the HIJING Event Generation...

parton 1



hadrons A



**FastJet
anti- k_T**



**jet A(R=0.2)
jet A(R=0.3)
jet A(R=0.4)**

Identifying truth jets

deep within the HIJING Event Generation...

parton 1



hadrons A



FastJet
anti- k_T



jet A(R=0.2)
jet A(R=0.3)
jet A(R=0.4)

parton 2



hadrons B



FastJet
anti- k_T



jet B(R=0.2)
jet B(R=0.3)
jet B(R=0.4)

Identifying truth jets

deep within the HIJING Event Generation...

parton 1



hadrons A



**FastJet
anti- k_T**



jet A(R=0.2)
jet A(R=0.3)
jet A(R=0.4)

parton 2



hadrons B



**FastJet
anti- k_T**



jet B(R=0.2)
jet B(R=0.3)
jet B(R=0.4)

parton 3



hadrons C




**FastJet
anti- k_T**



jet C(R=0.2)
jet C(R=0.3)
jet C(R=0.4)


Identifying truth jets

deep within the HIJING Event Generation...




Well reconstructed jets

True HIJING



Reconstructed




- $b = 1.8\text{fm}$ HIJING event, $R=0.2$ anti- k_T dijet
- well reconstructed above background fluctuations after subtraction


Fake jets

- $b=2.4$ HIJING event, no true jets
- Find 30 & 10 GeV fake jets with $R=0.4$ anti- k_T



True HIJING



Reconstructed



Jets at RHIC, really?



- smearing from very, very low p_T
- jets are not independent fluc's over full solid angle
- w/calorimetry what matters is no. of jet "patches"
- R can be smaller than 0.4 (say, 0.2)
 - agree that $R = 0.4$ at 20 GeV/c is B.G.
- energy in a cone doesn't look like a jet
- jets from soft fluctuations \Rightarrow modified FF's
- internal CMS study: jets have a high p_T hadron
- ATLAS is going to ~ 40 GeV/c with fake rejector

Jörn Putschke (HP'12, RHIC/AGS '11); also Cacciari, Salaam, Soyez, Eur.Phys.J.C71:1692,2011