Contribution ID: 37 Type: Submitted

Radiative strength of neutron-rich nuclei and its astrophysical implications

Wednesday 19 December 2012 12:30 (10 minutes)

F. Giacoppo 1, L. Bernstein 2, D. Bleuel 2, P.A. Butler 3, T. K. Eriksen 1, A. Görgen 1, M. Guttormsen 1, T. W. Hagen 1, A.C. Larsen 1, H. T. Nyhus 1, T. Renstrøm 1, S. Rose 1, S. Siem 1, G.M. Tveten 1, A. Voinov 4, M. Wiedeking 5, and J. Wilson 6

1Department of Physics, University of Oslo, N-0316 Oslo, Norway 2Lawrence Livermore National Laboratory, USA 3Oliver Lodge Laboratory, University of Liverpool, United Kingdom 4Department of Physics and Astronomy, Ohio University, USA 5 iThemba LABS, South Africa 6 IPN Orsay, CNRS-IN2P3, Orsay, France

The density and width of nuclear excited states increase with excitation energy towards the particle separation. The NLD and the RSF are fundamental input parameters for calculating nuclear reaction cross-sections and react. The Oslo nuclear physics group has developed a method to determine simultaneously the NLD and the RSF from part. Recently an unexpected low-energy increase in the radiative strength function of light and medium-mass nuclei, An experimental program to study the gamma-ray strength function of neutron reach nuclei has been recently propries.

- [1] A. Schiller et al., Nucl. Instrum. Methods Phys. A 447, 498 (2000).
- [2] A. C. Larsen et al., Phys. Rev. C 83, 034 315 (2011).
- [3] A.Voinov et al., Phys. Rev. Lett. 93, 142504 (2004).
- [4] M. Guttormsen et al., Phys. Rev. C 71, 044307 (2005);
- [5] M. Wiedeking et al., Phys. Rev. Lett. 108, 162503 (2012).
- [6] A. C. Larsen and S. Goriely, Phys. Rev. C 82, 014318 (2010).
- [7] S. Siem et al. "Statistical properties of warm nuclei: Investigating the low-energy enhancement in the gamma strength function of neutron rich nuclei". Submitted proposal to INTC –October 2012

Author: GIACOPPO, Francesca (University of Oslo)

Presenter: GIACOPPO, Francesca (University of Oslo)

Session Classification: New techniques