



Bose Institute


# **Resonance excitation in ${}^7\text{Be} + \text{d}$ reaction to study ${}^7\text{Li}$ abundance anomaly**

***Dhruba Gupta***

**Department of Physics, Bose Institute, Kolkata**

ISOLDE Workshop and Users Meeting, CERN, December 17-19, 2012

# Physics case



*Observed values represented by bands,  
predicted values represented by lines.*

$\eta_b^{WMAP} = n_B/n_\gamma = (6.23 \pm 0.17) \times 10^{-10}$   
ratio of the baryon and photon number densities

**Serious discrepancy** of a factor of about 4 in primordial  ${}^7\text{Li}$  abundance, while good agreement of D,  ${}^3,4\text{He}$  abundances

**BBN theory using  $\eta_b^{WMAP}$ :**  ${}^7\text{Li}/\text{H} = 5.12_{-0.62}^{+0.71} \times 10^{-10}$

**Observationally extracted:**  ${}^7\text{Li}/\text{H} = 1.23_{-0.16}^{+0.34} \times 10^{-10}$

## The Cosmological ${}^7\text{Li}$ problem

### Aim of the experiment:


Study discrepancy of  ${}^7\text{Li}$  abundance in the context of **resonance enhancement** of nuclear reactions

### Importance:

One of the **important unresolved problems** of present-day astrophysics.

**Existing data:** nuclear/astrophysical/new effects beyond standard BBN- inconclusive


# Incomplete nuclear physics input for BBN calculations: Can resonant enhancement alleviate this discrepancy?



${}^7\text{Li}$  mostly produced as  ${}^7\text{Be}$  during Big Bang. One narrow nuclear level in  ${}^9\text{B}$ ,  $E_{5/2^+} \approx 16.7$  MeV, not sufficiently studied experimentally, and just  $\sim 200$  keV above the  ${}^7\text{Be}+d$  threshold, may lead to resonant enhancement of  ${}^7\text{Be}(d,\gamma){}^9\text{B}$  and  ${}^7\text{Be}(d,p){}^8\text{Be}$  reactions.

*Cyburt et al*

arXiv:0906.4373v1 (2009)



${}^7\text{Be}(d,p){}^8\text{Be}^* \rightarrow 2\alpha$  ( $Q = 16.490$  MeV)


*R. W. Kavanagh*

*Nuclear Physics 18 (1960) 492*


**upto  $E_x = 11$  MeV**

$E_{\text{cm}} = 0.6 - 1.3$  MeV, reaction rate relied on an extrapolation to BBN energies. Differential cross section multiplied by  $4\pi$  (assuming isotropic angular distribution) and **arbitrarily** by 3 (to estimate contribution of higher energy  ${}^8\text{Be}$  states)

Angulo et al  
Astrophys. Jour. 630 (2005) L105



O'Malley et al  
Phys. Rev. C 84, 042801( R) (2011)



${}^7\text{Be}(d,p){}^8\text{Be}^* \rightarrow 2\alpha$  ( $Q = 16.490$  MeV)

( $E_{7\text{Be}} = 5.55, 1.71$  MeV) **upto  $E_x = 13.8$  MeV**

- Kavanagh (1960)
- Angulo (2005),  ${}^8\text{Be}^*$  (g.s + 1<sup>st</sup> ex.s)

*Cross section overestimated previously*

Due to Coulomb barrier, contributions of the higher states were negligible. Small angular range covered ( $\sim 7$ - $17$  deg) and full isotropy for proton angular distribution **assumed** in calculating average cross section

Resonance enhancement of the  ${}^7\text{Be}(d,p)2\alpha$  reaction through the  $5/2^+$  16.7 MeV resonance state in  ${}^9\text{B}$  ?  ${}^2\text{H}({}^7\text{Be},d){}^7\text{Be}$  ( $E_{7\text{Be}} = 10$  MeV)


**No evidence for a resonance** observed

*Scholl et al Phys. Rev. C 84, 014308 (2011)*

${}^9\text{Be}({}^3\text{He},t){}^9\text{B}$ ,  $E = 140$  MeV/A, the state is strongly excited.

Energy: 16.800(10) MeV, width: 81(5) keV

Cybert et al  
arXiv:0906.4373v1 (2009)



Reaction rates of the  ${}^7\text{Be}$  destruction by deuterons could be large, owing to a narrow resonance 16.7 MeV ( $5/2^+$ ) in  ${}^9\text{B}$ . **This resonance may be very strong**, and at the very limit of the quantum mechanically allowed value for the deuteron separation width. This would be responsible for a factor of  $\sim 2$  suppression of the primordial  ${}^7\text{Be}$  yield, **resolving  ${}^7\text{Li}$  problem**.

(Resonant energy  $E_r$ , deuteron separation width  $\Gamma_d$ )  $\approx$  **(170-220, 10-40) keV** can **eliminate current discrepancy**. Such a large width at this resonant energy can only be achieved if the interaction radius for the deuteron entrance channel is very large,  $a_{27} \geq 9$  fm

Chakraborty et al  
Phys. Rev. D 83, 063006 (2011)

16.7 MeV ( $5/2^+$ ) in  ${}^9\text{B}$ , a large channel radius ( $a > 10$  fm) needed to give sufficiently large widths

**Clearly more study of possible resonances in  ${}^7\text{Be} + d$  reactions and the 16.7 MeV state in  ${}^9\text{B}$  needed. This state has the potential to significantly influence  ${}^7\text{Be}$  destruction.**

# Merit of the experiment

*Measurements of the 16.7 MeV (5/2+) resonance in  $^9\text{B}$  at HIE-ISOLDE offering higher beam energies and intensity of  $^7\text{Be}$  can resolve the  $^7\text{Li}$  issue. Direct experimental information about its  $p$  and  $\alpha$  decay properties unknown. Dominant single proton-decay branch to the 16.626 MeV ( $2^+$ ) state in  $^8\text{Be}$  ?*

## FIRST TIME

*higher beam energy* : measure higher excitation energies in  $^8\text{Be}$  up to about **20 MeV**

*wider angular coverage* : 5-50 deg, improved average cross section measurements without assuming isotropy done in earlier works


## POSSIBLE AT HIE-ISOLDE (beam energy and intensity)

**35 MeV  $^7\text{Be}$**  : measure (d,p) (d,d) with T-REX

$\text{CD}_2$  target of thickness 1 mg/cm<sup>2</sup>,  $I = 10^8$  pps



Estimated cross section  $\sim 500$  mb

Expected count rate  $\sim 1000$ /s





# Kinematics





*Kinematics of the  ${}^7\text{Be}(d,d)$ ,  ${}^7\text{Be}(d,p)$  reactions for ground state ( $d$ ) ( $p$ ), excitation of 20 MeV ( $p'$ ), 5 MeV ( $d'$ ) respectively. The forward hashed (backward hashed) areas cover the detected  $p$  ( $d$ ) energies and angles in CD and barrel detectors of T-REX.*


CD ( $\Delta E=500 \mu\text{m}$ ,  $E = 1.5 \text{ mm}$ ), Forward Barrel ( $\Delta E=140 \mu\text{m}$ ,  $E = 1.0 \text{ mm}$ )

**Protons** :  $E_x = 0$ ,  $E_p = 20\text{-}40 \text{ MeV}$ , do not stop

$E_x = 20 \text{ MeV}$ ,  $E_p = 4\text{-}14 \text{ MeV}$  at  $5 - 50 \text{ deg}$ , corresponding  $\alpha$ -particles stop in  $\Delta E$



*Bildstein et al*  
*Eur. Phys. J. A 48, 85 (2012)*



*Energy resolution of T-REX would be sufficient for particle identification in this experiment*

Excitation energies of the populated states of interest can be reconstructed from deposited energies in various detectors for stopping particles. Breakup of  ${}^8\text{Be} \rightarrow 2\alpha$  particles that stop in  $\Delta E$  detectors of T-REX, can be identified by using a multiplicity trigger.



At ISOLDE,  ${}^7\text{Be}$  is produced with very high yields and even with a modest charge breeding efficiency 2-3% we should get plenty of beam at the end of the linac. Energy should be well above 5.5 MeV/A for this A/q.

Concerning contaminations of the beam, both  ${}^{14}\text{N}$  and  ${}^{21}\text{Ne}$  can be **suppressed completely** by selecting  ${}^7\text{Be}^{4+}$  after a stripping foil inserted at the end of the linac ( ${}^7\text{Be}^{3+}$  being accelerated up to the stripper foil).

As the halflife is long we will not reach saturation for the radiation level during the run. There are no restriction from a radiation protection point of view. If (for example) we take  $10^8$  pps for 3 x 8 h shifts we will end up with  $\sim 9 \times 10^{12}$  ions collected which lead to an activity of 0.2 MBq. Regarding the radiological hazards of radioactive nuclei produced at ISOLDE, the activity we need for our experiment is only a very small fraction of the authorization limit LA (**very small internal exposure risk**).

The hazard related to **external exposure**, assessed via the h10 quantity which for  ${}^7\text{Be} = 0.008$  (mSv/h)/GBq at 1 m quantity which give the dose rate at 1 m per GBq of activity.

# Outcome of the Experiment

- **Higher excitation energies in  $^8\text{Be}$**  - measurement up to about **20 MeV** from (d,p)
- **Properties of the 16.7 MeV ( $5/2^+$ ) resonance in  $^9\text{B}$**  from (d,d)

Direct experimental determination of corresponding reaction rates might either **support/refute** a nuclear physics solution to the lithium problem. Other effects may include astrophysical effects, new effects (beyond standard BBN model) or a combination of all. Conventional solutions to the lithium problem ask for experimental testing.

While **astrophysical solutions** are not ruled out, they **are increasingly constrained**. Thus a serious and thorough evaluation of **all possible nuclear physics aspects** of primordial lithium production is urgent in order to determine whether the lithium problem truly points to new fundamental physics.

- **Possible at HIE-ISOLDE.**

EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH  
Proposal to the ISOLDE and Neutron Time-of-Flight Committee

Search for higher excited states of  $^8\text{Be}^*$  to study the  
cosmological  $^7\text{Li}$  problem

October 2, 2012

D. Gupta<sup>1</sup>, S. K. Saha<sup>1</sup>, Maria JG Borge<sup>2</sup>, M. Kowalska<sup>2</sup>, D. Muecher<sup>3</sup>, R. Raabe<sup>4</sup>,  
G. Randisi<sup>4</sup>, P. Reiter<sup>3</sup>, T. Stora<sup>2</sup>, D. Voulot<sup>2</sup>, F. Wenander<sup>2</sup>

<sup>1</sup>Department of Physics, Bose Institute, 93/1 A. P. C. Road, Kolkata 700009, India

<sup>2</sup>CERN, CH-1211 Geneva 23, Switzerland

<sup>3</sup>Institut fuer Kernphysik, Universitaet zu Koeln, Zulpicher Str. 77, D-50937 Koeln, Germany

<sup>4</sup>Instituut voor Kern- en Stralingsfysica, KU Leuven, Celestijnenlaan 200D, B-3001 Leuven, Belgium

**Spokesperson:** [Dr. Dhruba Gupta] [dhruba@bosemain.boseinst.ac.in]  
**Contact person:** [Dr. Magdalena Kowalska] [Magdalena.Kowalska@cern.ch]

**Abstract:** We would like to study the unresolved  $^7\text{Li}$  abundance anomaly by carrying out experiments that destroy the rare isotope  $^7\text{Be}$ , the main source of  $^7\text{Li}$ . Utilizing a 35 MeV  $^7\text{Be}$  beam from HIE-ISOLDE, we would like to measure the (d,p) and (d,d) reactions with T-REX. The higher beam energy, for the first time, would allow us to measure higher excitation energies in  $^8\text{Be}$  up to about 20 MeV. With a wider angular coverage, we can make improved average cross-section measurement without assuming isotropy done in earlier works.

CERN-INTC-2012-049 / INTC-P-350  
02/10/2012

Requested shifts: [15] shifts, (split into [1] runs over [1] years)

**Collaborators:**

***S. K. Saha (Bose Institute)***

***Maria JG Borge, M. Kowalska, T. Stora, D. Voulot, F. Wenander (CERN)***

***D. Muecher (Technischen Universität Munchen, Munchen)***

***R. Raabe, G. Randisi (IKS, KU Leuven)***

***P. Reiter (IfK, Koeln)***

**Thank You**