

ISOLDE Workshop and Users meeting 2007/2008

Monday, 17 December 2007 - Wednesday, 19 December 2007

CERN

Book of Abstracts

Contents

Sub-Barrier Coulomb Excitation of ^{110}Sn and First Results on $^{108,106}\text{Sn}$	1
Shape coexistence in n-rich Strontium isotope at REX+ Miniball	1
IS456 - Ground and isomer state properties of the neutron-deficient polonium isotopes using the ISOLDE RILIS	2
Coulomb excitation on the $^{184,186,188}\text{Hg}$ nuclei	2
ISOLDE news	3
Spin determination and nuclear moment measurements of ^{71}Cu and ^{72}Cu	3
Hydrodynamics of ISOLDE liquid metal targets	4
Selective contaminant adsorption for RIB purification at ISOLDE	5
The $d(^{30}\text{Mg}, ^{31}\text{Mg})p$ transfer reaction at REX-ISOLDE: Results from IS454	5
Breakup of the 18.2 MeV state in ^{11}Be : New decay modes.	6
Towards High Intensity Energy ISOLDE: extrapolation of yields for accelerated isotopes at REX-ISOLDE	6
Transfer reactions with HIE-ISOLDE	7
First proof of shape coexistence at the borderline of the 'Island of Inversion': Identification of the second 0^+ state in ^{30}Mg and measurement of its electric monopole strength	7
Lattice location of implanted ions and characterization of implantation-induced damage in Ge	8
FEBIAD ion source operation modes to tune its selectivity: physics processes, numerical simulations and experimental data.	9
ECR charge breeding at ISOLDE	9
Local Probe Studies on Highly Distorted Rare-Earth Manganites	10
First results from electron emission channeling on-line experiments	10
Nuclear orientation in transfer reactions	11
WITCH: first measurement and recent developments	11

Rare isotope beams at MSU - present and future	12
20 Years of ISOLTRAP	12
Mössbauer studies of dilute magnetic semiconductors (IS-443)	12
Results with ISCOOL a new Radio Frequency Quadrupole Cooler and Buncher at ISOLDE	13
News from the RIKEN RI beam factory	14
Bremsstrahlung accompanied α -decay ^{210}Po	14
Nuclear mass measurements for nucleosynthesis studies at ISOLTRAP	15
Status report on g-factor measurements on Coulomb excited ^{138}Xe nuclei using the Transient-Field technique (IS415)	15
REX-ISOLDE - now also heavyweight champion	15
Beta-asymmetry measurements as a probe for non standard model physics	16
First laser measurements on radioactive beams using ISCOOL	16
Studies of the ^{12}C nucleus using beta-decay of ^{12}N and ^{12}B	17
JYFLTRAP at IGISOL - Atomic masses from nuclear structure physics to fundamental physics	18
Coulomb excitation of neutron-rich ^{44}Ar at SPIRAL	18
TRIGA-TRAP: Mass measurements on exotic and heavy nuclides	19
Shell structure and shape changes in neutron rich krypton isotopes	19
Magneto-resistive materials: hyperfine studies using radioactive isotopes.	20
Hyperdeformation in Xe Nuclei.	20
Between Bose-Einstein condensate and fermionic superfluid.	21
Double-beta decay and neutrino physics	21
Coulomb excitation of ^{31}Mg - at the shore of the island of inversion	22
Inelastic Branch of the Stellar Reaction $^{14}\text{O}(\alpha,p)^{17}\text{F}$	22
Future physics at HIE-ISOLDE: a personal view	23
Gamma-ray nucleosynthesis: results from the INTEGRAL satellite	23
Masses of halo nuclides and decay-spectroscopy in the lead region: new developments at ISOLTRAP	23
The first 40 years of physics at ISOLDE	24
The ISAC facility at TRIUMF and Halo-nuclei mass measurements at TITAN	24
MAYA active target for the study of extremely exotic nuclei	25

Shell structure evolution in nuclei : future collaboration between GANIL and	25
The HIE-ISOLDE project	25
Soft-landing of Radioactive Probes on Clean Metal Surfaces and at Interfaces	25
Target and Ion Source Development	26
Pb(II) and Hg(II) binding to proteins and model systems studied by ^{204}mPb - and $^{199\text{mHg}}$ - Perturbed Angular Correlation of γ -rays (PAC) spectroscopy	26
Welcome	26

Nuclear Physics III / 0

Sub-Barrier Coulomb Excitation of ^{110}Sn and First Results on $^{108,106}\text{Sn}$ **Author:** Andreas Jan Ekstrom¹**Co-authors:** Claes Fahlander¹; Joakim Cederkall²; Morten Hjorth-Jensen³; al et⁴¹ *Physics Department, University of Lund, Sweden*² *PH Department, CERN 1211, Geneva 23, Switzerland.*³ *Physics Department and Center of Mathematics for Applications, University of Oslo, Norway*⁴ .

The advent of radioactive ion beams (RIBs) places exotic nuclei within reach of experimental study. In particular, RIBs enables a new investigation of the low energy structure of isotopes in the vicinity of the doubly magic ^{100}Sn nucleus. We present here the first results from a series of measurements of the reduced transition probability - the $B(E2)$ value - for the neutron deficient Sn-isotopes $^{110,108,106}\text{Sn}$ [1]. All three experiments were performed using sub-barrier Coulomb excitation at 2.8 MeV/u at REX-ISOLDE with the MINIBALL germanium detector array. This technique provides a measurement accuracy of ~10%. The $B(E2)$ value can presently difficult to obtain using any other method, due to a high lying $6+$ isomer present in the neutron-deficient even Sn isotopes.

Doubly-magic nuclei are of significant theoretical importance, since they provide a good testing ground for shell-model calculations. The Sn isotopic chain comprises nuclei between neutron numbers $N=82$ and $N=50$. Therefore it provides a unique opportunity to study the shell structure evolution as a function of the neutron degree of freedom. The spectroscopy of the low lying states in the even Sn isotopes have been explained within the generalized seniority scheme [2]. Large scale shell model calculations using a model space confining the neutrons to the *gdsh* orbitals support this picture [3]. Our recent experiments together with results from refs. [3,4] indicate a deviation from theoretical predictions manifested in a stronger than expected collectivity towards the proton dripline. This might imply that further core-polarization effects and a refined effective interaction are needed. One perhaps important effect recently discussed in refs. [5,6,7] is that of an enhanced attractive neutron-proton interaction in spin- isospin-flip vertices contained in the monopole part of the tensor force. Hence, with a decrease of neutrons in the orbital $0g_{7/2}$, the proton $0g_{9/2}$ becomes less bound, implying an increased core-excitation probability. Thus, this effect could compete with a reduction of the $B(E2)$ values originating in the decrease in neutron number. We will present the latest results and data analysis from $^{110,108,106}\text{Sn}$, and where they stand in comparison with present theoretical models.

[1] J. Cederkall et al., Phys. Rev. Lett. 98, 172501 (2007).

[2] I. Talmi, Nucl. Phys. bf A172, 1 (1971).

[3] A. Banu et al., Phys. Rev. C 72, 061305(R) (2005).

[4] C. Vaman et al., arXiv:nucl-ex/0612011v1.

[5] T. Otsuka et al., Phys. Rev. Lett. 87, 082502 (2001).

[6] T. Otsuka et al., Phys. Rev. Lett. 95, 232502 (2005).

[7] B. L. Cohen et al., Phys. Rev. 127, 597 (1962).

Nuclear Physics III / 1

Shape coexistence in n-rich Strontium isotope at REX+ Miniball**Author:** Clement Emmanuel¹**Co-author:** Andreas Goergen²¹ *European Organization for Nuclear Research (CERN)*

² CEA-Saclay

Corresponding Author: emmanuel.clement@cern.ch

The nuclei in the mass region $A=100$ around Sr and Zr show a dramatic change of the nuclear ground-state shape from near spherical for $N \leq 58$ to strongly deformed for $N > 60$. Theoretical calculations predict the coexistence of slightly oblate and strongly prolate deformed configurations in the transitional region. However, excited rotational structures based on the highly deformed configuration, which becomes the ground state at $N = 60$, are not firmly established in the lighter isotopes, and the earlier interpretation of a very abrupt change of shape has been challenged by recent experimental results in favor of a rather gradual change. The low spins electromagnetic properties of the neutron-rich ^{96}Sr were studied by low-energy Coulomb excitation using the REX-ISOLDE facility and the MINIBALL detector array. The experiment was performed in June 2007 using a molecular extraction and for the first time at the new position of the Miniball array. Preliminary results will be presented.

Nuclear Physics II / 2

IS456 - Ground and isomer state properties of the neutron-deficient polonium isotopes using the ISOLDE RILIS

Author: Thomas E. Cocolios¹

Co-authors: Andrei N. Andreyev¹; Bruce A. Marsh²; Dima Fedorov³; Ernesto Mane Jr.⁴; Francois Le Blanc⁵; Gerhard Huber⁶; Iulian Stefan⁵; Jarno Van de Walle²; Jeroen Buescher¹; Mark Huyse¹; Maxim Seliverstov⁶; Pavel Molkov³; Piet Van Duppen¹; Serge Franchoo⁵; Sergei Zemlyanoy⁷; Stanislav Antalic⁸; Tomas Smets¹; Ulli Koester⁹; Ulli Kudryavtsev¹; Valentin Fedosseev²

¹ *Instituut voor Kern- en Stralingsfysica*

² *CERN*

³ *Petersburg Nuclear Physics Institute*

⁴ *University of Manchester*

⁵ *Institut de Physique Nucleaire d'Orsay*

⁶ *IP - Johannes Gutenberg Universitaet*

⁷ *Joint Institute of Nuclear Research*

⁸ *Comenius University*

⁹ *Institut Laue Langevin*

Corresponding Author: thomas.elias.cocolios@cern.ch

For many years, the RILIS has provided very clean beams for ISOLDE. It is now also a tool used for precision measurement of nuclear ground and isomer state properties. Combining the laser spectroscopy technique in the ion source itself with the high sensitivity and precision of nuclear spectroscopy, it is possible to extract optical spectra for very exotic nuclei as was demonstrated with the lead and bismuth studies at the RILIS. This last summer, a similar study was performed on the neutron-deficient $^{193-200,202,204}\text{Po}$ isotopes using recently developed laser ionized beams of polonium. In this presentation, we shall report on the achievements of experiment IS456 as well as on the challenges of the analysis of such data. The first insight on the nuclear properties will be discussed.

Nuclear Physics III / 3

Coulomb excitation on the $^{184,186,188}\text{Hg}$ nuclei

Authors: A. Petts¹; N. Bree²

Co-authors: A. Larsen ³; B. Butler ¹; D. Jenkins ⁴; D. Voulot ⁵; E. Clément ⁵; E. Piselli ⁵; F. Wenander ⁵; I. Stefanescu ²; J. Cederkäll ⁵; J. Pakarinen ¹; J. Van de Walle ⁵; K. Hadynska ⁶; L. Fraile ⁵; M. Guttormsen ³; M. Huyse ²; M. Scheck ⁷; M. Zielinska ⁸; N. Patronis ²; P. Marley ⁴; P. Napiorkowski ⁶; P. Peura ¹; P. Van Duppen ²; R. Herzberg ¹; R. Julin ⁹; R. Krücken ¹⁰; S. Knapen ²; S. Siem ³; T. Cocolios ²; T. Grahn ¹; Th. Kroell ¹⁰

¹ *Oliver Lodge Laboratory, University of Liverpool, United Kingdom*

² *Instituut voor Kern- en Stralingsfysica, K.U. Leuven, Celestijnenlaan 200D, B-3001 Leuven, Belgium*

³ *Department of Physics, University of Oslo, Blindern, N-0316 Oslo, Norway*

⁴ *Department of Physics, University of York, Heslington, York YO10 5DD, United Kingdom*

⁵ *ISOLDE, CERN, CH-1211 Geneva 23, Switzerland*

⁶ *Heavy Ion Laboratory, Warsaw University, Pasteura 5A, 02-093 Warsaw, Poland*

⁷ *Department of Physics and Astronomy, University of Kentucky, Lexington, Kentucky 40506-0055, USA*

⁸ *DAPNIA, CEA Saclay, F-91191 Gif-sur-Yvette Cedex, France*

⁹ *Department of Physics, University of Jyväskylä, SF-40100 Jyväskylä, Finland*

¹⁰ *Technische Universität München, James-Franck-Str., D-85748 Garching, Germany*

Corresponding Author: nick.bree@cern.ch

We report on a Coulomb excitation experiment at 'safe' energies on the even-even neutron-deficient ^{184,186,188}Hg isotopes at the REX-ISOLDE radioactive beam facility. For the first time a Coulomb excitation experiment using post-accelerated radioactive ion beams was performed in the heavy mass region. Apart from the first 2+ to the 0+ ground state transition, also Coulex towards the second 2+ and the 4+ state was observed and gamma-rays from the different decay paths were detected in the Miniball gamma-spectrometer. From the intensity of these gamma-rays the matrix elements connecting the different states in the different bands will be deduced for these nuclei. The obtained results should lead to a better understanding of the shape coexistence in this mass region of the nuclear chart.

HIE-ISOLDE / 4

ISOLDE news

Author: Karsten Riisager¹

¹ *CERN*

Corresponding Author: karsten.riisager@cern.ch

I shall review briefly recent (and foreseen) administrative changes at CERN that are relevant for the ISOLDE users, including registration of users and safety courses.

HIE-ISOLDE / 5

Spin determination and nuclear moment measurements of ⁷¹Cu and ⁷²Cu

Author: Kieran Flanagan¹

Co-authors: Bradley Cheal ²; Christopher Geppert ³; David Forest ⁴; Deyan Yordanov ⁵; Garry Tungate ⁴; Gerda Neyens ⁶; Henry Stroke ⁷; Joerg Kraemer ³; Jon Billowes ²; Klaus Blaum ⁵; Magda Kowalska ⁸; Marieke De Rydt ⁶; Paul Campbell ²; Peter Lievens ⁶; Pieter Vingerhoets ⁶; Rainer Neugart ⁵; Wilfried Nörtershäuser ⁵

¹ *KU-University of Leuven*

² *University of Manchester*

³ GSI⁴ University of Birmingham⁵ University of Mainz⁶ Katholieke University of Leuven⁷ New York University⁸ CERN**Corresponding Author:** kieran.flanagan@cern.ch

As part of the IS439 experiment the COLLAPS collaboration has recently measured the hyperfine structure of $^{71,72}\text{Cu}$. From this work we are able to unambiguously assign the spin of ^{72}Cu and extract the magnetic and quadrupole moments for both isotopes. This work is part of the ongoing investigation of the evolution of nuclear structure with neutron excess in the copper isotope chain. A central motivation of the original proposal was to measure the spin/parity of ^{72}Cu , and thus resolve the inconsistency between recent results from in-source spectroscopy and beta-decay studies [1,2]. This experiment uses high-resolution laser spectroscopy, which can unambiguously measure the nuclear spin and provide model-independent measurements of the nuclear moments. The interpretation of these recent measurements in conjunction with the hyperfine structure measurements from 2006 will be presented and discussed.

[1] J.-C. Thomas et al., Phys. Rev. C 74, 054309 (2006).

[2] U. Koster, Private Communication, 2005.

Technical Developments / 6

Hydrodynamics of ISOLDE liquid metal targets

Author: Etam Noah Messomo¹**Co-authors:** Jacques Lettry¹; Luca Bruno¹; Richard Catherall¹; Thierry Stora¹¹ CERN**Corresponding Author:** etam.noah@cern.ch

The proton beam from the CERN Proton-Synchrotron Booster (PSB) is highly pulsed. A typical 1.4 GeV PSB pulse consists of four bunches, with a maximum bunch power of 12 GW. This is more than six orders of magnitude higher than the often quoted average beam power of ~ 2.7 kW. The lifetimes of early liquid metal targets exposed to the PSB beam were severely compromised through broken welds, corroded proton beam windows and flow of liquid metal into the ion source. To reduce the energy densities for liquid metal target units, the current practice is to spatially defocus the beam and to apply a time-staggered extraction mode to the PSB bunches. Hydrodynamic simulations of proton-induced transient effects such as splashing were carried out and showed that the 10 ns bunch spacing which matches the target relaxation time leads to constructive interference of proton-induced pressure waves. Measurements of the release of ^{84}Kr from a Pb target and ^{111}Cd from a Sn target confirm the dependence of release on bunch spacing. Small changes in the PSB bunch spacing can significantly reduce pressure amplitudes and may be a solution to further improve ISOLDE liquid metal target release and yield properties by increasing the practical limit of 8×10^{12} protons per pulse whilst ensuring target lifetimes remain unaffected. With the ongoing design phase and planned construction of the CERN accelerator complex upgrade, this study introduces some of the engineering tools which will be required in the development of the new target and ion source units compatible with the proton beam characteristics.

Summary:

Acknowledgements: We acknowledge the financial support of the EC under FP6 "Research Infrastructure Action – Structuring the European Research Area" EURISOL DS Project; Contract No. 515768 RIDS. The EC is not liable for any use that can be made on the information contained herein.

Technical Developments / 7**Selective contaminant adsorption for RIB purification at ISOLDE****Author:** Elian Bouquerel¹**Co-authors:** Jacques Lettry²; Martin Eller²; Richard Catherall²; Stefano Marzari²; Thierry Stora²¹ CERN, Université Paris 11, Orsay.² CERN**Corresponding Author:** elian.bouquerel@cern.ch

Target and ion source units have been recently developed at CERN-ISOLDE with the aim of delivering isobarically clean beams of short lived neutron-rich Cd and Zn. Chemical selectivity is achieved by specific interaction of the contaminant with a catching material inserted in the transfer line [1, 2]: a quartz insert allowed the retention and the decay of alkali contaminants by controlling its temperature. The temperature dependence showed to originate from a longer effusion time, and higher decay, of the impurities by selective adsorption onto the quartz. The design of this transfer lines is presented. Alkali suppression factors are deduced from 80Rb, 8Li, 46K and 126,142Cs yield measurements. The enthalpy of adsorption for Rb and Cs is calculated from 80Rb and 126Cs suppression factors.

[1] E. Bouquerel et al., 'Purification of a Zn Radioactive Ion Beam by alkali suppression in a quartz line target prototype', European Phys. Journal A (in press).

[2] – K.L. Kratz et al., 'The Beta-decay half-life of 13048Cd82 and its importance for astrophysical r-process scenarios', Z. Phys. A325 (1986) 489.

Nuclear Physics I / 8**The d(³⁰Mg, ³¹Mg)p transfer reaction at REX-ISOLDE: Results from IS454****Author:** Vinzenz Bildstein¹**Co-authors:** Jan Diriken²; Kathrin Wimmer¹; Mark Huyse²; Nick Bree²; Nikolas Patronis²; Piet Van Dupper²; Reiner Krücken¹; Riccardo Raabe²; Roman Gernhäuser¹; Thorsten Kröll¹¹ Physik-Department E12, TU München² Instituut voor kern- en stralingsfysica, KU Leuven**Corresponding Author:** vinzenz.bildstein@cern.ch

Thirty years after the discovery of the "island of inversion" [1] the borders of the island are still not well determined and in particular the evolution of the single-particle structure is not well investigated.

Transfer reactions yield important spectroscopic information, i.e. spin and parity assignments as well as spectroscopic factors, complementary to the information obtained in Coulomb excitation [2].

In order to study transfer reactions in inverse kinematics at REX-ISOLDE with MINIBALL a new setup was built covering a large solid angle. This new setup overcomes the limitations of previous transfer experiments performed at REX-ISOLDE [3].

In the first experiment the nucleus ³¹Mg which is right on the edge of the "island of inversion" was studied via the d(³⁰Mg, ³¹Mg)p reaction.

First results of this beamtime which took place at the end of October this year will be shown as well as future plans for transfer experiments at REX-ISOLDE.

[1] C. Thibault et al., Phys. Rev. C 12, 644 (1975)

[2] O. Niedermaier et al., Phys. Rev. Lett. 94, 172501 (2005)

[3] M. Pantea, PhD Thesis, TU Darmstadt, Germany (2005)

*supported by BMBF 06MT238, DFG Cluster of Excellence "Origin of the Universe" and the EU through RII3-EURONS (contract no. 506065).

Nuclear Physics II / 9

Breakup of the 18.2 MeV state in ^{11}Be : New decay modes.

Authors: Hans Fynbo¹; Karsten Riisager²; Maria Jose Garcia Borge³; Miguel Madurga Flores³; Olof Tengblad³

Co-authors: Bjorn Jonson⁴; Goran Nyman⁴

¹ *Physics and Astronomy (Univ. of Aarhus)*

² *ISOLDE*

³ *Consejo Superior de Investigaciones Cientificas (CSIC)*

⁴ *Fundamental Physics (Chalmers Univ.)*

Corresponding Author: miguel.madurga@cern.ch

The ^{11}Li β -decay offers a unique window to understand the nuclear structure far beyond the valley of stability. The β -delayed charged particle emission of ^{11}Li has been the subject of several previous studies [1-4]. The established channels involve the emission α particles ($2\alpha+3n$), ^6He ($^6\text{He}+\alpha+n$), tritons ($^8\text{Li}+t$), deuterons ($^9\text{Li}+d$) and the emission of $1n$ and $2n$ feeding the ground states of ^{10}Be and ^9Be respectively. The two channels involving α particles were previously studied in a ^{11}Li β -decay coincidence experiment by Langevin et al.[4]. The coincidence charged particle spectrum was explained as due to the breakup of two states in ^{11}Be at 10.6 and 18.2 MeV excitation energy. The breakup of the latter was assumed to occur by the three body channel $n\alpha^6\text{He}$ and the 5-body $3n2\alpha$. However, a recent ^{11}Li β -decay experiment performed by our collaboration [7] observed structures in the coincidence scatter plot interpreted as the sequential break-up of ^{11}Be through intermediate ^4He resonances.

In this work we present the first results of a new ^{11}Li β -decay experiment we performed at ISOLDE to clarify our previous interpretation [7]. The experimental set-up, consisting of 3 DSSSD's, was optimized for maximum solid angle coverage while having good spatial resolution. This improvements allowed us to record five times as many statistics as in the previous experiment, and to obtain direct evidence of sequential three-body, $n\alpha^6\text{He}$, break-up of ^{11}Be through the ground state of ^7He . Moreover, by reconstructing the neutron energy using energy and momentum conservation, we have identified two new states in ^{11}Be decaying through this new ^7He channel at 15.25, 16.18 MeV on top a previously known state at 18.0 MeV.

1] M. Langevin et al., Phys Lett. B146 (1984) 176.

[2] M. Langevin et al., Nucl. Phys. A366 (1981) 449.

[3] I. Mukha et al., Phys. Lett. B367 (1996) 65.

[4] M.J.G. Borge et al., Nucl. Phys. A613 (1997) 199.

[5] M. Madurga et al., in preparation

Technical Developments / 10

Towards High Intensity Energy ISOLDE: extrapolation of yields for accelerated isotopes at REX-ISOLDE

Author: Anna Gustafsson¹

¹ *Subatomic Physics Group - Department of Physics - Chalmers University of Technology*

Since the first year of operation, REX-ISOLDE has been delivering numerous post-accelerated radioactive beams [1,2]. An update of the ISOLDE database 3 is being done, including REX efficiencies for more than 50 isotopes corresponding to more than 15 elements. Based on this data the yields of post-accelerated beams was extrapolated for the upgrade of the ISOLDE facility, the so-called High Intensity Energy ISOLDE project.

A method will be demonstrated to predict yields for post accelerated isotopes at HIE-ISOLDE starting from the existing low energy ISOLDE yield database. The REX post acceleration system has the potential to transport and accelerate most of the 700+ radioisotopes from 65 elements of ISOLDE's repertoire thanks to the generic approach used for charge breeding and acceleration.

[1]: J. Cederkall et al., *nucl. Phys. A746* (2004) 17c-21c

[2]: D. Voulot et. al. *Nucl. Instrum. and Meth. A*, proceeding of the EMIS 2007 conference, to be published

HIE-ISOLDE / 11

Transfer reactions with HIE-ISOLDE

Author: Riccardo Raabe¹

¹ *Instituut voor Kern- en Stralingsfysica, K.U.Leuven*

The evolution of the shell structure far from stability shows striking effects, due to changes in the underlying interaction. Spin-orbit and proton-neutron interactions, pairing, the vicinity of the continuum, induce effects that translate in observables such as the migration of single-particle levels, reduced or disappearing shell gaps, deformation. Precise spectroscopic information has been so far mainly provided by decay studies; with the development of intense and pure post-accelerated beams of unstable ions, however, the use of reaction methods has also become possible.

Transfer reactions, thanks to their selectivity, are a particularly valuable tool, as they can give access to the most complete information about the structure of the populated levels. They have been used extensively for the investigation of the structure of stable nuclei; with radioactive nuclei, the experimental conditions set severe requirements on the quality of the ion beams and the detection setup.

HIE ISOLDE will provide an important opportunity to apply transfer reaction methods in regions of the chart of nuclei which are not accessible at other facilities. One-nucleon transfers can be used to study the single-particle states in the vicinity of a closed shell nucleus, or their evolution along a series of isotopes between closed shells. We will present some concrete examples for the N=16, 20, 28 and Z=28 regions, discussing the feasibility of measurements utilizing the present detection arrays and possible upgrades.

Nuclear Physics I / 12

First proof of shape coexistence at the borderline of the 'Island of Inversion': Identification of the second 0+ state in ³⁰Mg and measurement of its electric monopole strength

Author: Wolfgang Schwerdtfeger¹

Co-authors: Dietrich Habs ¹; Henryk Mach ²; Herbert Huebel ³; Kathrin Wimmer ¹; Kris Heyde ⁴; Luis Fraile ⁵; Per Hoff ⁶; Peter Ring ⁷; Peter Thirolf ¹; Ralf Hertenberg ¹; Reiner Kruecken ⁷; Roman Gernhaeuser ⁷; Rudolph Lutter ¹; Thomas Morgan ¹; Thorsten Kroell ⁷; Ulrich Koester ⁸; Vinzenz Bildstein ⁷

¹ *Ludwig-Maximilians-University, Munich, Germany*

² *Department of Radiation Sciences, Uppsala University, Sweden*

³ *Institut of Radiation and Nuclear Physics, University Bonn, Germany*

⁴ *Department of Subatomic and Radiation Physics, Universiteit Gent, Belgium*

⁵ *CERN, Geneva, Switzerland*

⁶ *Department of Chemistry, University of Oslo, Norway*

⁷ *Technical University, Munich, Germany*

⁸ *ILL, Grenoble, France*

Corresponding Author: wolfgang.schwerdtfeger@physik.uni-muenchen.de

The 1789-keV level in ³⁰Mg was unambiguously identified as the second excited 0+ state by measuring its E0 transition to the ground state via conversion electron spectroscopy. This level corresponds to the strongly deformed intruder 0+ state, thus proving for the first time shape coexistence in this nucleus at the borderline of the 'Island of Inversion' around N ~20. The matrix element $\rho^2(E0, 0+_{2} \rightarrow 0+_{1}) = 3.6(7) E^{-3}$ allows to deduce the squared mixing amplitude a^2 between the unperturbed lowest two 0+ states, indicating a rather weak coupling between the spherical and deformed configurations. One week of beamtime was dedicated for the conversion electron measurement in ³⁰Mg following the beta decay of ³⁰Na which took place in August 2007 at ISOLDE.

Solid State Physics / 13

Lattice location of implanted ions and characterization of implantation-induced damage in Ge

Author: Stefan Decoster¹

Co-authors: André Vantomme ¹; Bart De Vries ¹; Joao Guilherme Correia ²; Ulrich Wahl ³

¹ *Instituut voor Kern & Stralingsfysica and INPAC, K.U.Leuven, Belgium*

² *Instituto Tecnológico e Nuclear, Lisboa, Portugal & CERN-PPE, Geneva, Switzerland*

³ *Instituto Tecnológico e Nuclear, Lisboa, Portugal*

Corresponding Author: stefan.decoester@fys.kuleuven.ac.be

Due to its high mobility of charge carriers, Ge is seen as an important material in future semiconductor technology. Despite the fact that this elemental semiconductor is known for a long time, the knowledge about the lattice location of impurities, the induced structural and electrical damage of Ge after ion implantation is relatively scarce.

Here we present a study of the lattice location of several impurities after ion implantation, in correlation to the structural and electrical implantation-induced damage. The lattice location of several elements (i.e. Na, Fe, Cu, In, Ag, Sn and Er) was examined in detail with the emission channeling (EC) technique using radioactive isotopes produced at ISOLDE.

The EC-experiments clearly indicate that for most of the investigated elements, the substitutional (S) Ge-position is not the overall preferred lattice site, even after annealing up to 500°C. With the exception of Er, no elements occupy the tetrahedral (T) lattice site. However, a somewhat striking result is that next to the S-site, other 'more exotic' lattice sites such as the bond-centered (BC), split

(SP) and anti-bond-centered (AB) site get occupied by some of the implanted ions, before and even after annealing.

The structural quality of implanted Ge samples was also investigated by the Rutherford Backscattering/Channeling (RBS/C) technique as a function of energy, mass and fluence of implanted ions, which helped in explaining the EC results.

Technical Developments / 14

FEBIAD ion source operation modes to tune its selectivity: physics processes, numerical simulations and experimental data.

Author: Liviu Penescu¹

Co-authors: Jacques Lettry¹; Richard Catherall¹; Thierry Stora¹

¹ CERN

Corresponding Author: liviu.penescu@cern.ch

The FEBIAD ("Forced Electron Beam Induced Arc Discharge") ion sources are used for the production of radioactive 1+ ion beams for a wide range of chemical elements. Their small volume and high operating temperature provide good confinement times and ionization efficiencies. Nevertheless, the source lacks in selectivity, ionizing elements regardless of their ionization potential. Presently, the in-source selectivity can be tuned by making use of the different adsorption enthalpy by optimizing the operating temperature, or by creating molecular compounds of the element of interest.

Within the HIGHINT Marie Curie project, theoretical, numerical and experimental investigations are ongoing for the ion source upgrading towards higher operating pressures and, consequently, higher extracted currents.

These investigations let us also tune the selectivity of the source, by operating the source in different modes, which will favor the ionization of a specific class of elements.

When creating a good plasma confinement, the ionization of a gas with a lower ionization potential than the buffer gas should be favored, to the detriment of its residence time.

By using a weak plasma confinement, the ionization selectivity should be increased for elements having different volatilities.

The theoretical model of this approach and the ongoing simulations are presented, as well as some concrete on-line results such as the Chlorine suppression for Argon ionization.

Technical Developments / 15

ECR charge breeding at ISOLDE

Authors: Melanie Marie-Jeanne¹; Pierre Delahaye¹

Co-authors: Charles Barton²; Joakim Cederkall¹; Oliver Kester³

¹ CERN

² University of York

³ GSI

Corresponding Author: pierre.delahaye@cern.ch

At ISOLDE, CERN, an online test bench is dedicated to radioactive ion charge breeding with the Daresbury Phoenix booster.

Two successful experiments have concluded the program of investigation of the performances of the booster undertaken in the frame of the IS397 proposal. The first one consisted in the trapping and charge breeding of ⁶¹Fe daughter nuclides from the beta decay of ⁶¹Mn. As a metallic element, Fe is a

beam difficult to produce at ISOLDE. The second one consisted in the injection of a “cocktail” beam of isobars providing new information on the processes involved in the capture of the 1+ beam.

Nuclear Physics III / 16

Local Probe Studies on Highly Distorted Rare-Earth Manganites

Author: Tania Manuela De Melo Mendonca¹

Co-authors: Andre Pereira ¹; Armandina Lima Lopes ²; Joao Guilherme Correia ³; Joao Pedro Araujo ¹; Joao Sequeira Amaral ²; Pedro Bandeira Tavares ⁴; Vitor Sequeira Amaral ²

¹ IFIMUP, Phys. Dept., FCP, Rua do Campo Alegre 687, P-4150 Porto, Portugal

² Phys. Dept. & CICECO, Universidade de Aveiro, 3810-193 Aveiro, Portugal

³ Instituto Tecnológico e Nuclear, E. N. 10, P-2686-953 Sacavém, Portugal

⁴ Chemistry Dept. and CQ-VR, Universidade de Trás-os-Montes e Alto Douro, P-5000-911 Vila Real, Portugal

Corresponding Author: tania.de.melo.mendonca@cern.ch

The perovskite manganites of the family RMnO₃ (R=Eu, Gd, Ho, Y, Er and Lu) show different competing orders, e.g. orbital, magnetic, and dielectric, that lead to intriguing properties as colossal magnetoresistance and multiferroic behaviour.

Looking forward to investigate locally the interplay between structural, electric and magnetic degrees of freedom, samples have been synthesized via sol-gel methods and studied with the Perturbed Angular Correlation (PAC) technique.

In this first work we have limited the use of PAC to measure the Electric Field Gradient (EFG) / Magnetic Hyperfine Field (MHF) in a series of samples with different rare-earths (R), aiming to find a correlation between these local parameters and the ionic radius of the R element.

At ISOLDE small pellets were implanted with ^{111m}Cd and subsequently annealed to remove implantation defects. The PAC measurements have been performed from 10K up to 473K for the same samples. These results are compared with the ones previously obtained for other perovskite manganites like LaMnO₃, PrMnO₃ and CaMnO₃. Complementary information is obtained through X-ray diffraction data and SQUID measurements on the same samples. The unit-cell parameters, the bond distances and angles are also factors that can be related to the effects under study, and their revision is also proposed in order to complete the microscopic characterization of these compounds.

Solid State Physics / 17

First results from electron emission channeling on-line experiments

Author: Ulrich Wahl¹

Co-authors: Ana Claudia Marques ²; Carlos Pedro Marques ¹; João Guilherme Correia ¹; Lino Pereira ³; Manuel Ribeiro da Silva ⁴; Stefan Decoster ⁵; Tomas Kana ⁶

¹ Instituto Tecnológico e Nuclear ITN

² Centro de Física Nuclear da Universidade de Lisboa and CERN

³ Faculdade de Ciências, Universidade do Porto

⁴ Instituto Superior Tecnico and Centro de Física Nuclear da Universidade de Lisboa

⁵ Instituut voor Kern- en Stralingsfysica Leuven

⁶ Instituut voor Kern- en Stralingsfysica

Corresponding Author: ulrich.wahl@cern.ch

This talk reports on the results from the first electron emission channeling on-line run of the EC-SLI collaboration, which took place during this year's Mn beam time in June. Using our new on-line setup which is equipped with a position-sensitive Si pad detector with fast readout system and which was mounted on the LA2 beam line, two isotopes were successfully used for beta- emission channeling experiments for the first time: ^{56}Mn (2.58 h) and ^{61}Co . While ^{56}Mn was available directly, ^{61}Co was obtained by means of implanting the short-lived precursor isotope ^{61}Mn and exploiting the decay chain ^{61}Mn (4.6 s) \rightarrow ^{61}Fe (6 min) \rightarrow ^{61}Co (1.6 h). We were thus able to determine the lattice location of Mn in GaN and of Co in ZnO in the as-implanted state and following annealing up to 900°C. In both cases it was found that the transition metals preferred substitutional cation (i.e. Ga or Zn) sites.

In order to explore the feasibility of using the isotope ^{27}Mg for lattice location experiments, we also did a brief test of the decay chain ^{27}Na (295 ms) \rightarrow ^{27}Mg (4.6 s). However, in this case it was found that the ISOLDE beam consisted mainly of stable ^{27}Al , making emission channeling experiments using this isotope from UC-W surface ionization targets not feasible.

Nuclear Physics I / 18

Nuclear orientation in transfer reactions

Authors: Dimiter Balabanski¹; Enrico Fiori²; Georgi Georgiev²

¹ Univ. of Camerino & INRNE Sofia

² Cen. de Spect. Nucl. & Spect. Masse (CSNSM)

Corresponding Author: enrico.fiori@cern.ch

Measurements of nuclear magnetic moments are of current interest at RIB facilities, because they provide key information about the states of interest, e.g. about their wave functions.

A basic requirement for the measurement of magnetic moments is the existence of an initially spin-aligned nuclear ensemble. We report here that transfer reactions are applicable for such studies. As the orientation of the nuclear spins is induced directly by the reaction, transfer reactions allow the investigation of nuclear moments of very short lived states, in the range of nanosecond to microseconds. With the development of post-accelerated radioactive beams, available in ISOL facilities, highly-excited neutron-rich isomers will become accessible for measurement. These studies will yield basic information about the configuration of these states and thus provide a stringent test for nuclear theories.

The possibility of using transfer reactions for nuclear moment measurements of isomeric states in PAD experiment will be discussed. The results of four experiments held in the Tandem facility of the Institut de Physique Nucleaire (IPN Orsay, Paris), involving single-nucleon and multi-nucleon reactions will be illustrated, along with advantages and disadvantages of each reaction method. The experiments aimed at studies of the g-factors of the ^{65}mNi 9/2- and ^{66}mCu 6- isomeric states, which were populated in single nucleon transfer reaction (d,p). In the ^{43}mSc case, the 19/2- isomer was populated in the exchange of two neutrons and a proton from the target to the projectile nuclei. The 6- isomeric state in ^{64}mCu was studied with a single-nucleon transfer in inverse kinematics. In all the cases the amplitude of the R(t) function, a measure of the nuclear spin orientation in TDPAD experiment, will be provided. For the case of multinucleon transfer, as different beam energies were used, the orientation is investigated as function of projectile's energy. The results from the fourth experiment have demonstrated the level of difficulties one can encounter using inverse kinematic reactions.

Trapping for Nuclear Physics / 19

WITCH: first measurement and recent developments

Author: Michaël Tandecki¹

¹ *Inst. voor Kern- en Stralingsfysica - Katholieke Universiteit Le*

Corresponding Author: michael.tandecki@cern.ch

We present in detail the status of the WITCH experiment (Weak Interaction Trap for Charged particles). The primary aim is to investigate exotic weak currents in nuclear beta decay by measuring a recoil ion spectrum and deriving the beta neutrino angular correlation coefficient from its shape. At the end of the year 2006, we succeeded for the first time in measuring a recoil ion spectrum using ¹²⁴In. The data have, in the meantime, been thoroughly analysed and the charge state distribution of the recoiling ions has been derived from it. In 2007 a lot effort has been made to improve the precision of the experiment; one of the topics in this respect was the improvement of the Penning traps. A direct result of this is a significant increase in the mass resolution, which is useful to remove isobaric contamination; currently a mass resolution of 200,000-300,000 can be routinely achieved in test conditions.

News from Other Laboratories / 20

Rare isotope beams at MSU - present and future

Author: Georg Bollen¹

¹ *Michigan State University*

Corresponding Author: bollen@nscl.msu.edu

Status and future plans for research with rare isotope beams at the coupled cyclotron facility (CCF) at the National Superconducting Cyclotron Laboratory (NSCL) at Michigan State University (MSU) will be presented with emphasis on the re-acceleration of thermalized beams from projectile fragmentation.

Trapping for Nuclear Physics / 21

20 Years of ISOLTRAP

Author: H.-Juergen Kluge¹

¹ *GSI*

Corresponding Author: j.kluge@gsi.de

Twenty years ago, the first publication on a Penning trap mass measurement of a radionuclide appeared in *Hyperfine Interactions* 38 (1987) 793 with the title "FIRST ABSOLUTE MASS MEASUREMENTS OF SHORT-LIVED ISOTOPES". The authors were Georg Bollen, Phillip Dabkiewicz, Peter Egelhof, Thomas Hilberath, Hartmut Kalinowsky, Franz Kern, Harald Schnatz, Lutz Schweikhard and Helmut Stolzenberg from the Institut für Physik, Universität Mainz, Robert B. Moore from the Foster Radiation Laboratory, McGill University, Montreal, H.-Jürgen Kluge from CERN, Geneva, Switzerland and the Institut für Physik, Universität Mainz and George M. TEMMER, and Gerhard ULM from CERN, Geneva, Switzerland and, last but not least, The ISOLDE Collaboration, CERN. Since then, the masses of several hundred radionuclides have been determined at ISOLDE with ever increasing accuracy, sensitivity and applicability. The talk presents an overview on these developments and on the results obtained by use of ISOLTRAP.

Solid State Physics / 22

Mössbauer studies of dilute magnetic semiconductors (IS-443)

Author: Roberto Mantovan¹

Co-authors: Deena Naidoo²; Gerd Weyer³; Guido Langouche⁴; Haraldur Pall Gunnlaugsson³; Karl Johnston⁵; Krish Bharuth-Ram⁶; Marco Fanciulli¹; Rainer Sielemann⁷; Sveinn Olafsson⁸

¹ CNR-INFM MDM National Laboratory

² School of Physics, University of the Witwatersrand, South Africa

³ Department of Physics and Astronomy, University of Aarhus

⁴ Department of Physics, Katholieke Universiteit, Leuven, Belgium

⁵ EP Division, CERN, Geneva, Switzerland

⁶ School of Physics, University of KwaZulu-Natal, Durban, South Africa

⁷ Hahn-Meitner Institute, Berlin, Germany

⁸ Physics Department, Science Institute University of Iceland, Iceland

Corresponding Author: roberto.mantovan@mdm.infm.it

Mössbauer studies of dilute magnetic semiconductors (IS-443)

R. Mantovan¹, M. Fanciulli¹, H.P. Gunnlaugsson², G.Weyer², R. Sielemann³, D. Naidoo⁴, K. Bharuth-Ram⁵, S. Olafsson⁶, G. Langouche⁷, K. Johnston⁸

¹Laboratorio Nazionale MDM CNR-INFM, Agrate Brianza (MI) 20041, Italy

²Department of Physics and Astronomy, University of Aarhus, DK-8000 Århus C, Denmark

³Hahn-Meitner Institute, 14109 Berlin, Germany

⁴School of Physics, University of the Witwatersrand, WITS 2050, South Africa

⁵School of Physics, University of KwaZulu-Natal, Durban 4041, South Africa

⁶Physics Department, Science Institute University of Iceland, Iceland

⁷Department of Physics, Katholieke Universiteit, Leuven, Belgium

⁸EP Division, CERN, CH-1211 Geneva 23, Switzerland

The study of materials showing multifunctional properties at room temperature (RT) is a challenging task for material science and of importance for future applications in spin-based electronics. The prediction of a Curie temperature above RT in Mn-doped ZnO [1] started a new field of research, with the aim to investigate magnetism in normally non-magnetic oxides. Despite the enormous experimental and theoretical efforts, the origin of the magnetism in (3d-elements) doped and pure ZnO is still a matter of debate in the scientific community. Calculations showing a vacancy-driven mechanism for the magnetism in ZnO have been proposed [2, 3], but clear experimental evidences for such mechanisms are still lacking.

In the framework of the experiment IS-443 at ISOLDE-CERN, we apply Mössbauer spectroscopy to investigate the magnetic properties in ZnO upon implantation with radioactive $^{57}\text{Mn}^+$ ($T_{1/2} = 1.5$ min), decaying to the ^{57}mFe Mössbauer state ($T_{1/2} = 100$ ns). We study the electronic and magnetic configurations of Fe atoms in the ZnO crystals, and the interaction between the Mn/Fe atoms with the defects induced during the implantation process [4]. Our results show that the majority of the Fe atoms are located on Zn sites in a high-spin Fe^{3+} state at ≈ 600 K, giving a strong magnetic contribution in ZnO. The formation/annealing of the magnetism in ZnO is interpreted as occurring/disappearing upon the association/dissociation of Mn/Fe complexes with the lattice defects created in the implantation process [4]. We present an overview of our experimental findings focusing on the essential role played by the lattice defects in observing magnetism in ZnO.

[1] T. Dietl, H. Ohno, F. Matsukura, J. Cibert., and D. Ferrand, *Science* 287, 1019 (2000).

[2] N. A. Spaldin, *Phys. Rev. B* 69, 125201 (2004).

[3] A. Debernardi and M. Fanciulli, *Appl. Phys. Lett.* 90, 212510 (2007).

[4] G. Weyer, H.P. Gunnlaugsson, R. Mantovan, M. Fanciulli, D. Naidoo, K. Bharuth-Ram, T. Agne, *J. Appl. Phys.*, in print.

Results with ISCOOL a new Radio Frequency Quadrupole Cooler and Buncher at ISOLDE

Author: Hanna Franberg¹

Co-authors: Erwin Siesling¹; Ivan Podadera²; Magnus Eriksson¹; Pascal Fernier¹; Pierre Delahaye¹; Tim Giles¹

¹ ISOLDE/CERN

² Accelerator Group, CIEMAT, Madrid, Spain

Corresponding Author: hanna.franberg@cern.ch

We will present the installation of the ISCOOL in the HRS beam line section and give the results from the tests performed during the on-line commissioning in November 2007.

News from Other Laboratories / 24

News from the RIKEN RI beam factory

Author: Heiko Scheit¹

¹ RIKEN Nishina Center

Corresponding Author: scheit@ribf.riken.jp

The RIKEN Radioactive Ion Beam Factory, a major upgrade of the existing heavy-ion accelerator facility at RIKEN, has recently been commissioned. Three new ring cyclotrons boost the beam energies to 440 and 350 MeV/u for light and heavy ions, respectively with an intensity goal for Uranium beams of 1p micro A.

From these primary beams intense secondary radioactive ion beams will be produced by the large acceptance, two-stage in-flight RI fragment separator BigRIPS.

A wide array of experiment equipment is existing, under construction or planned. The facility, major experimental equipment and first results of their operation will be presented.

25

Bremsstrahlung accompanied α -decay ^{210}Po

Author: Heiko Scheit¹

¹ Max-Planck-Institut fuer Kernphysik (MPI)

Corresponding Author: scheit@ribf.riken.jp

A high-statistics measurement of bremsstrahlung emitted in the alpha-decay of ^{210}Po has been performed. The measured differential emission probabilities are in good agreement with our theoretical results obtained within the quasiclassical approximation as well as with an exact quantum mechanical treatment. Due to the small effective electric dipole charge of the radiating system, a significant interference between the electric dipole and quadrupole

contributions occurs, altering substantially the angular correlation between the alpha particle and the emitted photon.

Trapping for Nuclear Physics / 26

Nuclear mass measurements for nucleosynthesis studies at ISOLTRAP

Author: Martin Breitenfeldt¹

Co-author: Lutz Schweikhard¹

¹ *Ernst-Moritz-Arndt-Universitaet Greifswald*

Corresponding Author: martin.breitenfeldt@cern.ch

In 2007 the mass measurements at the tandem Penning trap mass spectrometer ISOLTRAP focussed on nuclear structure and nucleosynthesis studies. The masses of about 20 nuclides have been measured with relative mass uncertainties as low as $1 \cdot 10^{-8}$. The beam times were dedicated to the mass determination of neutron deficient and neutron rich nuclides relevant for the investigation of the rp- and r-process, respectively. For neutron-deficient Cd isotopes, close to the doubly magic 100Sn and the end-point region of the rp-process, the first direct mass measurement of 99Cd was performed, giving accurate data at the neutron shell closure $N=50$. Furthermore, measurements aimed at the masses of neutron-rich Ag and Cd nuclides which will give input values for astrophysics calculations for the r-process, especially at the waiting point 130Cd .

Nuclear Physics III / 27

Status report on g-factor measurements on Coulomb excited ^{138}Xe nuclei using the Transient-Field technique (IS415)

Author: Joerg Leske¹

Co-authors: Andrea Jungclaus²; Karl-Heinz Speidel³

¹ *IKP, TU Darmstadt, Germany*

² *UAM, Madrid, Spain*

³ *HISKP, University of Bonn, Germany*

Corresponding Author: joerg.leske@cern.ch

Two experimental campaigns have been carried out at REX-ISOLDE and MINIBALL in order to determine the magnetic moment of the first excited state in ^{138}Xe with the technique of transient magnetic fields. ^{138}Xe ion beams of 2.9 MeV/u and different multilayered targets were used for Coulomb excitation. De-excitation gamma rays were measured in four MINIBALL cluster detectors in coincidence with forward scattered target ions. We will report on our achievements and present first results. Technical improvements as well as experimental limits will be discussed followed by a vision of future g-factor projects on REX-ISOLDE.

Nuclear Physics I / 28

REX-ISOLDE - now also heavyweight champion

Author: Fredrik John Carl Wenander¹

¹ CERN

Corresponding Author: fredrik.wenander@cern.ch

A summary of the advances at the REX-ISOLDE facility during 2007 will be given. In total 3 new elements and 8 new isotopes were delivered for physics, coming either as atomic ions or molecular ions from ISOLDE. The REX-ISOLDE post accelerator has this year also seriously entered the arena of heavy ion beams by accelerating light Hg isotopes to full linac energy. The potential and constraints in terms of the machine for this 'heavyweight' region will be discussed. One of the experimental runs could be carried out in spite of a malfunctioning trap by operating the EBIS in continuous injection mode. This opens up future possibilities with the RFQ cooler now installed and operational. A number of technical developments were carried out at REX, for example: the extension of the beam transport line and move of the experimental stations were completed; new control systems were implemented and the linac RF amplifiers were consolidated. In addition, we will discuss foreseen accelerator improvements and beam developments.

Nuclear Physics II / 29

Beta-asymmetry measurements as a probe for non standard model physics

Author: Frederik Wauters¹

Co-author: Nathal Severijns¹

¹ *katholieke universiteit leuven*

Corresponding Author: frederik.wauters@fys.kuleuven.be

The status of the beta-asymmetry measurements project will be presented. In these experiments we measure the beta-asymmetry parameter which is sensitive to deviations from the $V - A$ character of the weak interaction, more precisely to tensor currents. The experimental technique we use to create a polarized ensemble of nuclei is low temperature nuclear orientation (LTNO).

The first (preliminary) results for the beta-asymmetry parameter will be presented for ^{60}Co and ^{114}In . These measurements were performed in Leuven and Louvain-la-Neuve preceding the on-line measurements with ^{67}Cu at ISOLDE in the summer of 2006 and 2007. The first run ^{67}Cu was successful but had a lack of statistics due to the low yield. This year the statistics was increased with a factor of 4 to 5. The data analysis is ongoing. ^{67}Cu has a high sensitivity to possible tensor contributions in the weak interaction. We aim to improve the limits for tensor couplings which are currently of the order of 8 % in the amplitudes.

To reach the required precision we are using extensively GEANT4 Monte-Carlo simulations to get control of systematic effects such as scattering and magnetic field effects. A part of our research is therefore focussing on the performance of GEANT4 for beta decay measurements.

HIE-ISOLDE / 30

First laser measurements on radioactive beams using ISCOOL

Author: Mane Junior Ernesto Batista¹

Co-authors: Ari Jokinen²; Bradley Cheal³; Christopher Geppert⁴; David Forest⁵; Deyan Yordanov⁶; Gary Tungate⁵; Gerda Neyens⁷; Hannah Franberg⁸; Iain Moore²; Ivan Podadera⁹; Jonathan Billowes³; Jorg Kramer⁶; Juha Aysto²; Kieran Flanagan¹⁰; Klaus Blaum⁴; Magdalena Kowalska⁸; Mats Lindroos⁸; Paul Campbell³; Peter Lievens¹⁰; Pierre Delahaye⁸; Pieter Vingerhoets¹⁰; Reiner Neugart¹¹; Richard Catherall⁸; Wilfried Noertershaeuser⁴

¹ *Dept. of Phys. & Astronomy, Schuster Lab, University of Manchester*

² *University of Jyväskylä*

³ *Dept. of Phys. & Astronomy, Schuster Lab, University of Manchester*

⁴ *Gesellschaft fuer Schwerionen forschung mbH (GSI)*

⁵ *School of Physics and Astronomy, University of Birmingham*

⁶ *Johannes-Gutenberg-Universitaet*

⁷ *Inst. voor Kern- en Stralingsfysica, Katholieke Universiteit Leuven*

⁸ *CERN*

⁹ *Cent. Invest. Energ. Medioamb. Tec. (CIEMAT), Consejo Sup. de Investigac. Cientif (CSIC)*

¹⁰ *Katholieke Universiteit Leuven*

¹¹ *Institut fur Physik, Johannes-Gutenberg-Universitaet*

Corresponding Author: ernesto.mane.junior@cern.ch

Collinear laser spectroscopy provides an excellent diagnostic tool for characterizing the ion beam extracted from the ISOLDE RFQ cooler-buncher.

Potassium and rubidium beams prepared in ISCOOL were delivered to the COLLAPS beam line and focussed to give 50% transmission through a 1mm iris. The iris was used to overlap a co-propagating Ti:Sa laser. The laser was tuned to resonance on a D-line component (766 nm for K, 780nm for Rb). The time-structure of the ion beam was determined from the fluorescent photon arrival times. This naturally eliminated any contribution from molecular ions formed in the RFQ, which cannot be easily discriminated if ion-detection were used.

The effects of using different extraction potential gradients on the ion beam bunch width were studied. The energy spread was characterized by taking fluorescence spectra obtained using the COLLAPS laser setup. This was measured for each of the extraction potential gradients. An upper limit of the ion energy spread was determined.

To complete these tests, the cooler was tuned to provide bunches of 20 micro-seconds with low energy spread for a demonstration of the bunched-beam laser spectroscopy application on neutron-rich radioisotopes of potassium.

Nuclear Physics II / 31

Studies of the ^{12}C nucleus using beta-decay of ^{12}N and ^{12}B

Author: Solveig Hyldegaard¹

¹ *Department of Physics and Astronomy, University of Aarhus*

Corresponding Author: sgp@phys.au.dk

Experimental studies of the ^{12}C nucleus are important to both nuclear theory and astrophysics. Improved knowledge of states in ^{12}C is necessary for our understanding of the triple alpha reaction in stars. Especially the Hoyle state at just 0.38 MeV above the triple alpha threshold is vital for this process, but also a predicted 2^+ state at 1.7 MeV above the triple alpha threshold in ^{12}C has been included in the NACRE reaction rate calculations, and its existence needs to be verified experimentally.

The structure of the ^{12}C nucleus is not fully understood. Some properties are well described by cluster models and others by mean field theory. New ab-initio calculations have been published and the results are promising giving good agreement with experimental data in the literature.

In a new experiment at KVI in Groningen, the Netherlands, ^{12}C was populated in beta-decay of ^{12}B and ^{12}N . Beams of ^{12}B and ^{12}N ions were implanted in a 48 times 48 strip detector (DSSSD) and let decay. Because of the segmentation of the DSSSD into very small pixels the background of beta particles is significantly reduced and essentially confined to low energies. The implanted ions and emitted alpha particles are stopped in one pixel of the detector so very precise absolute branching ratios and decay spectra can be obtained. Measurements at very low energies are possible because detector deadlayer effects are avoided in the implantation method, so new information about the Hoyle state can be achieved. Results of the analysis will be presented including new energy spectra, branching ratios and comparisons to earlier work and theory.

Trapping for Nuclear Physics / 32

JYFLTRAP at IGISOL - Atomic masses from nuclear structure physics to fundamental physics

Author: Ari Jokinen¹

¹ *University of Jyväskylä*

Corresponding Author: ari.jokinen@phys.jyu.fi

JYFLTRAP and IGISOL-facility has been in high-precision operation now for few years producing vast amount of atomic mass data for nuclear structure studies and fundamental physics. All in all more than 200 atomic masses of ground and isomeric states has been measured from sub-keV to few keV precision. In this contribution the recent highlights from the facility will be reviewed and few examples will be described in more detail.

In neutron-rich side of the nuclide chart proton-induced fission of ^{238}U has been applied to produce isotopes from Ni to Pd for mass spectroscopy in JYFL. In this contribution, new data on the evolution of the N=50 shell gap will be discussed.

On neutron-deficient side, a large fraction of our mass measurement program has concentrated on the refractory isotopes below 100Sn. These measurements contribute to nuclear structure and nuclear astrophysics questions in this region. In this contribution, recent measurements around ^{94}Ag will be discussed in detail to shed light on the puzzle of highly disputed two-proton decaying high-spin (21+) isomer in ^{94}Ag .

Finally, precision Q-value measurements for super allowed beta decay and double beta decay will be discussed in context of CVC-hypothesis, unitarity of CKM-matrix and neutrino physics.

Nuclear Physics III / 33

Coulomb excitation of neutron-rich ^{44}Ar at SPIRAL

Author: Magda Zielinska¹

Co-authors: Alexander Buerger²; Andreas Goergen¹; Cedric Dossat¹; Christophe Theisen¹; Daniel Pietak³; Emmanuel Clement⁴; Geirr Sletten⁵; Jędrzej Iwanicki⁶; Joa Ljungvall¹; Julian Srebrny⁶; Katarzyna Wrzosek⁶; Pawel Jan Napiorkowski⁶; Wilton Catford⁷; Wolfram Korten¹

¹ *CEA Saclay*

² *University of Bonn*

³ *Warsaw University of Technology*

⁴ GANIL⁵ NBI Copenhagen⁶ HIL Warsaw⁷ University of Surrey**Corresponding Author:** magda@slcj.uw.edu.pl

The weakening of the N=28 shell closure for neutron-rich nuclei and the development of deformation and shape coexistence in this mass region were addressed in a low-energy Coulomb excitation experiment using a radioactive ⁴⁴Ar beam from SPIRAL facility of GANIL. The Ar⁴⁴ ions were post-accelerated to 2.7 and 3.7 MeV/nucleon and Coulomb excited on ¹⁰⁹Ag and ²⁰⁸Pb targets, respectively. The scattered projectiles and recoiling target nuclei were detected in a highly segmented double-sided silicon detector and the gamma rays were detected with the EXOGAM germanium detector array. In addition to the first excited 2+ state, one higher-lying level was populated and two gamma lines from its deexcitation were observed. With the ¹⁰⁹Ag target both projectile and target nuclei were excited, which has been used to normalize the excitation probability in ⁴⁴Ar with the well known transition strengths in ¹⁰⁹Ag. The level of statistics was sufficient to subdivide the data into several sub-sets corresponding to different ranges of scattering angles. Since the influence of the quadrupole moment of the first 2+ state on its excitation probability varies significantly with scattering angle, it was possible to obtain from one experiment information on both the transitional and diagonal matrix elements involved in the excitation process.

Trapping for Nuclear Physics / 34

TRIGA-TRAP: Mass measurements on exotic and heavy nuclides

Author: Dennis Neidherr¹**Co-authors:** Christian Smorra¹; Christine Weber²; Frank Herfurth³; Georg Eitel¹; Jens Ketelaer¹; Julia Repp¹; Klaus Blaum⁴; Klaus Eberhardt⁵; Rafael Ferrer¹; Sebastian George⁴; Szilard Nagy¹; Wilfried Noertershaeuser⁵¹ Institut für Physik - Johannes Gutenberg-Universität² Department of Physics - University of Jyväskylä³ GSI Darmstadt⁴ Institut für Physik - Johannes Gutenberg-Universität, GSI Darmstadt⁵ Institut für Kernchemie - Johannes Gutenberg-Universität**Corresponding Author:** dennis.neidherr@cern.ch

The TRIGA Mainz research reactor offers unique possibilities for online mass measurements on neutron-rich isotopes as produced by fission of U-235, Pu-239 or Cf-249 targets. A new Penning trap mass spectrometer will be installed, featuring not only the commonly used time-of-flight resonance technique, but also the non-destructive narrow-band image current technique, enabling the detection of a single singly-charged ion stored in the trap. TRIGA-TRAP is the first online mass spectrometer for singly-charged heavy ions using this image current detection technique in combination with cryogenic Penning traps. In case of many heavy and superheavy nuclides, the production rates are often less than a few ions per second, but some isotopes exhibit comparably long half-lives in the order of seconds, which allows for repeated measurement cycles of the same trap content. The status of TRIGA-TRAP will be presented.

Nuclear Physics I / 35

Shell structure and shape changes in neutron rich krypton isotopes

Author: Dennis Muecher¹

Co-author: Jędrzej Iwanicki²

¹ *Institut for Nuclear Physics, University of Cologne*

² *Heavy Ion Laboratory, Warsaw*

Corresponding Author: muecher@ikp.uni-koeln.de

The krypton isotopes are located roughly in the middle of the Z=28 and Z=50 shell closures, where competition of various structures and shapes at low-excitation energy occurs. For example, this can be seen in the case of Z=40 Zirconium Isotopes. In ⁹⁶Zr (N=56) a strong shell closure was found, interpreted to be reinforced by the Z=40 shell gap. Such a shell gap should quickly disappear moving away from Z=40. Indeed, no influence of this gap was found for isotopes Z>40. Adding more neutrons to the Zirconium Isotopes leads to a unique increase of B(E2) values from 8 W.u. to about 100 W.u. by adding only a few neutrons. This abrupt phase transition from spherical to deformed nuclei may demonstrate the complexity of this region of the nuclear chart.

In the case of Krypton isotopes (Z=36) the energy of the first excited state peaks at ⁹²Kr (N=56) like in the case of Zirconium isotopes. In the past it was extensively discussed whether the N=56 shell gap is active or deformation starts to set in. Up to now, only information about energies and masses were available for the neutron rich krypton isotopes to discuss this topic. We were able to measure B(E2) values for ⁸⁸Kr (N=52) and ⁹²Kr (N=56) for the first time by Coulomb excitation in inverse kinematics at REX-ISOLDE using the MINIBALL array. The analysis was done using the newly developed Coulex Code Gosia2, that can handle target and projectile excitations simultaneously. We present our results and discuss the consequences for the understanding of the structure of neutron-rich krypton isotopes.

Solid State Physics / 36

Magneto-resistive materials: hyperfine studies using radioactive isotopes.

Author: Vitor Amaral¹

Co-authors: Armandina Lima Lopes²; João Amaral³; João Guilherme Martins Correia⁴; João Nuno Gonçalves¹; João Pedro Araújo⁵; Pedro Tavares⁶; Tânia Mendonça⁴

¹ *Universidade de Aveiro*

² *Div. PH-UIS, CERN, Universidade do Porto and Universidade de Aveiro, Portugal*

³ *Universidade de Aveiro, Portugal*

⁴ *Div. PH-UIS, CERN and ITN Sacavém, Portugal*

⁵ *IFIMUP, Universidade do Porto, Portugal*

⁶ *Departamento de Química, UTAD, Portugal*

Corresponding Author: vamaral@ua.pt

This year the Physics Nobel Prize was awarded to Albert Fert and Peter Grunberg for the discovery of the magnetic effect of giant magnetoresistance, the basis of today's magnetic recording in computer hard disks. This discovery is an example of using magnetism to control the electrical current flow through materials built at the nanometer scale. More recently, colossal magnetoresistance was discovered in magnetic oxides such as mixed valence manganites, leading to even larger changes of resistance (up to several orders of magnitude). In this talk the properties associated to spin dependent electron transport are addressed. The relation to local structure, spin-polarized electron band and charge localization phenomena is discussed. The talk will point out the application of hyperfine studies, performed using radioactive isotopes, mainly at ISOLDE, that contribute to explore this exciting research field.

News from Other Laboratories / 37**Hyperdeformation in Xe Nuclei.****Author:** Geirr Sletten¹¹ *NBI University of Copenhagen*

Nuclear structure physics is testing nuclei to the limits of their existence by imposing extreme values of Isospin, Excitation energy and Angular momentum on them. In hyperdeformed nuclei the deformation parameter is the central issue. Theoretical predictions are that nuclei at extreme angular momentum might acquire shapes with small axis ratio 3:1 and that rotational bands might exist in the corresponding potential well. Signals from hyperdeformed structures have been in the experimental focus for more than a decade and whether the existence of hyperdeformation at high spin has been firmly identified, is still debated. A recent experiment at ANL using the GAMMASPHERE array provides preliminary results on hyperdeformation in ¹²⁴Xe, and forms the basis for the present report. So far, discrete line rotational bands corresponding to hyperdeformation have not been identified, but weak signals from the analysis of unresolved spectra indicate cascades with a moment of inertia $128h^2 \text{ MeV}^{-1}$ which is expected for hyperdeformed rotors.

Opening Session / 38**Between Bose-Einstein condensate and fermionic superfluid.****Author:** Piotr Magierski¹¹ *Warsaw University of Technology***Corresponding Author:** magiersk@if.pw.edu.pl

In the last couple of years we have witnessed a tremendous progress in the field of cold fermionic atoms. Ultra cold atomic gases provide a remarkable opportunity to investigate strongly correlated Fermi systems. They are dilute and their interactions can be precisely controlled over an enormous range. In particular, they form unique laboratories where the crossover between the Bose-Einstein condensate and the BCS superfluid can be explored. In this limit, which is relevant for the dilute neutron matter, the scattering length greatly exceeds the average inter-particle separations. Consequently, the system is believed to be strongly paired and the size of Cooper pairs is comparable to the Fermi wavelength.

On the theoretical side, our overall understanding of these remarkable many body systems has improved tremendously, even though many questions remain yet unanswered. During the talk I shall discuss the properties of dilute and strongly interacting Fermi system, emphasizing the link with the nuclear matter properties and the neutron star structure. I will discuss the equation of state for fermions in the so-called unitary regime, where the scattering length tends to infinity. The comparison with recent experiments concerning trapped atomic gases will be presented.

Opening Session / 39

Double-beta decay and neutrino physics

Author: Christina Volpe¹

¹ *IPN, Orsay*

Corresponding Author: volpe@ipno.in2p3.fr

In the last decade many spectacular advances have been performed in neutrino physics. Still, crucial questions remain open, in particular the knowledge of the Dirac versus Majorana nature of neutrinos. The most feasible way of knowing the answer to this issue is through the observation of a nuclear process called neutrinoless double-beta decay.

Its search presents serious experimental and theoretical challenges, in particular, the precise knowledge of the nuclear matrix elements determining the half-lives.

In this talk, we will review the status in this field and emphasize the difficulties as well as the crucial role played by nuclear physics.

We will discuss that beta-decay, muon capture and charge-exchange studies offer important constraints to the neutrinoless half-life predictions.

We will conclude discussing the interest of neutrino-nucleus interaction studies.

Nuclear Physics I / 40

Coulomb excitation of ³¹Mg - at the shore of the island of inversion

Author: Peter Reiter¹

¹ *IKP, Cologne*

The first safe Coulomb excitation measurements of ³⁰Mg and ³²Mg at safe energies with REX-ISOLDE and MINIBALL clearly confirms the transition in the neutron-rich magnesium isotopes from normal to intruder-dominated configurations between these two Mg isotopes by measuring the B(E2;

0g_s⁺-21⁺) values of the even-even nuclei. A recent Coulomb excitation

experiment with a post-accelerated odd ³¹Mg beam from REX-ISOLDE showed the transition from the 945 keV state into the first excited state at 50.5 keV. The ongoing analysis will concentrate on the quadrupole properties of the 945 keV state, the intriguing interpretation as a 5/2 state of the positive yrast band of ³¹Mg on top of the 1/2⁺ ground state and comparison with recent shell model calculations [1].

[1] F. Marechal, et al., Phys. Rev. C 72, 044314 (2005)

Nuclear Physics III / 41

Inelastic Branch of the Stellar Reaction ¹⁴O(α,p)¹⁷F

Author: Philip Woods¹

¹ *University of Edinburgh*

The miniball and CD array have been used on Rex Isolde to measure particle gamma-coincidences between inelastically scattered protons and gamma-rays from the first excited state of ¹⁷F. This method can be used to infer the significance of the inelastic branch in the ¹⁴O(α p)¹⁷F reaction which is a key reaction in X-ray bursters scenarios. Current analysis of the results from this recent experiment will be reported.

HIE-ISOLDE / 42

Future physics at HIE-ISOLDE: a personal view

Author: Philip Woods¹¹ *University of Edinburgh*

The Rex Isolde, and Miniball and CD arrays, already constitute a world class facility for nuclear structure physics with radioactive beams, with nuclear astrophysics studies also starting up. The future HIE-ISOLDE project promises to open up significantly the possibilities for new physics with the uniquely wide range of beam species available on Rex Isolde. The talk will explore these possibilities and the implications for plans for new equipment developments.

Opening Session / 43

Gamma-ray nucleosynthesis: results from the INTEGRAL satellite

Author: Nami Mowlavi¹¹ *Observatory of Geneva***Corresponding Author:** nami.mowlavi@obs.unige.ch

The International Gamma Ray Astrophysics Laboratory has been launched in 2002 to observe the Universe at energies between a few keV to a few MeV. Among the objectives was the study of gamma-ray lines, probing, among others, specific nucleosynthesis processes that occur in stars. I present a review of the observations performed by INTEGRAL in this respect, and discuss them in relation with the astrophysical sites at the origin of those gamma-ray nuclei.

Trapping for Nuclear Physics / 44

Masses of halo nuclides and decay-spectroscopy in the lead region: new developments at ISOLTRAP

Author: Magdalena Kowalska¹¹ *CERN***Corresponding Author:** kowalska@cern.ch

The ISOLTRAP experiment has up to now provided precise values of masses of more than 300 radionuclides with relative uncertainty below 10^{-7} and many cases reaching 10^{-8} . Presently, there are two main projects under way at ISOLTRAP, which aim at widening the number of nuclides to study and the type of the performed measurements. The first is devoted to studying masses of halo nuclei, which provide important input e.g. for deriving their charge radii. Since the present cooling scheme using helium is not effective for such light nuclides, hydrogen cooling has to be used instead, and necessary simulations and tests are presently under way. The other development will allow using ISOLTRAP for isobaric and even isomeric purification to performing beta- and gamma-decay studies. Presently, the setup is being prepared for the measurements on neutron-rich Hg and Tl isotopes beyond $N=126$. In this talk I will present the physics motivation of these two developments, their status and future plans.

Opening Session / 45**The first 40 years of physics at ISOLDE****Author:** Björn Jonson¹¹ *Chalmers University of Technology***Corresponding Author:** bjn@fy.chalmers.se

In this talk I will give some selected comments to the present and future physics at ISOLDE, REX ISOLDE and HIE-ISOLDE. I will also give examples of some of the pioneering experiments at ISOLDE that have led to the success of this field of nuclear physics. As a basis for my talk I give below some dates and developments that have been important on the way towards the present Facility. 1951: The first experiment of ISOL type was carried out by Otto Kofoed-Hansen and Karl-Ove Nielsen in Copenhagen, Denmark. 1961: The Nuclear Chemistry Group built an electromagnetic isotope separator and showed that the capacity for production of radioactive nuclides in an internal beam of the SC was very high. 1964: December 17. CERN decides to build an ISOL(DE) experiment at the SC. 1967: September 23. The first experiment was performed at ISOLDE. The results were published in a paper with 29 authors [1]. 1973: Shutdown and reconstruction of ISOLDE. The external beam from the SC was increased to 4 uA. 1974: First experiments at ISOLDE 2. 1987: Installation of a high-resolution separator in the SC proton hall. 1989: Start of the civil engineering work for the ISOLDE site close to the PS Booster. 1990: December 17, 12.00. The SC was shut down. 1992: June 26. The first experiment was performed at the ISOLDE PSB Facility. 2001: October 30. The first beams were delivered from REX ISOLDE. [1] Decay Characteristics of Short-Lived Radio-Nuclides Studied by On-Line Isotope Separator Techniques. P.G.Hansen, P.Hornshøj, H.L.Nielsen, K.Wilsky, H.Kugler, G.Astner, E.Hagebo, J.Hudis, A.Kjelberg, F.Munnich, P.Patzelt, M.Alpsten, G.Andersson, A.Appelqvist, B.Bengtsson, R.A.Naumann, O.B.Nielsen, E.Beck, R.Foucher, J.P.Husson, J.Jastrzebski, A.Johnson, J.Alstad, T.Jahnsen, A.C.Pappas, T.Tunaal, R.Henck, P.Siffert, G.Rudstam, Phys.Lett. 28B, 415 (1969).

News from Other Laboratories / 46**The ISAC facility at TRIUMF and Halo-nuclei mass measurements at TITAN****Author:** Jens Dilling¹¹ *TRIUMF***Corresponding Author:** jdilling@triumf.ca

The ISAC facility at TRIUMF is providing over 200 different radioactive beams to users at energies up to 5 MeV/u. Plans exist to extend the facility by an additional 200 μA proton beam line and a photo-fission facility based on a mega-watt class superconducting electron linac.

One of the experimental facilities at ISAC is TITAN, a high precision Penning trap mass spectrometer.

First experiments with radioactive beams were carried out this year with the aim to determine the masses of halo nuclei. In a first run, the mass of Li-8 and Li-9 were investigated with a precision of about $2 \cdot 10^{-8}$.

The goal of an upcoming experiment (Dec 2007) is the mass measurement of Li-11.

The talk will give an overview of ISAC and present the future plans. It will introduce the TITAN facility and show first experimental results.

Nuclear Physics II / 47**MAYA active target for the study of extremely exotic nuclei**

Author: Herve Savajols¹

¹ *GANIL*

Corresponding Author: savajols@ganil.fr

This talk will discuss some recent experiments using the MAYA active target at GANIL and TRIUMF.

48

Shell structure evolution in nuclei : future collaboration between GANIL and

Author: Faical Azaiez¹

¹ *IPNO*

Corresponding Author: azaiez@ipno.in2p3.fr

I will try in my talk to single out some ideas of collaboration between GANIL and REX-ISOLDE for the study of shell structure evolution in nuclei. Among other possibilities, this could be based on the exchange and/or merge of detections systems available in both facilities which should increase the experimental capabilities and allow the study of weekly produced keys nuclei.

HIE-ISOLDE / 49**The HIE-ISOLDE project**

Author: Matteo Pasini¹

¹ *CERN*

Corresponding Author: matteo.pasini@cern.ch

This talk will present the plans for the HIE-ISOLDE project.

Solid State Physics / 50**Soft-landing of Radioactive Probes on Clean Metal Surfaces and at Interfaces**

Author: W.-D. Zeitz¹

¹ *Hahn Meitner Institute*

Soft-landing techniques of radioactive probe atoms have been developed in the ASPIC facility which is installed at ISOLDE. The probe atoms were deposited at interfaces and on surfaces of metals and ferromagnets where they can be manipulated to occupy specific sites. Utilizing sensitive nuclear techniques for analysis, magnetic hyperfine fields and electric field gradients have been determined. In the light of density-functional methods, these parameters can be linked to parameters which are commonly used in complementary techniques in solid state physics.

A review shall be given over the achievements of the measurements and targets for further investigations shall be identified.

Technical Developments / 51

Target and Ion Source Development

Author: Thierry Stora¹

¹ CERN

The activities related to the development of new radioactive ion beams at ISOLDE will be reviewed, with an emphasis on high power targetry. Some items will be more precisely presented. New sub-micrometric and nanomaterials are under development at ISOLDE and within the EURISOL Design Study. Their use will be illustrated with the delivered ²¹Mg and ¹⁷F beams produced with pure submicrometric SiC targets. Post-irradiation analysis of the target material has started. Preserving good extraction efficiencies when dealing with high incoming primary driver beams is a challenging task. A prototype testing the characteristics of a multibody target unit linked to a single FEBIAD ion source has been operated for the first time this year at ISOLDE. This test opens up the possibility of accommodating 100kW incoming primary beam driver while preserving compact target characteristics and good ionization efficiencies.

Nuclear Physics III / 52

Pb(II) and Hg(II) binding to proteins and model systems studied by ²⁰⁴mPb- and ¹⁹⁹mHg-Perturbed Angular Correlation of γ -rays (PAC) spectroscopy

Author: Johan Vibenholt¹

¹ University of Copenhagen

We aim to elucidate the fundamental chemistry of heavy metal - protein interactions. This involves both studies of model systems, designed proteins and naturally occurring proteins, as well as the interpretation of the spectroscopic data using quantum mechanical calculations. We will present data on model systems (coordination compounds) and proteins recorded recently at ISOLDE.

Opening Session / 54

Welcome

Author: Karsten Riisager¹

¹ CERN