

Particle correlations in ALICE

Mónika Kőfaragó
Eötvös Loránd University, Hungary

Supervisor:
Jan Fiete Grosse-Oetringhaus
CERN, ALICE

16th August 2012




- Focusing on Pb-Pb collisions
- Looking for properties of quark-gluon plasma
- Searching for modifications by the medium




- Focusing on Pb-Pb collisions
- Looking for properties of quark-gluon plasma
- Searching for modifications by the medium

- $$R_{AA} = \frac{d^2 N_{ch}^{AA} / d\eta dp_T}{\langle N_{coll} \rangle d^2 N_{ch}^{PP} / d\eta dp_T}$$




Phys. Lett. B 696 (2011) 30-39



- Looking for modification of jets compared to pp collisions
- $\Delta\varphi = \varphi_1 - \varphi_2$, $\Delta\eta = \eta_1 - \eta_2$



- Looking for modification of jets compared to pp collisions
- $\Delta\varphi = \varphi_1 - \varphi_2$, $\Delta\eta = \eta_1 - \eta_2$



- Tracking correction





- Tracking correction



ALICE



- Mixed event





- Tracking correction




- Mixed event




- Background subtraction (η gap subtraction)






0 – 10%




60 – 70%




ALICE


0 – 10%




60 – 70%



pp



- Fit with different functions
- Best: sum of two Gaussian functions
- Width (σ) extracted from the fit





- Add more high p_T bins
- Hard to fit



- Energy: $\sqrt{s_{NN}} = 2.76$ TeV
- 2010: ~ 15 M events
- 2011: centrality trigger \implies 10 times more in central bins
- Goal: have higher statistics \implies unite 2010 and 2011 data \implies cross checks \implies Mixed event, efficiency should be similar
- 2011: Some TPC modules were inactive in a fraction of the data \implies cross checks are needed


- Energy: $\sqrt{s_{NN}} = 2.76$ TeV
- 2010: ~ 15 M events
- 2011: centrality trigger \implies 10 times more in central bins
- Goal: have higher statistics \implies unite 2010 and 2011 data \implies cross checks \implies Mixed event, efficiency should be similar
- 2011: Some TPC modules were inactive in a fraction of the data \implies cross checks are needed

$2.0 < p_{T, \text{trig}} < 3.0 - 1.00 < p_{T, \text{assoc}} < 2.00 - 0-10\%$





- Similar, but $\sim 3\%$ difference \implies have to understand

Efficiency



Efficiency



Summary

- Working mostly on 2010 data
- Started comparison of 2010 and 2011 data

Plans

- Unite all 2011 data
- Unite 2010 and 2011 data
- Include higher p_T bins
- Comparison to AMPT model (**A** **M**ulti**P**hase **T**ransport)
- Comparison to STAR experiment

Thank you for your
attention!