








QCD from CMS



- 1) General problematics
- 2) Where pQCD works.
- 3) Where the pQCD is challenged.
- 4) Pushing pQCD to the limits.
- 5) Few words about the photons.




1.1) Common wisdom about QCD

- QCD is ONE theory: SU3 + free coupling parameter.
 - Well established in ee and ep colliders: α_s value and running, interaction mediator (gluon), SU3 gauge group, pQCD at NNLO, factorization theorem for hadrons PDFs...
- But MANY regimes with associated approximations:
 - NP regime (lattice QCD, phenomenology): spectroscopy
 - Common wisdom: complicated chemistry. 
 - Intermediate (phenomenology): fragmentation functions, UE, Min bias collisions.
 - Common wisdom: “tuning and witchcraft”. 
 - Hard interaction (pQCD): jets physics, inelastic hadrons PDFs.
 - Common wisdom: “well understood, just need to cross check it works”. 
- At CMS: till now O(50) dedicated papers.
- <https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsSMP>
- <https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsFSQ>



1.2) pQCD before the hadron colliders

- I would try to dispel the common wisdom that pQCD is so well understood and show where/why hadron colliders may and shall contribute.
- At LEP/HERA we got used that NLO (or NNLO when possible) contains most of the perturbative effects. UE and parton showering are minor corrections.
- It seems that this is not always true for the hadron colliders. Let's have a look on it.



1.3) Kingdom of the pQCD

- Factorization theorem: hard matrix element calculated at NLO factorized from proton PDFs, parametrized (mainly) in ep collision and evolved by DGLAP evolution at NLO (or NNLO).




“Higgs and SM region”

Terra incognita: forward physics

“Exotica searches region”

Where pQCD works


Central jets production and “Exotica region”




2.1.1) Inclusive hard jets production

$$x = \frac{2p_t}{\sqrt{s}} \exp(\pm y)$$

- pp collisions: 2 ladders.
 - Central jets: symmetric collisions $x_1 \sim x_2$. DGLAP (strong k_T ordering) expected to perform well.
 - Forward jets: asymmetric collisions $x_1 \ll x_2$. May need $\log(1/x)$ resummation and alternative evolution (BFKL, CCFM)



7 TeV LHC parton kinematics



2.1.2) Experimental setup


Trackers
 $|\eta| < 2.5$

**Calorimeters
(EM and Hadronic)**
 $|\eta| < 3.0$

**Hadronic Forward (HF)
Calorimeters**
 $2.9 < |\eta| < 5.2$

Muon Chambers
 $|\eta| < 2.5$

A Compact Solenoidal Detector for LHC



CMS-PARA-001-11/07/97 JLB.PP



- Central region $|\eta, y| < 2.5$: tracker + calorimeters. Particle Flow reconstruction allow very precise measurement by combining all detectors and allow pile-up (PU) removal.
- Forward region: $|\eta, y| > 3.0$: calorimeters only, but jets collimated and have large energy. Low handle on PU and large UE.

2.2.1) Measurement of inclusive jets

$$\frac{d\sigma_{Data}^2}{dp_T dy} = \frac{d\sigma_{Det}^2}{dp_T dy} C_{Det}$$


$$\frac{d\sigma_{Thr}^2}{dp_T dy} = \frac{d\sigma_{NLO}^2}{dp_T dy} C_{NP}$$

- Measurement: corrected for detector effect to “hadrons level” - C_{DET} .
- NLO calculation: corrected for hadronization and MPI effects estimated from LO+PS MC - C_{NP} .



2.2.2) Inclusive jets : 7 TeV with R=0.7

CMS-PAS-11-004





Jets reconstructed with sequential anti-k_T algorithm with R=0.7



- Generally excellent agreement over 12 orders of magnitude with NLO+NP calculations.
- Let's look in more details.

2.2.3) Inclusive jets : 7 TeV with R=0.7

CMS-PAS-11-004





- Exp. Uncertainties – dominated by Jet Energy Scale.
- Theory uncertainty : low p_T - NP, middle/high p_T – scales, PDF.
- Promising data to constraint proton PDF within DGLAP evolution.



2.3) Dijet angular correlation


arXiv:1202.5535



- LO matrix element at LHC : dijet production.
- Anti- k_T with $R=0.5$ used.
- Each M_{jj} region normalised to 1.
- « Unfortunately » well described by pQCD in shape and no CI or resonances observed.

2.4) Inclusive jets : 7 TeV with R=0.5

Phys. Rev. Lett. 107
(2011) 132001





- The results for anti- k_T with $R=0.5$ are not so glorious:(Even if the difference is covered by uncertainties.
- Why ?

Where pQCD is challenged Small radius and/or forward jets







3.1) Dijet azimuthal decorrelation



- At LO 2 jets are back to back : $\Delta\phi \sim \pi$. True at large p_T .
- At low p_T ISR and FSR play a significant rôle. NLO+NP starts to fail to describe decorrelation. But LO+PS MC describe well.
- Simple evidence of importance of PS : large corrections beyond NLO!

3.2) POWHEG : NLO + PS + NP




- PS corrections are more important in forward direction and for smaller R (leaking out effect).
- Applying NP corrections from LO+PS MC to NLO is inconsistent when PS effects are large.


3.3) POWHEG vs NLO+NP

Phys. Rev. Lett. 107
(2011) 132001


S. Dooling, H. Jung



$R=0.5 \quad |y| < 0.5$



$R=0.5 \quad 2.0 < |y| < 2.5$





- POWHEG by itself describes better $R=0.5$ data than NLO+NP. PS represent part of the missing orders.
- Agreement not perfect at large y , but covered by systematics.

CMS Experiment at the LHC, CERN

3.4) Inclusive jets : Forward region



arXiv 1202.0704




- $3.2 < |\eta| < 4.7$
- JES dominating systematic.
- All models agree within systematics.

3.5) Inclusive jets : 7 TeV with R=0.7

FWD-PAS-10-003




- Access to very low x region. But sensitive to ISR there. To be understood.
- May be used to measure proton PDF ?



Pushing pQCD to the limits


large $|\Delta\eta|$



4.1) Inclusive to exclusive dijet ratio

arXiv 1204.0696



- Jets with $p_T > 35$ GeV and $|y| < 4.7$, $R = 0.5$.
- Measure ration of inclusive dijets events to exclusive dijet events : cancleration of systematics
- $|\Delta y| \sim |\Delta \eta| < 9$!!! between the two most external jets (Mueller-Navlett).
- Hope to see non DGLAP dynamics in the ladders between two jets.
- Strategic region for VBF physics : large $|\Delta y|$ and central jets veto !!!



4.2) Inclusive to exclusive dijet ratio

arXiv 1204.0696

- Ratio only described by PYTHIA (surprising ?). Influence of the tune and MPI small.
- Deviation of most of the other models at large $|\Delta y|$.
- Cascade, HEJ : include elements of CCFM or BFKL like dynamics.




Few worlds about Photons production



5.1) Single photons production




- Predictions from the NLO pQCD (JETPHOX)) agrees with Data except low p_T photons where NLO predictions tends to overestimated data.



5.2) Double photons production

JHEP01(2012)133

- Annihilation: $qq \rightarrow \gamma\gamma$
- Fusion: $gg \rightarrow \gamma\gamma$
- Fragmentation: $qg \rightarrow \gamma\gamma q$
- Calculation is done at NLO with DIPHOX, GAMMA2MC
- The overall agreement in diphoton mass spectrum.
The theoretical predictions underestimate the measured cross section for $\Delta\phi_{\gamma\gamma} < 2.8$.




SUMMARY

- 1) The understanding of QCD at LHC is a still challenging question even when considering simple observables as inclusive jets with low jet radius.
- 2) There are some QCD observables which are sufficiently well understood to use LHC data to constraint proton PDFs.
- 3) But for many others the NLO calculations are not sufficient in pp in contrary to ee, ep collisions. Parton Showers and other kind of leading log resummations seems to play an appreciable role.
- 4) This subject have to be understood since QCD is a background to non leptonic QCD searches and to the VBF production.
- 5) Photons program contrubuts/benefits but also suffers from the effort dedicated to the $H \rightarrow \gamma\gamma$.

BACKUP



4) Dijets : 1 central jet + 1 forward jet



- All models fails. The closest is HEJ