



Contribution ID: 94

Type: **Talk**

Open and closed heavy-flavour suppression in heavy-ion collisions with CMS

Tuesday 23 July 2013 11:30 (30 minutes)

The Compact Muon Solenoid (CMS) is fully equipped to measure quarkonia in the di-muon decay channel in the high multiplicity environment of nucleus-nucleus collisions. Quarkonia are especially relevant for studying the quark-gluon plasma since they are produced at early times and propagate through the medium, mapping its evolution. CMS has measured the nuclear modification factors of non-prompt J/psi (from b-hadron decays), prompt J/psi, inclusive psi(2S), and the first three Y states in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. A suppression of non-prompt J/psi, which is sensitive to the in-medium b-quark energy loss, has been measured at relatively high p_T ($6.5 < p_T < 30$ GeV/c) in PbPb collisions, compared to the yield in pp collisions scaled by the number of inelastic nucleon-nucleon collisions. For prompt J/psi in the same kinematic range, a strong, centrality-dependent suppression is observed. Such strong suppression at high p_T has previously not been observed at RHIC. At midrapidity and high p_T , psi(2S) show an even stronger suppression than J/psi. Furthermore, CMS has measured the suppression of the three Y states, separately, down to $p_T = 0$. A clear ordering of the suppression with binding energy is observed, as expected from sequential melting. The status of quarkonium measurements in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV will be also presented.

Primary authors: JO, Mihee (Korea University (KR)); CMS COLLABORATION, The (CMS)

Presenters: JO, Mihee (Korea University (KR)); CMS COLLABORATION, The (CMS)

Session Classification: Plenary 5