


$\Upsilon(nS)$ Polarization Results from CMS and CDF

Ilse Krätschmer*
(Hephy Vienna)
on behalf of the
CMS and CDF Collaborations

Beauty 2013
14th International Conference on B-Physics at Hadron Machines
8 - 14 April 2013




HEPHY
Institute of High Energy Physics




* supported by Austrian Science Fund (FWF): P24167

Introduction



Motivation

- Even after decades of theoretical and experimental research, quarkonium production is still not well understood
- Quarkonium polarization is sensitive to the production mechanism and therefore important to the theoretical understanding



NRQCD factorization

Braaten & Lee, Phys. Rev. D63, 071501(R) (2001)

DØ

DØ Collaboration, Phys. Rev. Letters 101, 182004 (2008)

CDF Run I

CDF Collaboration, Phys. Rev. Lett. 88, 161802 (2002)

k_T factorization

Baranov & Zotov, JETP Lett. 86, 435 (2007)

Quarkonium Polarization

- Polarization is measured through the average angular decay distribution - for vector particles most generally written as

$$\frac{dN}{d \cos \theta d\phi} \propto 1 + \lambda_{\theta} \cos^2 \theta + \lambda_{\phi} \sin^2 \theta \cos 2\phi + \lambda_{\theta\phi} \sin 2\theta \cos \phi$$

where λ_{θ} , λ_{ϕ} , $\lambda_{\theta\phi}$ are the polarization parameters

- Two extreme angular decay distributions:


Longitudinal polarization

$$J_z = 0$$

$$\lambda_{\theta} = -1$$

$$\lambda_{\phi} = 0$$

$$\lambda_{\theta\phi} = 0$$



Transverse polarization

$$J_z = \pm 1$$

$$\lambda_{\theta} = +1$$



$$\lambda_{\phi} = 0$$

$$\lambda_{\theta\phi} = 0$$




Definitions

- Angular decay distribution measured with respect to a certain reference frame:
 - center-of-mass helicity HX (polar axis $z_{HX} \approx$ direction of quarkonium momentum)
 - Collins-Soper CS ($z_{CS} \approx$ direction of relative velocity of colliding particles)
 - perpendicular helicity PX ($z_{PX} \perp z_{CS}$)
- Usage of the dimuon decay channel $\Upsilon(nS) \rightarrow \mu^+\mu^-$



Need to Measure Full Angular Distribution

- Only λ_θ in one reference frame was measured in the past
- The full angular decay distribution (three polarization parameters) should be measured: Two very different physical cases are indistinguishable if only λ_θ is measured.
- Observed polarization depends on the frame



$$\lambda_\theta = +1$$
$$\lambda_\phi = -1$$

$$\lambda_\theta = -1$$
$$\lambda_\phi = 0$$




$$\lambda_\theta = +1$$
$$\lambda_\phi = 0$$

Frame Independent Parameter

- Define frame invariant parameters such as $\tilde{\lambda}$ from the angular decay distribution of a given frame

$$\tilde{\lambda} = \frac{\lambda_{\theta} + 3\lambda_{\phi}}{1 - \lambda_{\phi}}$$




$$\tilde{\lambda} = +1$$



$$\tilde{\lambda} = -1$$

Eur. Phys. J. C 69 (2010) 657






CDF Run II $\Upsilon(nS)$ Polarization Analysis


- Previous results only measured λ_θ in the HX frame
- In this analysis, λ_θ , λ_ϕ and $\lambda_{\theta\phi}$ are determined simultaneously in two different reference frames (CS, HX) for $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ mesons - self-consistency of the results is tested with $\tilde{\lambda}$
- A dimuon sample collected in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV, corresponding to a total integrated luminosity of 6.7 fb^{-1} , is used
- Estimated number of signal events in the kinematic phase space under consideration ($p_T < 40$ GeV, $|y| < 0.6$):

$\Upsilon(1S)$	$\Upsilon(2S)$	$\Upsilon(3S)$
550k	150k	76k

Details in PRL 108, 151802 (2012): “Measurements of the Angular Distributions of Muons from Υ Decays in $p\bar{p}$ Collisions at $\sqrt{s}=1.96$ TeV”


$\Upsilon(nS)$ Polarization in the HX Frame, $|y| < 0.6$





- Results of the two reference frames are consistent
- No evidence of strong polarization over a wide range of p_T

- New CDF Run II result agrees with the previous CDF Run I result




CMS $\Upsilon(nS)$ Polarization Analysis

- λ_θ , λ_ϕ , $\lambda_{\theta\phi}$ and $\tilde{\lambda}$ are measured in three different reference frames (PX, CS, HX) for $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ mesons
- Analysis based on dimuon sample collected in pp collisions in 2011 at $\sqrt{s} = 7$ TeV, corresponding to a total integrated luminosity of 4.9 fb^{-1}
- Estimated number of signal events in the kinematic phase space under consideration ($p_T > 10$ GeV, $|y| < 1.2$):


$\Upsilon(1S)$	$\Upsilon(2S)$	$\Upsilon(3S)$
252k	94k	58k

- Analysis performed independently in five transverse momentum p_T bins for two dimuon rapidity $|y|$ cells


Details in PRL 110, 081802 (2013):
 “Measurement of the $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ polarizations in pp collisions at $\sqrt{s} = 7$ TeV”




$\Upsilon(nS)$ Polarization in the HX Frame, $|y| < 0.6$



$\Upsilon(nS)$ Polarization in the HX Frame, $0.6 < |y| < 1.2$




Frame Invariant Parameter $\tilde{\lambda}$



- Results of all three reference frames are consistent
- No evidence of unaccounted systematic uncertainties


CDF vs CMS



- Results from CDF and CMS are consistent
- CMS extends the measurements beyond the p_T and rapidity ranges probed by CDF
CDF: $p_T < 40 \text{ GeV}$, $|y| < 0.6$
CMS: $10 < p_T < 50 \text{ GeV}$, $|y| < 1.2$
- Measurements do not show strong polarizations: puzzling!

Comparison to Theory

- $\Upsilon(1S)$ suffers from large χ_b feed-down contribution, with unknown polarization
- $\Upsilon(3S)$ polarization calculated more reliably
- Theory predictions do not agree with experimental results



Summary and Conclusions



- CDF as well as CMS measured the polarization of $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ mesons
- Three frame dependent anisotropy parameters λ_θ , λ_ϕ , $\lambda_{\theta\phi}$ and the frame invariant parameter $\tilde{\lambda}$ have been measured: CDF determined these parameters in two reference frames (CS, HX) while CMS obtained results in three different frames (CS, HX, PX)
- CMS results ($10 < p_T < 50$ GeV, $|y| < 1.2$) extended the measurements beyond the p_T and rapidity ranges probed by CDF ($p_T < 40$ GeV, $|y| < 0.6$)
- No evidence of strong longitudinal or transverse polarizations has been observed




Other CMS Talks and Posters

- Torsten Dahms, "Y melting in CMS / ALICE", Tuesday, 9:30 - 9:50
- Marco de Mattia, " $B_s \rightarrow \mu^+\mu^-$ in CMS", Tuesday, 14:30-14:50
- Mauro Dinardo, "Measurement of $B \rightarrow K^{*0} \mu^+\mu^-$ in CMS", Tuesday, 16:35 - 16:55
- Luigi Guiducci, "ATLAS, CMS and LHCb Trigger/DAQ systems for Flavour Physics", Wednesday, 14:40 - 15:10
- Sara Fiorendi, "Heavy Flavour Spectroscopy and Production in CMS", Thursday, 11:35 - 11:55
- Alexis Pompili, "Observation of two structures in the $J/\Psi - \phi$ mass spectrum of exclusive $B^+ \rightarrow J/\Psi \phi K^+$ decays in CMS"
- Martino Dall'Osso, "Measurement of the X(3872) production cross section via decays to $J/\Psi \pi \pi$ "

BACKUP

CDF Detector




$\Upsilon(nS)$ Polarization in the CS Frame, $|y| < 0.6$


CDF Run II, 6.7 fb⁻¹

I stat.

I stat. + syst.





CMS Detector



Obtaining Polarization Parameters

Full and direct calculation of the Posterior Probability Distribution (PPD) of the polarization parameters λ_θ , λ_ϕ , $\lambda_{\theta\phi}$

1. Events distributed as in the background model are subtracted from the data sample
2. Definition of the PPD from the remaining signal-like events
3. Numerical results and graphical representations are determined from 1D and 2D projections of the PPD



$\Upsilon(1S)$
 $|y| < 0.6$
 $30 < p_T < 50$ GeV

Definition of the PPD


$$\mathcal{P}(\vec{\lambda}) \propto \prod_i \frac{1}{\mathcal{N}(\vec{\lambda})} W(\cos \theta^{(i)}, \phi^{(i)} | \vec{\lambda}) \varepsilon(p_1^{(i)}, p_2^{(i)})$$

\mathcal{N} : normalization

W : general angular distribution

ε : dimuon efficiency as a function of the muon momenta

Background Subtraction




- Signal region is defined as $\pm 1\sigma$ around mass peak
- Background fraction is determined by fits to the dimuon mass distribution
- Angular distribution of the background events are modeled as weighted sums of the distributions in the sidebands, left of Y(1S) and right of Y(3S) peak

- Event-by-event background subtraction of background-like events using a likelihood ratio criterion

Efficiencies


- Data-driven single muon efficiencies measured with the *Tag&Probe* method
- Precise knowledge of efficiencies needed to avoid introducing artificial polarization
- Dimuon efficiencies are calculated as the product of single muon efficiencies
- Correlations between muons are negligible as seen in detailed MC studies
- Efficiencies are accounted for on an event-by-event basis




Systematic Effects

- Sources of systematic effects:
 - Analysis method
 - Background model
 - Muon efficiencies
- Systematic uncertainties are propagated to the PPD
- Total uncertainties of the measurements are dominated by systematics at low p_T and statistics at high p_T
- $Y(2S)$ and $Y(3S)$ systematic uncertainties are dominated by the background model uncertainty, especially at low p_T


$\Upsilon(nS)$ Polarization in the PX Frame, $|y| < 0.6$



Y(nS) Polarization in the PX Frame, $0.6 < |y| < 1.2$



Y(nS) Polarization in the CS Frame, $|y| < 0.6$



$\Upsilon(nS)$ Polarization in the CS Frame, $0.6 < |y| < 1.2$

