






# RECENT UPC RESULTS FROM STAR

JANET SEGER (FOR THE STAR COLLABORATION)




# Ultrapерipheral Collisions

- Intense electromagnetic fields act for very short time
- Weizsacker-Williams model
  - Photon flux  $\sim Z^2$
- Photon emitted by projectile nucleus fluctuates into  $q\bar{q}$  pair, which then scatters from target nucleus
- Coherent coupling to both nuclei
  - small transverse momentum
    - Max photon energy  $\sim 3$  GeV at RHIC
- Exclusive vector meson production ( $\gamma+A \rightarrow V+A$ ) of interest as a probe of the nuclear gluon distribution  $g(x, Q^2)$  and GPDs



Klein & Nystrand, PRC60 014903  
 Baltz et al PRL89 012301 (2002)  
 Bauer et al NP A729 787 2003




# STAR Experiment




- TPC and TOF provide tracking and PID for  $\eta < 1.2$ 
  - TOF also used for triggering
- Beam-beam counters cover  $2 < \eta < 5$ 
  - Used as veto to define rapidity gap
- Zero degree calorimeters 18 m from interaction point
  - used for triggering

# STAR UPC triggers

- UPC\_Main
  - Low multiplicity in TOF
    - $2 < \# \text{ of TOF hits} < 6$
  - Veto on small-tile BBCs ( $3 < \eta < 5$ )
  - Signal in both ZDCs
    - $1 < \text{beam neutrons} < 6$  in each ZDC




- UPC\_Topo (small sample)
  - Low multiplicity in TOF
    - $2 < \# \text{ of TOF hits} < 6$
  - Veto on small-tile BBCs ( $3 < \eta < 5$ )
  - Back-to-back tracks in TPC, with veto for up-down tracks in TOF to reduce cosmic rays




# J/Ψ Photoproduction

- 2- gluon interaction
- Data Selection
  - $|Z_{\text{vertex}}| < 100 \text{ cm}$
  - # of tracks/ event =2
  - Connect vertex to TOF trigger to eliminate pile-up events
  - # of TPC hits  $\geq 14$
  - Pair  $p_T < 0.15 \text{ GeV}/c$
  - Vector meson rapidity:  $|y| < 1$
- Backgrounds:
  - Cosmic Rays
    - Rejected by ZDC trigger requirement
  - Beam gas interactions
    - Rejected with multiplicity, vertex cuts
  - Peripheral AA collisions
    - Reduced with multiplicity,  $p_T$  cuts




# 2010 Data ( $\sim 37\text{M}$ UPC triggers)

Pairs with  $3.0 < M_{ee} < 3.2 \text{ GeV}/c^2$



Peak at low  $p_T$   
consistent with  
coherent production

No PID  
Electron continuum not subtracted




$\sim 145$  candidates in  $J/\Psi$  peak

Chanaka De Silva

# J/ $\Psi$ rapidity distribution


- Cross section for photoproduction of J/ $\Psi$  at mid-rapidity can provide insight into nuclear gluon distribution ( $d\sigma/dy \sim [g(x, Q^2)]^2$ )



- Pairs with  $3.0 < M_{ee} < 3.2 \text{ GeV}/c^2$
- Not efficiency corrected

# J/ $\Psi$ t- distribution

- More statistics needed to observe diffraction pattern





Pairs with  
 $3.0 < M_{ee} < 3.2 \text{ GeV}/c^2$

Chanaka De Silva





# Rho diffraction

- $p_T$  of photon is very small
  - $p_T$  of  $\rho$  approx. balances the  $p_T$  of the recoiling gold
- Can expect to see diffraction pattern in  $t$ - distribution of coherently produced  $\rho$  mesons
- Fourier transform of  $d\sigma/dt$  is related to GPDs




# Rho meson photoproduction – 2010 data

- Data Selection
  - Exactly one neutron in each ZDC
  - Connect vertex to TOF trigger to eliminate pile-up events
  - Exactly two tracks out of selected vertex
    - Exclusive production
  - at least 14 hits in TPC
  - Select pions with TPC  $dE/dx$
- 2.82M  $\rho$  candidates



# $\rho$ t-distribution


- $t = -p_T^2$
- Like-sign background has been subtracted
- Data normalized using preliminary efficiencies & luminosity
- Incoherent tail is fit to a power law
  - Black line
- This is then subtracted to reveal the coherent distribution



Ramiro Debbé

ZEUS results for  $\gamma p \rightarrow \rho N$  scaled by 10.6 match the measured tail.

# Diffraction pattern




Sartre: Monte Carlo event generator based on an impact parameter dependent dipole model

**T. Ulrich and T. Toll**, arXiv:1108.1713 [nucl-th]  
<https://code.google.com/p/sartre-mc/>


Ramiro Debbe

- Sartre simulations give  $t$  of recoiling gold; data is measured  $t$  of  $\rho^0$ 
  - Data normalized using preliminary efficiencies & luminosity
- Can see diffraction pattern in data up through third dip
- Slope of first peak, and peak location are consistent with coherent interaction with an object with size of Au nucleus




Blue is Wood-Saxon with  $R_{\text{Au}} = 6.38$  fm,  $a = 0.53$  fm

# Details of diffraction pattern constrain dipole cross section models




Data normalized using preliminary efficiencies & luminosity

Ramiro Debbe



PhysRevC.87.024913


# Comparison to Starlight



Ramiro Debbe

- To match the locations of the dips, Starlight requires  $R_{\text{Au}} = 7.5$  fm
- Starlight allows photon transverse momentum  $\rightarrow$  no sharp valleys in the diffraction pattern

# Comparison to Starlight



- With gold radius of 6.38 fm, Starlight systematically shifted to higher  $p_T$  than data
  - Data normalized using preliminary efficiencies & luminosity


# Summary and Future Prospects

- $J/\Psi$  photoproduction in two units of rapidity around  $y = 0$  observed at RHIC
- Large sample of exclusive  $\rho$  photoproduction
  - Able to separate events where  $\rho$  scatters coherently off nucleus
  - Diffraction pattern of  $\rho$  mesons observed
- Corrections and normalizations need to be finalized
- U+U data at 193 GeV has been collected
- We are preparing for the 2015 p+Au run with the Roman Pots.



# Back up

# Model comparisons with $R_{Au} = 6.38$ fm



- With gold radius of 6.38 fm, models systematically shifted to higher  $p_T$  than data
  - Data normalized using preliminary efficiencies & luminosity