

Heavy Quarks in Heavy-Ion Collisions

Hendrik van Hees
with T. Lang, J. Steinheimer, M. Bleicher

Goethe University Frankfurt and FIAS


July 18, 2013



- 1 Heavy-quark interactions in the sQGP
 - Heavy quarks in heavy-ion collisions
 - Heavy-quark diffusion: The Langevin Equation
- 2 Non-perturbative HQ interactions
 - Resonance model for HQ-q Scattering
 - T-matrix approach with IQCD potentials
- 3 Comparison with data
 - Nonphotonic electrons at RHIC
 - D mesons at LHC
 - Predictions for D mesons at FAIR
 - Dileptons from correlated $D \bar{D}$ decays
- 4 Summary and Outlook


- Fast equilibration of hot and dense matter in heavy-ion collisions: collective flow (nearly ideal hydrodynamics) \Rightarrow sQGP
- Heavy quarks as calibrated probe of QGP properties
 - produced in early hard collisions: well-defined initial conditions
 - not fully equilibrated due to large masses
 - **heavy-quark diffusion** \Rightarrow probes for QGP-transport properties
- Langevin simulation within UrQMD-hydro hybrid model
- drag and diffusion coefficients
 - T -matrix approach with static lattice-QCD **heavy-quark potentials**
 - **resonance formation** close to T_c
 - mechanism for **non-perturbative strong interactions**

Heavy Quarks in Heavy-Ion collisions



hard production of HQs
described by PDF's + pQCD (PYTHIA)


c, b quark



HQ rescattering in QGP: Langevin simulation
drag and diffusion coefficients from
microscopic model for HQ interactions in the sQGP



Hadronization to D, B mesons via
quark coalescence + fragmentation



semileptonic decay \Rightarrow
"non-photonic" electron observables
 $R_{AA}^{e^+e^-}(p_T), v_2^{e^+e^-}(p_T)$

Relativistic Langevin process

- **Langevin process: friction force** + **Gaussian random force**
- in the (local) rest frame of the heat bath


$$d\vec{x} = \frac{\vec{p}}{E_p} dt,$$

$$d\vec{p} = -A\vec{p} dt + \sqrt{2dt}[\sqrt{B_0}P_\perp + \sqrt{B_1}P_\parallel]\vec{w}$$


- \vec{w} : normal-distributed random variable
- A : friction (drag) coefficient
- $B_{0,1}$: diffusion coefficients
- Einstein dissipation-fluctuation relation $B_1 = E_p T A$.
- flow via Lorentz boosts between “heat-bath frame” and “lab frame”
- A and B_0 from **microscopic models for qQ, gQ scattering**
- **background medium**: UrQMD \rightarrow hydro \rightarrow UrQMD

Non-perturbative interactions: Resonance Scattering

- General idea: Survival of D - and B -meson like **resonances** above T_c
- model based on chiral symmetry (light quarks) HQ-effective theory
- **elastic heavy-light-(anti-)quark scattering**




- D - and B -meson like resonances in **s**QGP



- parameters
 - $m_D = 2 \text{ GeV}, \Gamma_D = 0.4 \dots 0.75 \text{ GeV}$
 - $m_B = 5 \text{ GeV}, \Gamma_B = 0.4 \dots 0.75 \text{ GeV}$

T-matrix


- Brueckner many-body approach for elastic $Qq, Q\bar{q}$ scattering



- V : static $q\bar{q}$ potential from lattice QCD (F and U)
- reduction scheme: 4D Bethe-Salpeter \rightarrow 3D Lipmann-Schwinger
- S - and P waves
- Relation to invariant **matrix elements**

$$\sum |\mathcal{M}(s)|^2 \propto \sum_q d_a \left(|T_{a,l=0}(s)|^2 + 3|T_{a,l=1}(s)|^2 \cos \theta_{\text{cm}} \right)$$


T-matrix results




- **resonance formation** at lower temperatures $T \simeq T_c$
- melting of resonances at higher T
- model-independent assessment of elastic $Qq, Q\bar{q}$ scattering!

Nonphotonic electrons at RHIC


- form D and B mesons via **quark-antiquark coalescence**
- use PYTHIA for semi-leptonic decays
- comparison to single-electron data from PHENIX (200 AGeV Au-Au collisions)




[T. Lang, Hvh, J. Steinheimer, M. Bleicher, arXiv: 1211.6912 [hep-ph]]



- form D via **quark-antiquark coalescence**
- comparison to D-meson data from ALICE (2.76 ATeV Pb-Pb collisions)




[T. Lang, Hvh, J. Steinheimer, M. Bleicher, arXiv: 1211.6912 [hep-ph]]




D mesons at FAIR

- form D via **quark-antiquark coalescence**
- large sensitivity to initial HQ distributions (use estimates from HSD and PYTHIA)





[T. Lang, HvH, J. Steinheimer, M. Bleicher, arXiv: 1305.1797 [hep-ph]]



D mesons at FAIR



- form D via **quark-antiquark coalescence**
- large sensitivity to initial HQ distributions
(use estimates from HSD and PYTHIA)



[T. Lang, HvH, J. Steinheimer, M. Bleicher, arXiv: 1305.1797 [hep-ph]]

D mesons at FAIR


- form D via **quark-antiquark coalescence**
- large sensitivity to initial HQ distributions (use estimates from HSD and PYTHIA)
- large μ_B in resonance model: \bar{c} more dragged than c



[T. Lang, HvH, J. Steinheimer, M. Bleicher, arXiv: 1305.1797 [hep-ph]]

Dileptons from correlated $D\bar{D}$ decays

- for $m_\phi \lesssim M_{\ell+\ell^-} \lesssim m_{J/\psi}$:
dilepton emission from thermal QGP and from correlated $D\bar{D}$ decays
- medium modifications of D and \bar{D} destroy correlations



[T. Lang, HvH, J. Steinheimer, M. Bleicher, arXiv: 1305.7377 [hep-ph]]

Summary and Outlook

- Heavy quarks in the sQGP
- non-perturbative interactions
 - mechanism for strong coupling: resonance formation at $T \gtrsim T_c$
 - lattice-QCD potentials parameter free
 - also provides “natural” mechanism for quark coalescence
- Comparison to data and predictions for FAIR
 - R_{AA} and v_2 of non-photonic electrons at RHIC
 - R_{AA} and v_2 for D mesons at LHC
 - R_{AA} and v_2 for D mesons at FAIR (pp baseline mandatory!)
 - impact of medium modifications on correlated $D \bar{D}$ decays to dileptons
- Outlook
 - implementation of hadronic cross sections for D/B-meson diffusion
 - include inelastic heavy-quark processes (gluo-radiative processes)
 - implement resonance-recombination model for hadronization
 - charmonium/bottomonium suppression/regeneration