

OBSERVATION OF D^0 MIXING AT CDF

Jonathan Lewis

Fermilab

18 July 2013

Flavor mixing in the charm sector

- Neutral mesons can oscillate between matter and anti-matter

- Mass eigenstates \neq flavor eigenstates

$$|D^0\rangle = (|D_1\rangle + |D_2\rangle)/\sqrt{2} \quad |\bar{D}^0\rangle = (|D_1\rangle - |D_2\rangle)/\sqrt{2}$$

- Time evolution



$$|D_{1,2}(t)\rangle = |D_{1,2}(0)\rangle e^{-t(\Gamma_{1,2} + iM_{1,2})}$$

$$x = \frac{m_1 - m_2}{\Gamma_D} = 2 \frac{m_1 - m_2}{\Gamma_1 + \Gamma_2} \quad y = \frac{\Gamma_1 - \Gamma_2}{2\Gamma_D} = \frac{\Gamma_1 - \Gamma_2}{\Gamma_1 + \Gamma_2}$$

- Mixing well established in K^0 (1962), B^0 (1987), and B_s^0 (CDF 2006) decays
 - Evidence for D^0 from Belle (2006), Babar and CDF(2007)
 - Observation by LHCb (2012)

Mixing Characteristics

- ▣ Two sources
 - Long-range intermediate states
 - ▣ Dominant
 - ▣ Large theoretical uncertainties
 - Short-range box diagram
 - ▣ Suppressed
 - No GIM suppression from heavy top quark
 - ▣ Possible enhancement by New Physics
- ▣ Charm mixing slow
 - $x, y \ll 1$



Mixing Signature

- ▣ Compare rate of wrong-sign $D^0 \rightarrow K^+ \pi^-$ decays to right-sign $D^0 \rightarrow K^- \pi^+$ decays
 - Tag flavor at production with $D^{*+} \rightarrow D^0 \pi^+$ decays
 - Wrong sign events can come from mixing or doubly Cabibbo-suppressed (DCS) decays
 - ▣ DCS decays are time independent



- ▣ Time evolution

$$R(t) = R_D + \sqrt{R_D} y' t + \frac{x'^2 + y'^2}{4} t^2$$

$$x' = x \cos \delta_{K\pi} + y \sin \delta_{K\pi}$$

$$y' = y \cos \delta_{K\pi} - x \sin \delta_{K\pi}$$

$\delta_{K\pi}$: strong phase difference
between CF and DCS amplitudes

Data Sample

- ▣ Full CDF II data set
 - $9.6 \text{ pb}^{-1} \text{ p}\bar{\text{p}}$ at $\sqrt{s}=1960 \text{ GeV}$
- ▣ Heavy flavor hadronic decay trigger
 - Two opposite-charge tracks with $p_{\text{T}} > 2 \text{ GeV}/c$, $d_0 > 100 \mu\text{m}$


1.4T Solenoid

COT Drift Chamber

- $\delta p_{\text{T}}/p_{\text{T}}^2 \approx 0.07\%$
- PID from dE/dx with $\sim 1.5\sigma$ K- π separation at $p_{\text{T}}=2 \text{ GeV}$



SVX Vertex Detector

- $\delta d_0 \approx 25 \mu\text{m}$
- Beam spot rms $25 \mu\text{m}$




Reconstruction Method

- Reconstruct events with both RS and WS hypotheses







Large background to WS decays from incorrect charge assignment in RS decays





Reduce misassignment background

- Remove WS candidates consistent with RS hypothesis and vice versa
 - Removes 96% of background
 - 78% efficient for signal



D* Reconstruction


- ▣ Standard method
 - ▣ Fit $\Delta m = m(K\pi\pi_{\text{tag}}) - m(K\pi) - m_{\pi}$
- ▣ Must account for D^0 and D^* backgrounds
 - ▣ Cutting on D^0 peak would include fake D mesons in D^* fit
 - ▣ In 10 bins of D^0 decay time, for RS and WS candidates, find the D^0 yield in 60 bins of Δm
 - ▣ Fit the resulting Δm distribution to get the D^* yields



Wrong-Sign Ratio

- From D^* yields, get measured ratio in each time bin

$$R_i = \frac{N_i(D^{*+} \rightarrow [K^+ \pi^-] \pi^+) + N_i(D^{*-} \rightarrow [K^- \pi^+] \pi^-)}{N_i(D^{*+} \rightarrow [K^- \pi^+] \pi^+) + N_i(D^{*-} \rightarrow [K^+ \pi^-] \pi^-)}$$





B Contamination


- ▣ Must account for B decay component
 - D^0 will have larger apparent lifetime
 - Mixing sample is required to be consistent with prompt production, but some B contamination remains

$$R_m(t) = \frac{N^{WS}(t) + N_B^{WS}(t)}{N^{RS}(t) + N_B^{RS}(t)}$$


B Fraction Measurement



- Use impact parameter of reconstructed D^0 mesons



B Fraction



- ▣ Measured in RS data
- ▣ Fit to 4th order polynomial

Mixing Fit

- χ^2 fit vs. decay time
 - Correct for B component using MC for D decay time distribution as a function of apparent decay time

$$\chi^2 = \sum_{i=1}^{20} \left[\frac{r_i - R_m(t_i)}{\sigma_i} \right]^2 + C_{f_B}(\mathbf{p}) + C_{R_B}(\mathbf{h})$$

20 measured WS/RS points r_i with error σ_i

Gaussian constraint on f_B parameters (\mathbf{p})

Gaussian constraint on MC decay time distributions of D^* from B

$$R_m(t) = \frac{N^{WS}(t) + N_B^{WS}(t)}{N^{RS}(t) + N_B^{RS}(t)} = R(t) \left[1 + f_B^{RS}(t) \left(\frac{R_B(t)}{R(t)} - 1 \right) \right]$$



$$f_B^{RS}(t) = \frac{N_B^{RS}(t)}{N^{RS}(t) + N_B^{RS}(t)}$$

Fraction of RS D^* from B decays

$$R_B(t) = \frac{N_B^{WS}(t)}{N_B^{RS}(t)}$$

WS/RS ratio of non-prompt D^0
 Calculated by weighting $R(t)$ with the decay-time distribution of secondary D^0 from MC


Mixing Fit Results



Fit type	χ^2/ndf	Parameter	Fitted values $\times 10^{-3}$	Correlation coefficient		
				R_D	y'	x'^2
Mixing	16.91/17	R_D	3.51 ± 0.35	1	-0.967	0.900
		y'	4.3 ± 4.3		1	-0.975
		x'^2	0.08 ± 0.18			1
No-mixing	58.75/19	R_B	4.30 ± 0.06			

Significance

- ▣ Bayesian probability contour
 - No mixing hypothesis excluded at 6.1σ
- ▣ Frequentist method using toy-MC in samples without
 - $\Delta\chi^2$ difference exceeds observed in 6 of 10^{10} trials
 - No-mixing case excluded at 6.1σ



Summary

- CDF confirms observation of D^0 mixing at 6.1σ
 - Parameters similar to other experiments
- Nature is giving up her charms, but without indication of new physics

