

Recent H1 results on Heavy Flavour Photoproduction

Michel Sauter

Ruprecht-Karls-Universität Heidelberg
for the H1Collaboration



EPS 2013

The 2013 European Physical Society Conference on High Energy Physics
18-24 July 2013, Stockholm, Sweden




The HERA ep collider (1992 – 2007) at DESY in Hamburg

- ep collider:
- e^\pm energy: 27.6 GeV
- p energy: 920 GeV
- Center of mass energy: 318 GeV
- 2 collider experiments: H1 and ZEUS





Motivation to measure heavy flavour production

- Charm and beauty quarks at HERA are mainly produced in Photon-Gluon-Fusion → sensitive to the gluon in the proton.
- Hard scales for perturbative QCD:
 - $m_{c,b}^2, p_T^2, Q^2$
 - multi-scale problem, for example combined to $\mu_0 = \frac{1}{2} \sqrt{(m_{b,c}^2 + p_T^2)}$
- Interpretation of heavy flavour measurements:
 - Use the pQCD calculations and **constrain the gluon density of the proton.**
 - Take the gluon density from elsewhere and **test the consistency of the pQCD calculation.**



Two kinematic regimes:

- Photoproduction: $Q^2 \approx 0 \text{ GeV}^2$
- Deep Inelastic Scattering: $Q^2 > 1 \text{ GeV}^2$
(scattered electron detected)



QCD scheme:


- Massive scheme – Fixed Flavour Number Scheme (FFNS):
 - c and b quarks generated dynamically via boson-gluon-fusion.
 - c and b quarks treated massive.
 - Expected to be valid for **small scales** $\mu^2 \approx m_{b,c}^2$

QCD predictions:

- QCD LO + Parton shower Monte Carlo generators:
 - Collinear factorization, DGLAP evolution (PYTHIA).
 - k_T factorization, CCFM evolution (CASCADE).
 - Used for data corrections and model comparisons.
- QCD NLO calculations:
 - Massive scheme, NLO(α_s^2):
 - FMNR
 - MC@NLO
 - Used for comparisons and extrapolations to full phase space of beauty production.

Tagging methods for heavy flavours at HERA

- Rates at HERA behaved like $\sigma(b) : \sigma(c) : \sigma(uds) \approx 1 : 50 : 2000$
- Charm and beauty enrichment is possible with:
 - 1) Full reconstruction
 - Only possible for charm at HERA, eg. $D^* \rightarrow K\pi\pi$.
 - 2) Lepton tagging: Use semileptonic b/c decay channels
 - look for μ or e , high BR(c,b \rightarrow lepton + anything)
 - 3) p_T^{rel} tagging : b/c quark have large masses
 - look for decay leptons with a high transverse momentum w.r.t the b quark flight direction.
 - 4) Lifetime tagging: b/c quark have long lifetimes:
 - look for displaced vertices.
 - look for tracks with large impact parameters δ .
 - 5) Secondary vertex mass tagging: long lifetime and large masses
 - look for high secondary vertex masses.



Methods used in the analyses discussed today:

- **D* analysis:** D* reconstruction.
- **Low p_T analysis:** 2 electrons.
- **Muon analysis:** 1 μ for p_T^{rel} tag, lifetime with large impact parameters δ .







Data sample: $\mathcal{L}=93 \text{ pb}^{-1}$

Phase Space

$$Q^2 < 2 \text{ GeV}^2, p_T^{D^*} > 1.8 \text{ GeV}$$

Charm tagging

D^* meson reconstruction via:



- Very high precision of the data, compared to the uncertainties of the NLO predictions.
- NLO predicted shapes less sensitive to theoretical uncertainties, generally show a reasonable agreement with the data.




Data sample: $\mathcal{L}=93 \text{ pb}^{-1}$


Phase Space

$$Q^2 < 2\text{GeV}^2, \quad p_T^{D^*} > 2.1\text{GeV}$$

2 jets with: $p_T^{\text{jet } 1} > 3.5 \text{ GeV}$




resolved enhanced





direct enhanced

- Azimuthal correlation between the two jets, $\Delta\Phi$:



- Fraction of the photon energy entering the hard interaction (direct vs resolved), x_γ^{obs} :

$$x_\gamma^{\text{obs}} = \frac{\sum_{\text{Jet}1}(E - p_z) + \sum_{\text{Jet}2}(E - p_z)}{\sum_h(E - p_z)}$$



- MC@NLO predictions below the data for resolved photons, direct contribution reasonably well-described in normalization, shape not well described.



Focus of the measurement:


- Low $p_T(b)$ momentum
- Low experimental thresholds
- Extraction of the b-cross section from two low p_T -electrons, $p_T(e) \geq 1 \text{ GeV}$

Trigger for low p_T -Electrons:

Combination of **calorimeter** (Jet Trigger) and **tracker** (Fast Track Trigger):

- Topological match
- Cut on $E_{T,JT}/p_{T,FTT}$


Data sample: $\mathcal{L}=48 \text{ pb}^{-1}$



Offline Electron Identification:

Combination of:

- **calorimeter shower profiles**
- **dE/dx measured in the tracker**
- $\sim 90\%$ efficiency at a rejection of $\sim 99\%$




A. W. Jung et al., Proc. 15th IEEE-NPSS Real-Time Conference, (2007) 1.

B. Olivier et al. Nucl. Instrum. Meth. A 641 (2011) 58.



Heavy Flavour Tagging


- Exploit di-electron correlations:
 - Invariant mass m_{ee}
 - Azimuthal angle $\Delta\phi_{ee}$
 - Charge product $q(e1)*q(e2)$






Heavy Flavour Tagging

- Exploit di-electron correlations:
 - Invariant mass m_{ee}
 - Azimuthal angle $\Delta\phi_{ee}$
 - Charge product $q(e1) \cdot q(e2)$
- An additional background region (open electron identification cuts) constrains uds.
- Matrix unfolding of the differential beauty cross section (similar to 2d template fit).




$J/\psi \rightarrow e^+e^-$ mass peak





- Differential beauty cross section as function of the mean b quark momentum.

- Access to lowest $p_T(b)$ values ever measured in ep.
- Data in agreement with the NLO calculation, but slightly below.





Data sample: $\mathcal{L}=179 \text{ pb}^{-1}$




Phase Space

Events with:

- 1 muon, $p_T^\mu > 2.5 \text{ GeV}$
- 2 jets, $p_T^{\text{jet } 1(2)} > 7 \text{ (6) GeV}$





Heavy Flavour Tagging

- Momentum relative to μ jet, p_T^{rel} .
- Impact parameter δ .
- 2d template fit.






- Simultaneous measurement of differential charm and beauty jet cross sections.
- Azimuthal angle difference of jets $\Delta\phi^{jj} \rightarrow$ sensitive to higher orders.



- The data are in agreement with NLO calculation (MC@NLO).
- $\Delta\phi^{jj}$ described by all models (LO, NLO).

HERA



- Many measurements agreeing with each other over a wide $p_T(b)$ range.
- General NLO calculation (FMNR) consistent with data.

- Heavy flavour production at HERA allows to test QCD at different scales.
- New heavy flavour photoproduction measurements of H1 using different experimental techniques and having different systematics are in good agreement.
- The data is in general in agreement with NLO pQCD predictions.



- “Measurement of Inclusive and Dijet D^* Meson Cross Sections in Photoproduction at HERA”
[DESY-11-248, H1 Collab., F.D. Aaron et al., Eur. Phys. J. C72 \(2012\) 1995](#)
- “Measurement of Beauty and Charm Photoproduction using Semi-muonic Decays in Dijet Events at HERA”
[DESY-12-059, H1 Collab., F.D. Aaron et al., Eur. Phys. J. C72 \(2012\) 2047](#)
- “Measurement of Beauty Photoproduction near Threshold using Di-electron Events with the H1 Detector at HERA”
[DESY-12-072, H1 Collab., F.D. Aaron et al., Eur. Phys. J. C72 \(2012\) 2148](#)



Heavy Flavour Tagging

- An additional background region (open electron identification cuts) constrains uds.
- Electrons are identified by
 - Electron discriminator, D_{ele}
 - Isolation criterion, $R_{E,cone}$
- Definition of the background region:

$R_{E,cone}^{\max(e1,e2)}$	$D_{ele}^{\min(e1,e2)}$	
	0.825 – 0.875	0.875 – 1.0
150 – 350%	B1	B3
0 – 150%	B2	S

Open cut

Background templates

