




CMS Experiment at LHC, CERN
Data recorded: Mon Nov 8 11:30:53 2010 CEST
Run/Event: 150431 / 630470
Lumi section: 173

Two and Four Particle Correlations in pPb Collisions with CMS

Monika Sharma
For the CMS Collaboration




July 18, 2013

Monika Sharma at EPS HEP-2013, Stockholm, Sweden



Motivation: hydrodynamics

Azimuthal particle correlations in AA collisions



Elliptic, v_2



Triangular, v_3

Quantified by coefficients of a Fourier series

Initial spatial anisotropy




Final momentum anisotropy



$$\text{Fourier series: } dN/d\phi = 1 + 2v_1 \cos(\phi) + 2v_2 \cos(2\phi) + 2v_3 \cos(3\phi) + \dots$$



Motivation: establishing baseline of PbPb collisions

Controlled environment? pPb collisions



- ✓ No deconfined medium is expected to be formed?
- ✓ What about azimuthal anisotropy in pPb collisions?



Motivation



- Long range correlations can be explained by flow harmonics in PbPb
- Are these correlations in pPb also related to hydrodynamic flow?
- Or CGC?

Data sets, trigger selection and multiplicity distribution

- **Data sets**
 - 2013 pPb + Pbp, 31 nb^{-1}
 - 2011 PbPb, $2.3 \mu\text{b}^{-1}$ (50-100%)
- **Triggers**
 - Minimum bias trigger
 - High multiplicity triggers in 2013




Track ($p_T > 0.4 \text{ GeV}/c$, $|\eta| < 2.4$)
multiplicity distribution in pPb
for different triggers


The fraction of events in $300 \leq N_{\text{trk}}^{\text{offline}} < 350$ is
about 10^{-7} with respect to all MinBias events.

Data sets, trigger selection and multiplicity distribution

- **Data sets**
 - 2013 pPb + Pbp, 31 nb^{-1}
 - 2011 PbPb, $2.3 \mu\text{b}^{-1}$ (50-100%)
- **Triggers**
 - Minimum bias trigger
 - High multiplicity triggers in 2013



Track multiplicity distribution in
MinBias pPb and PbPb 50-100%




Results: Two-particle correlations in PbPb and pPb

PbPb

(a) CMS PbPb $\sqrt{s_{NN}} = 2.76$ TeV, $220 \leq N_{\text{trk}}^{\text{offline}} < 260$

$1 < p_{\text{T}}^{\text{trig}} < 3$ GeV/c

$1 < p_{\text{T}}^{\text{assoc}} < 3$ GeV/c




pPb

(b) CMS pPb $\sqrt{s_{NN}} = 5.02$ TeV, $220 \leq N_{\text{trk}}^{\text{offline}} < 260$

$1 < p_{\text{T}}^{\text{trig}} < 3$ GeV/c

$1 < p_{\text{T}}^{\text{assoc}} < 3$ GeV/c




PLB [724 \(2013\) 213](#)

- PbPb and pPb use the same multiplicity selection, $220 \leq N_{\text{trk}}^{\text{offline}} < 260$
- Very strong long-range correlations in pPb

Discovery: pPb pilot run

'Near Side long range correlations'

→ Long range rapidity correlations
→ Seen only for high multiplicity
pp and pPb collisions




[PLB 718 \(2013\) 795](#)

- Explore in detail the multiplicity and p_T dependence of the 2-particle correlations
- New observable: 4-particle correlations add greater sensitivity to collective effects

Discovery: pPb pilot run

'Near Side long range correlations'

→ Long range rapidity correlations
→ Seen only for high multiplicity
pp and pPb collisions



'Away Side JET', $\Delta\phi \sim \pi$

'Near Side JET', $\Delta\phi \sim 0$

What is new from the latest pPb run?


[PLB 718 \(2013\) 795](#)

- Explore in detail the multiplicity and p_T dependence of the 2-particle correlations
- New observable: 4-particle correlations add greater sensitivity to collective effects

2-particle correlation method


CMS pPb $\sqrt{s} = 5.02$ TeV, $N \geq 110$

$1 < p_T^{\text{trig}} < 2$ GeV/c
 $1 < p_T^{\text{assoc}} < 2$ GeV/c



Average over
 ridge region
 $(2 < |\Delta\eta| < 4)$

CMS pPb $\sqrt{s} = 5.02$ TeV, $N \geq 110$



Shift the distribution to zero yield at minimum (ZYAM)
 and extract the associated yield

Physics Letters B
 718 (2013) 795–814


$$v_n \{2, |\Delta\eta| > 2\} (p_T) = \frac{V_{n\Delta}(p_T, p_T^{\text{ref}})}{\sqrt{V_{n\Delta}(p_T^{\text{ref}}, p_T^{\text{ref}})}}$$

Fourier decomposition:



$$\frac{dN^{\text{pair}}}{d\Delta\phi} \sim 1 + 2 \sum_{n=1} V_{n\Delta} \cos(n\Delta\phi)$$

Assuming factorization:

$$V_{n\Delta} = v_n(p_T^{\text{trig}}) \times v_n(p_T^{\text{assoc}})$$



Multi-particle correlations



Four particle correlations (Q-cumulant method): remove 2 and 3 particle correlations


$$\begin{array}{c} \varphi_1 \\ \bullet \\ \varphi_2 \\ \bullet \end{array} \quad \begin{array}{c} \varphi_3 \\ \bullet \\ \varphi_4 \\ \bullet \end{array} = \begin{array}{c} \bullet \quad \bullet \\ \bullet \quad \bullet \end{array} + \begin{array}{c} \bullet \quad \bullet \\ \bullet \quad \bullet \end{array} + \begin{array}{c} \bullet \quad \bullet \\ \bullet \quad \bullet \end{array} \longrightarrow c_n\{4\} = \langle\langle 4 \rangle\rangle - 2 \cdot \langle\langle 2 \rangle\rangle^2$$

$$\langle e^{in(\varphi_1 + \varphi_2 - \varphi_3 - \varphi_4)} \rangle - \langle e^{in(\varphi_1 - \varphi_3)} \rangle \langle e^{in(\varphi_2 - \varphi_4)} \rangle - \langle e^{in(\varphi_1 - \varphi_4)} \rangle \langle e^{in(\varphi_2 - \varphi_3)} \rangle$$

Associated yield


Yield vs p_T

- Similar p_T dependence for PbPb and pPb
- Larger in PbPb ($|\Delta\eta|>2$)
- Expected behavior due to jets (Fig. b)




Yield vs multiplicity

- Yield becomes significant at $N \sim 40-50$, followed by a monotonic rise
- Larger in PbPb ($|\Delta\eta|>2$)



v_2 in PbPb and pPb

PLB 724 (2013) 213



- v_2 shows similar shape in pPb and PbPb, but is smaller in pPb
- hydro calculation agrees with $v_2\{4\}$



July 18, 2013


Monika Sharma at EPS HEP-2013, Stockholm, Sweden

13




v_3 in PbPb and pPb

PLB [724 \(2013\) 213](#)



- v_3 shows similar shape in pPb and PbPb; magnitude comparable
- Hydro prediction: $v_3\{PP\}$, not including fluctuations



July 18, 2013


Monika Sharma at EPS HEP-2013, Stockholm, Sweden

14



Multiplicity dependence of v_2


PLB [724 \(2013\) 213](#)



- v_2 is smaller in pPb than in PbPb; turns on at $N \sim 40-50$

Multiplicity dependence of v_3

PLB [724 \(2013\) 213](#)




- v_3 is essentially the same in pPb and PbPb; turns on at $N \sim 40-50$

Summary

Comparison of high statistics, high multiplicity pPb and PbPb data as a function of p_T and multiplicity

- Large $v_2\{4\}$ and $v_3\{2\}$ in pPb
- Associated yield, $v_2\{4\}$ and $v_3\{2\}$ become apparent at about the same multiplicity; $N \sim 50$
- $v_3\{2\}$ is essentially the same in pPb and PbPb at the same multiplicity



Back-up



July 18, 2013

Monika Sharma at EPS HEP-2013, Stockholm, Sweden

18



Peripheral subtraction


- Away-side correlations contain non-flow effects
- Subtract the data for high multiplicity by low multiplicity to correct for this

Fourier decomposition:

$$\frac{1}{N_{\text{trig}}} \frac{dN^{\text{pair}}}{d\Delta\phi} = \frac{N_{\text{assoc}}}{2\pi} \left\{ 1 + \sum_n 2V_{n\Delta} \cos(n\Delta\phi) \right\}$$

Subtracting peripheral correlations in v_2, v_3 calculations:


$$V_{n\Delta}(\text{cent}) - V_{n\Delta}(\text{peri}) \times \frac{N_{\text{assoc}}(\text{peri})}{N_{\text{assoc}}(\text{cent})} \times \frac{Y^{\text{jet}}(\text{cent})}{Y^{\text{jet}}(\text{peri})}$$



Subtract $N_{\text{trk}}^{\text{offline}} < 20$ (70-100%) to avoid removing signal ($N_{\text{trk}}^{\text{offline}} \sim 40$)

Account for the fact that jet correlation increases with multiplicity

Test our procedure in HIJING




Weighted by near-side jet yield, most of non-flow correlations are subtracted

PbPb vs pPb: p_T dependence

PbPb

$|\Delta\eta| > 2$


$|\Delta\eta| < 1$



v_2 in PbPb and pPb

PbPb

pPb



- v_2 is smaller in pPb than in PbPb
- Peripheral subtraction has a small effect at high multiplicity

$$v_2\{2\} = \sqrt{\langle v_2 \rangle^2 + \sigma_{v_2}^2} \quad v_2\{4\} = \sqrt{\langle v_2 \rangle^2 - \sigma_{v_2}^2} \quad \frac{\sigma_{v_2}}{v_2} = \sqrt{\frac{v_2^2\{2\} - v_2^2\{4\}}{v_2^2\{2\} + v_2^2\{4\}}}$$



- ❖ Hydro-flow is not incorporated in the HIJING MC model – $c_2\{4\}$ consistent with zero for small bin width (2 or 5), while becomes nonzero for big bin width (30)
- ❖ The effect becomes larger going to more peripheral collisions
- ❖ In pPb data, $c_2\{4\}$ crosses zero and becomes negative at certain multiplicity. This is an indication of the onset of multi-particle correlation effect
- ❖ A bin width of 5 is chosen for the $v_2\{4\}$ analysis

<https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsHIN13002>

