

# Development of Liquid Scintillator containing Zirconium Complex for Neutrinoless Double Beta Decay Experiment

The 2013 European Physics Society Conference on High Energy Physics

Stockholm, Sweden 18 – 24 July, 2013


Y.Fukuda<sup>a</sup> T.Muramatsu<sup>a</sup> S.Moriyama<sup>b</sup>, Narengerile<sup>a</sup>, A.Obata<sup>a</sup>, I.Ogawa<sup>c</sup>, A.Gunji<sup>\*d</sup>

<sup>a</sup>Miyagi University of Education <sup>b</sup>Kamioka Observatory ICRR, University of Tokyo <sup>c</sup>University of Fukui <sup>d</sup>Tokyo University of Science (special thanks)

## 1. Neutrinoless Double Beta Decay

### ◆ Neutrinoless double beta decay

- Lifetime and neutrino mass  
 $[T_{1/2}^{0\nu}(0^+ \rightarrow 0^+)]^{-1} = G_{0\nu}(E_{0\nu}, Z) |M_{0\nu}|^2 < m_{\nu} >^2$
- Energy spectrum and lifetime measurement  
 • monochromatic energy = Q-value  
 •  $T_{1/2} \sim a(M/\Delta E)^3$  a: abundance M: mass t: meas.time  $\Delta E$ : energy res. B: BG rate



**Requirement : Low background rate, Large target mass and High energy resolution.**


### ◆ Double beta decay candidates

$\beta\beta$ emitters with $Q_{\beta\beta} > 2$ Mev		
Transition	$Q_{\beta\beta}$ (keV)	Abundance (%) ( $^{232}\text{Th} = 100$ )
$^{110}\text{Pd} \rightarrow ^{110}\text{Cd}$	2013	12
$^{76}\text{Ge} \rightarrow ^{76}\text{Se}$	2040	8
$^{124}\text{Sn} \rightarrow ^{124}\text{Te}$	2288	6
$^{136}\text{Xe} \rightarrow ^{136}\text{Ba}$	2479	9
$^{130}\text{Te} \rightarrow ^{130}\text{Xe}$	2533	34
$^{116}\text{Cd} \rightarrow ^{116}\text{Sn}$	2802	7
$^{82}\text{Se} \rightarrow ^{82}\text{Kr}$	2995	9
$^{100}\text{Mo} \rightarrow ^{100}\text{Ru}$	3034	10
$^{96}\text{Zr} \rightarrow ^{96}\text{Mo}$	3350	3
$^{150}\text{Nd} \rightarrow ^{150}\text{Sm}$	3667	6
$^{48}\text{Ca} \rightarrow ^{48}\text{Ti}$	4271	0.2

- above  $^{208}\text{Tl}$   $\gamma$  line (2.614MeV) :  $^{48}\text{Ca}$ ,  $^{150}\text{Nd}$ ,  $^{96}\text{Zr}$ ,  $^{100}\text{Mo}$ ,  $^{82}\text{Se}$ ...
- large abundance :  $^{100}\text{Mo}$ ,  $^{82}\text{Se}$ ,  $^{150}\text{Nd}$ ,  $^{96}\text{Zr}$
- solved in liquid scintillator formed as metal complex

**Zirconium ( $^{96}\text{Zr}$ ) is possible candidate.**


### ◆ Experimental limits for neutrino mass



- high energy resolution :  $4\% @ 2.5\text{MeV} = 100\text{keV}$
- low background rate :  $0.01\text{count kg}^{-1}\text{y}^{-1}$
- large target mass :  $\sim \text{ton scale}$

**Goal:  $< m_{\nu} > \sim 10\text{meV}$**

### ◆ Detector design for Zr loaded liquid scintillator




**Assuming 10w.t.% solubility**

- high energy resolution :  $4\% @ 2.5\text{MeV} = 100\text{keV}$
- low background rate :  $0.01\text{count kg}^{-1}\text{y}^{-1}$
- large target mass :  $\sim \text{ton scale}$

**Zirconium Complex in Organic liquid Scintillator (ZICOS) experiment**

## 2. Zirconium complex


### ◆ Zirconium (IV) acetylacetonate



- good solubility (over 10w.t.% in Anisole)
- stable and cheap (commercial product)

Molecular weight (M.W.) : 487.66

### ◆ Scintillation yield with respect to concentration of $\text{Zr}(\text{acac})_4$



### ◆ Quenting


- Expected light yield

$$\text{Light yield} = L_0 \times \frac{\sigma_1 N_{\text{PPO}}}{\sigma_1 N_{\text{PPO}} + \sigma_2 N_{\text{Zr}}}$$

$L_0$  : Light yield of anisole  
 $N_{\text{PPO}}$  : No. of PPO molecular  
 $N_{\text{Zr}}$  : No. of  $\text{Zr}(\text{acac})_4$  molecular  
 $\sigma_1$  : absorbance of PPO  
 $\sigma_2$  : absorbance of  $\text{Zr}(\text{acac})_4$


concentration of $\text{Zr}(\text{acac})_4$	Observed ADC channel	Expected ADC channel
0 mg	3850	3850
50mg	3175	3138
100mg	2800	2651
200mg (1 w.t.%)	2000	2018 (52%)
300mg	1600	1613
500mg	900	1178
1000mg (5 w.t.%)	—	695 (18%)

### ◆ tetrakis (isopropyl acetoacetate) Zr



$\text{Zr}(\text{CH}_3\text{COCHCOOCH}(\text{CH}_3)_2)_4$   
 $= \text{Zr}(\text{iprac})_4$   
 M.W. : 711.92

### ◆ tetrakis (ethyl acetoacetate) Zr




$\text{Zr}(\text{CH}_3\text{CCOCHCOOCH}_2\text{CH}_3)_4$   
 $= \text{Zr}(\text{etac})_4$   
 M.W. : 665.81

## 3. Scintillation yield with $\beta$ -keto ester


### ◆ Liquid scintillator containing Zr $\beta$ -keto-ester complex

#### ● Absorbance



**Confirmed absorption peak moves 275nm  $\rightarrow$  245nm in Hexane, but in Acetonitrile .**


#### ● Scintillation light yield



**Observed scintillation light yield decreased (but improved). Still exist absorption peak around 270nm in Anisole.**


### ◆ Liquid scintillator containing $\beta$ -keto ester ligand

#### ● Absorbance




**The absorption peak of  $\beta$ -keto ester ligands were found around at 240nm (not 270nm) even though in an aromatic solvent. Therefore the scintillation light yield recovers almost double @ 5w.t.%.**

#### ● Scintillation light yield




## 4. Zirconium complex with luminescence

### ◆ Zirconium ODZ complex




- good solubility (5w.t.% in Benzonitrile)
- M.W. : 1040.18
- state : yellow powder

### ◆ Luminescence



### ◆ Scintillation light yield




- Emission wavelength : 430nm
- Absorption wavelength : 270nm and 320nm
- Third excitation ( $\sim 340\text{nm}$  from PPO) was used for the emission of  $\text{Zr}(\text{ODZ})_4$ .
- Quantum yield for first excitation ( $\sim 240\text{nm}$ ) was estimated as  $\sim 30\%$ .

**Need another solvent which has shorter emission spectrum than Benzonitrile.**


## 5. Other complex

### ◆ tetrakis (diethyl malonato) Zr



$\text{Zr}(\text{CH}_3\text{CH}_2\text{OCOCCHCOOCH}_2\text{CH}_3)_4$   
 $= \text{Zr}(\text{deml})_4$   
 M.W. : 727.84

### ◆ Absorbance of ligand



**short absorption peak ( $\sim 210\text{nm}$ )**

This work was supported by a Grant-in-Aid for Scientific Research on Innovative Areas (No.24104501) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), and a Grant-in-Aid for Scientific Research (C) (No.24540295) from the Japanese Society for the Promotion of Science (JSPS).