

Transverse Beam Dynamics

JUAS tutorial 2


16 January 2013

1 Exercise: geometry of a storage ring, thin lens, tune, dispersion

Consider a proton synchrotron accumulator made of identical cells. The relevant ring parameters are given in the following table:

Proton kinetic energy	2 GeV
Cell type	Symmetric triplet ^(*)
Ring circumference	960 m
Integrated quadrupole gradient ($\int Gdl$)	1.5 T

(*)Note: A thin lens symmetric triplet cell consists of a thin lens defocusing quadrupole of focal length $-f$, followed with a drift space of length L_1 , a thin lens focusing quadrupole of focal length f , a drift of length L_2 , a thin lens dipole of horizontal bending angle θ , a drift of length L_2 , a thin lens focusing quadrupole of focal length f , a drift of length L_1 , and a thin lens defocusing quadrupole of focal length $-f$ (see Figure below).



1. Compute the focal length f of the quadrupoles. The proton rest mass is 938 MeV.
2. Given the numerical values $L_1 = 1.5$ m and $L_2 = 6$ m:
 - Compute the horizontal and vertical transfer matrices of a triplet cell (take into account that the sign of the focal length changes when going from horizontal to vertical plane).

- Compute the horizontal and vertical machine tunes.
- Compute the horizontal and vertical betatron functions at the entrance of a triplet cell.
- Compute the horizontal and vertical dispersion functions at the entrance of a triplet cell.

Hint:

The 3×3 horizontal transfer matrix for one symmetric triplet cell is (in the thin lens approximation):

$$M_{triplet} = \begin{pmatrix} \frac{f^3 + 2L_1^2 L_2 - 2L_1 f(L_1 + L_2)}{f^3} & \frac{2(f - L_1)(L_1 f + L_2 f - L_1 L_2)}{f^2} & (L_1 + L_2 - \frac{L_1 L_2}{f})\theta \\ \frac{2L_1(L_1 L_2 - L_1 f - f^2)}{f^4} & \frac{f^3 + 2L_1^2 L_2 - 2L_1 f(L_1 + L_2)}{f^3} & \frac{(f^2 + L_1 f - L_1 L_2)}{f^2}\theta \\ 0 & 0 & 1 \end{pmatrix}$$

for the transport of a vector $\begin{pmatrix} x \\ x' \\ \Delta p/p_0 \end{pmatrix}$, where $\Delta p/p_0$ is the momentum offset with respect to the design momentum p_0 .

2 Exercise: emittance

From the solution of the trajectory equation,

$$x = \sqrt{\beta\epsilon} \cos[\phi + \phi_0]$$

- Derive the following relation:

$$\gamma x^2 + 2\alpha x x' + \beta x'^2 = \epsilon$$

where ϵ is the emittance, and β , α , and γ are the so-called Twiss parameters:

$$\alpha \equiv -\frac{\beta'}{2}, \text{ and } \gamma \equiv \frac{1 + \alpha^2}{\beta}$$

(DO NOT CONFUSE this γ with the relativistic Lorentz factor!).