


Michal Mereš\*  
for the ALICE Collaboration  
\*Comenius University


## Baryon number (BN) transport: String junction model

- Valence quarks are connected via non-perturbative configuration of gluon field (SJ).
- Generated baryon is formed around string junction with presence of several sea quarks.
- Deceleration of BN of the beam = BN transport.
- $R = N_{\bar{B}}/N_B$ ,  $B = p, \Lambda, \Xi, \Omega$
- Probability of BN transfer is  $\approx e^{(\alpha-1)\Delta y}$ ,  $\Delta y = y_{beam} - y_{baryon}$
- $\alpha$  depends on configuration in which BN is transported.
- $\alpha_{SJ} = 0.5$  (Veneziano, Rossi) [1]  $\leftrightarrow \alpha_{SJ} = 1$  (Kopeliovich) [2]



## BN transport in heavy-ion collisions

- BN transport can be enhanced by multiple nucleon collisions.
- If there is only small contribution for transport and tight limits on any additional contribution we will observe no centrality dependence for antibaryon-to-baryon ratios and consistent ratios for different baryons.




Pb-Pb collision at  $\sqrt{s_{NN}} = 2.76$  TeV  
p-Pb collision at  $\sqrt{s_{NN}} = 5.02$  TeV

## Analysis

### Reconstruction and identification of baryons

- Protons are identified using their specific ionization in the TPC gas.
- For this analysis, (anti)protons were selected within a band of  $3\sigma$  around the expected value.
- Strange baryons in ALICE are reconstructed by detecting the charged products of their weak decays.
- Each invariant mass distribution is fitted by a combination of Gauss function and polynomial function of second order.
- This fit is used for the determination of the peak position and width.
- Background area was fitted by 2nd order polynomial function. Integral of the function was subtracted from histogram content in the peak region.






## Corrections

- The TPC of the ALICE experiment is symmetric around mid-rapidity and has full azimuthal coverage.
- Many detector effects such as the acceptance, reconstruction and particle identification are the same for particles and anti-particles and thus cancel out in the ratio.
- Effects which we need to correct for are absorption and secondary particles produced in material.

### Absorption correction


- To correct for absorption, we multiply by a correction factor defined as the ratio of generated particles to reconstructible particles.
- Reconstructible particles are considered to be those that have an associated track reference in the TPC in simulation.
- Reconstructible mother is a mother which decayed in the considered charged channel.
- Contamination by secondary protons
- (Anti)protons contaminated by secondaries produced in the material and feed-down.
- Effect taken into account by studying the distribution of distances of closest approach ( $DCA_{xy}$ ) of tracks to the primary vertex.
- $DCA_{xy}$  distributions are very different for primaries and secondaries; data distribution can be fitted with MC template distributions to compute fractions of secondaries.




## Results

### Rapidity and transverse momentum dependence

- Preliminary results show no sign of rapidity and transverse momentum dependence.




- Experimental points are compared with DPMJet model predictions that include variation of BN transport mechanisms.




### Multiplicity density dependence

- Results show no sign of multiplicity density dependence



### Mid-rapidity ratios



## Conclusion

- We measured antibaryon-to-baryon ratios in pp and Pb-Pb collisions.
- Ratios in Pb-Pb and p-Pb are compatible within uncertainties.
- Results from pp, p-Pb and Pb-Pb collisions are very close to unity => The contribution of the BN transport process with  $\alpha_{SJ} = 1$  is compatible with zero for Pb-Pb at  $\sqrt{s_{NN}}=2.76$  and p-Pb at  $\sqrt{s_{NN}}=5.02$ .

[1] B. Z. Kopeliovich and B. Povh, Phys. Lett. B 446, 321 (1999)

[2] G.C.Rossi, G.Veneziano, Nucl. Phys. B123, (1977) 507

[3] STAR Collaboration, Phys. Rev. C 75 (2007) 064901