

Systematic studies of the centrality-dependence of soft photon production in Au+Au collisions with PHENIX

Benjamin Banner for the PHENIX collaboration

Department of Physics, Stony Brook University

2014-05-24

Soft photons $0.4 \text{ GeV}/c < p_T < 5.0 \text{ GeV}/c$

Probe (our understanding of) the full space-time evolution


- initial (hard) scattering
- approach to equilibrium
- QGP evolution
- interactions in HRG

$$R_\gamma = \frac{Y_{\text{incl}}/Y_{\pi^0}}{Y_{\text{decay}}/Y_{\pi^0}} = \frac{\langle \epsilon f \rangle N_{\text{incl}}^\gamma / N_{\pi^0}^\gamma}{Y_{\text{decay}}/Y_{\pi^0}}$$

$$N_{\text{incl}}^\gamma(p_{T\gamma}) = Y_{\text{incl}} a_{ee} \epsilon_{ee} p_{\text{conv}}$$

$$N_{\pi^0}^\gamma(p_{T\gamma}) = Y_{\pi^0} a_{ee} \epsilon_{ee} p_{\text{conv}} \langle \epsilon f \rangle$$

- soft photons difficult to measure in an EMCal
- use conversions of real photons $\gamma \rightarrow ee$



Soft photons $0.4 \text{ GeV}/c < p_T < 5.0 \text{ GeV}/c$

Probe (our understanding of) the full space-time evolution


- initial (hard) scattering
- approach to equilibrium
- QGP evolution
- interactions in HRG

$$R_\gamma = \frac{Y_{\text{incl}}/Y_{\pi^0}}{Y_{\text{decay}}/Y_{\pi^0}} = \frac{\langle \epsilon f \rangle N_{\text{incl}}^\gamma / N_{\pi^0}^\gamma}{Y_{\text{decay}}/Y_{\pi^0}}$$


$$N_{\text{incl}}^\gamma(p_{T,\gamma}) = Y_{\text{incl}} a_{ee} \epsilon_{ee} p_{\text{conv}}$$

$$N_{\pi^0}^\gamma(p_{T,\gamma}) = Y_{\pi^0} a_{ee} \epsilon_{ee} p_{\text{conv}} \langle \epsilon f \rangle$$

- soft photons difficult to measure in an EMCal
- use conversions of real photons $\gamma \rightarrow ee$




$$Y_{\text{direct}} = (R_Y - 1) Y_{\text{decay}}$$



centrality

inverse slope

0-20%	$(239 \pm 25 \pm 7) \text{ MeV}/c$
20-40%	$(260 \pm 33 \pm 8) \text{ MeV}/c$
40-60%	$(225 \pm 28 \pm 6) \text{ MeV}/c$
60-92%	$(238 \pm 50 \pm 6) \text{ MeV}/c$




Excess photon yield grows

$$\propto N_{\text{part}}^{1.48 \pm 0.08 (\text{stat}) \pm 0.04 (\text{syst})}$$

$$v_n^{\text{direct}} = \frac{R_\gamma v_n^{\text{inclusive}} - v_n^{\text{decay}}}{R_\gamma - 1}$$

v_n : weighted average of different sources



- v_2 deep into the “thermal” component of the direct photon signal with pure sample
- low p_T behavior?