Quark Matter 2014 - XXIV International Conference on Ultrarelativistic Nucleus-Nucleus Collisions



Contribution ID: 202

Type: Poster

Measurement of the space-time extent of the emission source in d+Au and Au+Au collisions through charged pion interferometry at $\sqrt{s_{NN}} = 200$ GeV

Tuesday 20 May 2014 16:30 (2 hours)

The recent observation of azimuthal correlations in p+Pb collisions at LHC

is consistent with an interpretation as either an initial state gluon saturation effect or the expansion dynamics of a QGP like medium. It is well known that the latter process is characterized by strong final state re-scattering effects. In this work two-pion interferometry measurements in d+Au and Au+Au collisions at $\sqrt{sNN} = 200 \text{ GeV}$ are performed to extract and compare the Gaussian source radii Rout, Rside and Rlong, which characterize the space-time extent of the emission sources. The comparisons which are carried out as a function of collision centrality and the mean transverse momentum for pion pairs, indicate strikingly similar patterns for the d+Au and Au+Au systems. They also indicate linear dependence of Rside on the initial traverse geometric size, as well

as a smaller freeze-out size for the d+Au system. These patterns point to the important role of final-state re-scattering effects in the reaction dynamics of d+Au collisions.

On behalf of collaboration:

PHENIX

Primary author: AJITANAND, Nuggehalli (SUNY Stony Brook)

Presenter: AJITANAND, Nuggehalli (SUNY Stony Brook)

Session Classification: Poster session

Track Classification: Correlations and Fluctuations