


# Electroweak Corrections to Gauge-Boson Pair Production at the LHC

Anastasiya Bierweiler



In collaboration with Tobias Kasprzik,  
Johann H. Kühn and Sandro Uccirati

JHEP 1211 (2012) 093  
PoS(ICHEP2012)078

Phenomenology Symposium 6.-8. May 2013




- I. Motivation and Introduction**
- II. Theoretical Status of QCD and Electroweak Corrections**
- III. Numerical Results**
- IV. Conclusions and Outlook**



- **WW/ZZ** is an important **irreducible background** to inclusive SM Higgs boson production.
- Gauge boson pair production provides an excellent opportunity to test **the non-abelian structure** of triple gauge boson couplings of the SM at high energies.
  - ↪ constraint on non-standard  $\gamma WW$  and  $ZWW$  couplings.
  - ↪ Search for anomalous triple and quartic couplings.
- Backgrounds to **new physics searches**, i.e. leptons +  $\cancel{E}_T$  signatures.
  - ↪ Pair production of supersymmetric particles.

## QCD Corrections to $V_1 V_2$ -pair production:

- NLO QCD corrections are available for full process including leptonic decays. [Dixon, Kunszt, Signer '99; Campbell, Ellis '99; De Florian, Signer '00]
- Results matched with parton showers and combined with soft-gluon resummation. [S. Frixione, B.R. Weber '06; P. Nason, G. Redolfi '06]
- QCD corrections dominated by tree-level  $q\bar{q}$  annihilation channels. Significant contributions of the channels  $gg \rightarrow V_1 V_2 \sim 10\%$  to LO, although formally at  $\mathcal{O}(\alpha_s^2)$ . [Dührssen et.al. '05; Glover, van der Bij '89; Kao, Dicus'91]



- By considering event selection for Higgs searches even corrections of 30% can be obtained. [Binoth et. al. '06]

## EW corrections to gauge-boson pair production:

- Complete  $\mathcal{O}(\alpha)$  corrections known to  $pp \rightarrow W\gamma \rightarrow l\bar{\nu}\gamma + X$  in single pole approximation. [Accomando, Denner, Pozzorini '01; Accomando, Denner, Meier '05]
- Complete  $\mathcal{O}(\alpha)$  corrections for on-shell  $Z$  bosons  $pp \rightarrow Z\gamma$ . [Hollik, Meier '04]
- Complete  $\mathcal{O}(\alpha)$  corrections for  $pp \rightarrow Z\gamma \rightarrow ll\gamma + X$  in a single pole approximation. [Accomando, Denner, Meier '05]
- Complete  $\mathcal{O}(\alpha)$  corrections for  $pp \rightarrow WW, ZZ, WZ \rightarrow 4l + X$  in high energy and pole-approximation.  
→ Large negative EW corrections at large transverse momentum.  
[Accomando, Denner, Pozzorini '01; Accomando, Denner, Kaiser '04]
- NNLL effects at two loops for  $WW$  channel.  
[Kühn, Metzler, Penin, Uccirati '11]



We have calculated the full one-loop corrections to gauge-boson pair production at the LHC.

[A. Bierweiler, T. Kasprzik, J.H. Kühn, S. Uccirati '12; A. Bierweiler, T. Kasprzik, J.H. Kühn'12]

---

**Default cuts:**  $p_{T,W^\pm} > 15 \text{ GeV}$ ,  $|y_{W^\pm}| < 2.5$

MSTW2008LO PDFs [Martin et al. 2009]



- assume  $\int \mathcal{L} dt = 200 \text{ fb}^{-1} \Rightarrow 1000 \text{ WW events with } M_{WW} > 2 \text{ TeV}$
- rapidly increasing admixture of  $\gamma\gamma \rightarrow WW$ , since  $\sigma(\gamma\gamma \rightarrow WW) \xrightarrow{\hat{s} \rightarrow \infty} 8\pi\alpha^2/M_W^2$
- **sizable** (up to -30%) **negative EW corrections**

## No compensation between $\gamma\gamma \rightarrow W^+W^-$ and weak corrections! ⇒ Different angular distributions

- $\gamma\gamma \rightarrow W^+W^-$  :  
strong enhancement in forward and backward directions
- **weak corrections:**



negative Sudakov Logarithms for large  $\hat{s}$  and  $\hat{t}$   
⇒ negative corrections for large scattering angles

⇒ Implications for  $d\sigma/d\Delta y_{WW}$  with  $\Delta y_{WW} = y_{W^+} - y_{W^-}$

(for fixed  $M_{WW}$  this corresponds to the angular distribution in the  $WW$ -rest frame)

**High energy cuts:**  $p_{T,W^\pm} > 15 \text{ GeV}$ ,  $|y_{W^\pm}| < 2.5$ ,  $M_{WW} > 1 \text{ TeV}$



MSTW2008LO PDFs [Martin et al. 2009]



- WW production dominated by small scattering angles
- drastic forward-backward peaking of  $\gamma\gamma \rightarrow WW$
- drastic distortion of angular distribution
- $\Sigma\delta$  varies from  $-30\%$  and  $+45\%$  for  $M_{WW} > 1 \text{ TeV}$

**Default cuts:**  $p_{T,V} > 15 \text{ GeV}$ ,  $|y_V| < 2.5$

MSTW2008LO PDFs [Martin et al. 2009]



- Largest EW corrections to ZZ production and small EW corrections for  $\gamma\gamma$  production
- EW corrections to  $W^+ Z$ -production coincide with those to  $W^- Z$ -production at the percent level as a consequence of the identical corresponding unpolarized partonic cross section.

## Conclusion and Outlook:

- **Understanding of gauge boson pair production processes crucial at the LHC:**
  - Understand SM at high energies
  - Understand background to Higgs- and BSM physic searches
- **We have computed the full EW corrections to V-pair production at the LHC:**
  - Small corrections to the total cross section
  - **But:** Sizable negative corrections at large transverse momenta and high invariant mass  $M_{VV}$
  - W-pair production: significant contribution of photon-induced channel at large scattering angles
- **Future work:**
  - Leptonic decays of the vector bosons should be included

## Conclusion and Outlook:

- **Understanding of gauge boson pair production processes crucial at the LHC:**
  - Understand SM at high energies
  - Understand background to Higgs- and BSM physic searches
- **We have computed the full EW corrections to V-pair production at the LHC:**
  - Small corrections to the total cross section
  - **But:** Sizable negative corrections at large transverse momenta and high invariant mass  $M_{VV}$
  - W-pair production: significant contribution of photon-induced channel at large scattering angles
- **Future work:**
  - Leptonic decays of the vector bosons should be included

**Thank you!**

---

# Backup

## EW Corrections at High Energies:

High-energy limit:

$$\hat{s} \sim |\hat{t}| \sim |\hat{u}| \gg M^2 := M_W^2 \simeq M_Z^2 \sim M_H^2 \sim M_t^2 \gg m_f^2 \gg \underbrace{m_\gamma^2}_{\text{IR-regulator}}$$

→ bosons are produced at large  $p_T$

- EW corrections are enhanced by universal large logarithms that originate from mass singularities  $\text{Log}(s/M^2)$  and grow with energy:


$$\underbrace{\alpha^L \text{Log}^{2L}(s/M^2)}_{\text{LL}}, \quad \underbrace{\alpha^L \text{Log}^{2L-1}(s/M^2)}_{\text{NLL}}, \dots$$

- Corrections are of  $\sim -50\%$  at  $p_T = 1\text{TeV}$  (WW production)


**Note:** Change of sign going from LL to NLL (to NNLL etc.)  
→ substantial cancellations are possible!

We have calculated the full one-loop corrections to W-pair production at the LHC. [A. Bierweiler, T. Kasprzik, J.H. Kühn, S. Uccirati '12]

- Partonic leading order contributions at  $\mathcal{O}(\alpha^2)$ :




- Photon-induced contributions at  $\mathcal{O}(\alpha^2)$ :



- MSTW2004qed PDF set is adopted [Martin et al. 2005]
- Potentially large contribution at large invariant masses

## NLO Electroweak Corrections:

### Virtual Electroweak Corrections:



- **Renormalization:**

- masses and fields renormalized in the **on-shell** scheme
- We work in  $G_\mu$  scheme: renormalization of  $\alpha_{G_\mu}$  is defined through the Fermi constant related to the muon life time

$$\alpha_{G_\mu} = \frac{\sqrt{2}G_\mu M_W^2 s_w^2}{\pi} = \frac{\alpha(0)}{1 - \Delta r}$$

$\Delta r$  includes the corrections to muon lifetime not contained in QED-improved Fermi model

⇒ The EW corrections are independent of logarithms of the light quark masses.

**Thomson-limit:** (zero momentum transfer)

$$\delta Z_e|_{\alpha(0)} = \frac{1}{2} \frac{\partial \Sigma_T^{\gamma\gamma}(k^2)}{\partial k^2} \Big|_{k^2=0} - \frac{s_w}{c_w} \frac{\partial \Sigma_T^{\gamma\gamma}(0)}{M_Z^2}$$

$\Sigma_T^{\gamma\gamma}(k^2)$  is transverse part of  $\gamma\gamma$  self-energy with momentum transfer  $k$ .

⇒ charge renormalization constant contains logarithms of light fermion masses including large corrections proportional to  $\sim \alpha \log(m_f^2/\hat{s})$

**$G_\mu$  scheme:** The transition from  $\alpha(0)$  to  $G_\mu$  is ruled by the weak corrections to muon decay  $\Delta r$ :

$$\alpha(G_\mu) = \frac{\sqrt{2} G_\mu M_W^2 s_w^2}{\pi} = \alpha(0)(1 + \Delta r) + \mathcal{O}(\alpha^3)$$

$$\Rightarrow \delta Z_e|_{\alpha(G_\mu)} = \delta Z_e|_{\alpha(0)} - \frac{\Delta r}{2}$$

⇒ large fermion mass logs are resummed in  $G_\mu$  scheme

⇒ the lowest order cross section in  $G_\mu$  parametrization absorbs large universal corrections to SU(2) gauge coupling introduced by  $\rho$  parameter

## IR Singularities:


- **Virtual Corrections:**
  - Soft singularities arise due to soft photons
  - Initial-state collinear singularities arise due to collinear photon radiation off initial-state quarks
  - introduce small **quark mass**  $m_f$  and **photon mass**  $m_\gamma$  to regularize divergences
    - results contain unphysical  $\log(m_f)$  and  $\log(m_\gamma)$
- **Real Radiation:**

IR divergent (also regularized by  $m_\gamma$  and  $m_f$ ), compensated partially by virtual corrections. Remaining collinear singularities have to be absorbed in PDFs.
  - renormalization of PDFs is required

⇒ **Apply phase-space slicing for numerically stable evaluation of phase-space integral**

**Default cuts:**  $p_{T,W^\pm} > 15 \text{ GeV}$ ,  $|y_{W^\pm}| < 2.5$



MSTW2008LO PDFs [Martin et al. 2009]



- Cross section dominated by central events with W-pair of low invariant mass and relatively isotropic distributions
- stronger preference of  $gg$ -induced process for small  $\Delta y_{WW}$
- enhancement of the  $\gamma\gamma$  process for small scattering angles



**Default cuts:**  $p_{T,V} > 15 \text{ GeV}$ ,  $|y_V| < 2.5$

MSTW2008LO PDFs [Martin et al. 2009]





**Default cuts:**  $p_{T,V} > 15 \text{ GeV}$ ,  $|y_V| < 2.5$

MSTW2008LO PDFs [Martin et al. 2009]





**Default cuts:**  $p_{T,V} > 15 \text{ GeV}$ ,  $|y_V| < 2.5$   
MSTW2008LO PDFs [Martin et al. 2009]





**Default cuts:**  $p_{T,V} > 15 \text{ GeV}$ ,  $|y_V| < 2.5$

MSTW2008LO PDFs [Martin et al. 2009]





**Default cuts:**  $p_{T,V} > 15 \text{ GeV}$ ,  $|y_V| < 2.5$

MSTW2008LO PDFs [Martin et al. 2009]





**Default cuts:**  $p_{T,W^\pm} > 15 \text{ GeV}$ ,  $|y_{W^\pm}| < 2.5$

MSTW2008LO PDFs [Martin et al. 2009]



**Default cuts:**  $p_{T,W^\pm} > 15 \text{ GeV}$ ,  $|y_{W^\pm}| < 2.5$

MSTW2008LO PDFs [Martin et al. 2009]



## Technicalities:

Two different approaches to calculate **virtual corrections** and **bremsstrahlung amplitudes** to  $pp \rightarrow VV'$ :

**Virtual corrections and bremsstrahlung amplitudes** computed in the **QGRAF** and **FORM** framework:

- **QGRAF**: [Nogueira] generation of Feynman diagrams
- **FORM**: [Vermaseren] implementation of Feynman rules, reconstruction of Dirac structures, introduction of tensor coefficients and their algebraical reduction to scalar integrals, analytical calculation of squared amplitudes, generation of **FORTRAN** code

Numerically-stable evaluation of scalar-integrals done within **LoopTools - 2.5**.

Numerical phase-space integration done within Fortran using the **Vegas** algorithm.

---

**Virtual corrections** computed in the **FeynArts/FormCalc/LoopTools(FF)** framework:

- **FeynArts** - 3.5: [Küblbeck et al '90] automatic generation of diagrams, calculation of amplitudes
- **FormCalc** - 6.1: [Hahn et al '99] introduction of tensor coefficients; analytical calculation of squared amplitudes; spin-, colour- and polarization sums; generation of **FORTRAN** code
- **LoopTools** - 2.5: [Vermaseren '90] numerical Passarino-Veltman reduction within **Fortran**;