

The Two-Loop Analog of the Passarino-Veltman Result And Beyond

Robert M. Schabinger

with Andreas von Manteuffel

The PRISMA Cluster of Excellence and
Mainz Institute of Theoretical Physics

A Basis for the Virtual Corrections to One-Loop Hard Scattering Processes

$$\begin{aligned}
 A^{1\text{-loop}}(k_1^{h_1}, \dots, k_n^{h_n}) = & \sum_{\alpha_5} C_5^{(\alpha_4)} I_5^{(\alpha_5)} + \sum_{\alpha_4} C_4^{(\alpha_4)} I_4^{(\alpha_4)} + \sum_{\alpha_3} C_3^{(\alpha_3)} I_3^{(\alpha_3)} \\
 & + \sum_{\alpha_2} C_2^{(\alpha_2)} I_2^{(\alpha_2)} + \sum_{\alpha_1} C_1^{(\alpha_1)} I_1^{(\alpha_1)}
 \end{aligned}$$

- It has been known for a long time (G. Passarino and M. J. G. Veltman, Nucl. Phys. **B160**, 151, 1979) that, for most phenomenologically interesting calculations, it is prohibitively inefficient to blindly calculate
- The program of finding an integral basis valid for arbitrary one-loop processes in renormalizable theories was begun by Passarino and Veltman ($2 \rightarrow 2$) and completed by Bern, Dixon, and Kosower (Z. Bern, L. J. Dixon, and D. A. Kosower, Phys. Lett. **B302**, 299, 1993)

A Basis for the Virtual Corrections to Two-Loop Hard Scattering Processes?

$$A^{2\text{-loop}}(k_1^{h_1}, \dots, k_n^{h_n}) = ???$$

A Basis for the Virtual Corrections to Two-Loop Hard Scattering Processes?

$$A^{2\text{-loop}}(k_1^{h_1}, \dots, k_n^{h_n}) = ???$$

**The two-loop integral basis is not known
in the generic case, even for $2 \rightarrow 2$!**

see J. Gluza, K. Kajda, and D. Kosower, Phys. Rev. **D83**, 045012, 2011

for partial results in a few special cases

A Basis for the Virtual Corrections to Two-Loop Hard Scattering Processes?

$$A^{2\text{-loop}}(k_1^{h_1}, \dots, k_n^{h_n}) = ???$$

**The two-loop integral basis is not known
in the generic case, even for $2 \rightarrow 2$!**

see J. Gluza, K. Kajda, and D. Kosower, Phys. Rev. **D83**, 045012, 2011

for partial results in a few special cases

How do we find it?

A Basis for the Virtual Corrections to Two-Loop Hard Scattering Processes?

$$A^{2\text{-loop}}(k_1^{h_1}, \dots, k_n^{h_n}) = ???$$

The two-loop integral basis is not known in the generic case, even for $2 \rightarrow 2$!

see J. Gluza, K. Kajda, and D. Kosower, Phys. Rev. **D83**, 045012, 2011

for partial results in a few special cases

How do we find it?

Integration By Parts Relations

K. Chetyrkin and F. Tkachov, Nucl. Phys. **B192**, 159, 1981

Can We Solve These IBP Relations?

Suppose we want to solve the system of IBP recurrence relations to determine the master integrals for a given multi-loop topology:

- For most interesting examples a highly non-trivial system of recurrence relations results
- Difficult or impossible to solve by hand
- A well-known algorithm due to Laporta (S. Laporta, *Int. J. Mod. Phys. A* **15**, 5087, 2000) reduces the problem to the solution of (usually) a very large system of linear equations


Laporta's algorithm, while very important, requires significant computational resources for most interesting two-loop topologies. What can we do to reduce the computational complexity?

It turns out that determining the master integrals is much simpler than actually solving the IBPs:


- Find all independent *sectors*, collections of integrals with the same set of propagator denominators, for the Feynman integrals that could potentially arise from Feynman diagrams
- Observe that the correlations between sectors can be ignored if all one wants are the master integrals. This allows for a sector-by-sector divide-and-conquer approach
- Use a phase space point where all internal masses, external masses, and Mandelstam invariants are set to primes. This effectively reduces a many-scale problem to a no-scale problem

It makes sense to work within the framework of **Reduze 2** (A. von Manteuffel and C. Studerus, arXiv:1201.4330) because most of the necessary code is already there

The Pentatriangle Topology




The Planar Double-Box Topology






$$\begin{aligned}
 & I \left[(q_1 - k_4)^2 - m_8^2 \right] \quad I \left[\left((q_1 - k_4)^2 - m_8^2 \right) \left((q_2 - k_1)^2 - m_9^2 \right) \right] \\
 & I \left[\left((q_1 - k_4)^2 - m_8^2 \right)^2 \right] \quad I \left[(q_2 - k_1)^2 - m_9^2 \right]
 \end{aligned}$$


The Non-Planar Double-Box Topology




$$\begin{aligned}
 & I \left[\left((q_1 - k_4)^2 - m_8^2 \right) \left((q_1 - q_2 + k_1 + k_2)^2 - m_9^2 \right) \right] \\
 & I \left[(q_1 - k_4)^2 - m_8^2 \right] \quad I \left[\left((q_1 - k_4)^2 - m_8^2 \right)^2 \right] \quad I \left[\left((q_1 - k_4)^2 - m_8^2 \right)^3 \right] \\
 & I \left[(q_1 - q_2 + k_1 + k_2)^2 - m_9^2 \right] \quad I \left[\left((q_1 - q_2 + k_1 + k_2)^2 - m_9^2 \right)^2 \right]
 \end{aligned}$$

Six-Propagator Topologies






$$I \left[(q_2 - k_4)^2 - m_2^2 \right] \quad I \left[\left((q_2 - k_4)^2 - m_2^2 \right)^2 \right]$$








$$I [q_1^2 - m_7^2] \quad I [(q_1^2 - m_7^2)^2] \quad I [((q_1 - k_4)^2 - m_8^2) ((q_1 - q_2 + k_1 + k_2)^2 - m_9^2)]$$





$$\begin{aligned}
 & I \left[(q_1 - k_4)^2 - m_8^2 \right] \quad I \left[\left((q_1 - k_4)^2 - m_8^2 \right) \left((q_1 - q_2 + k_1 + k_2)^2 - m_9^2 \right) \right] \\
 & I \left[\left((q_1 - k_4)^2 - m_8^2 \right)^2 \right] \quad I \left[\left((q_1 - k_4)^2 - m_8^2 \right) \left((q_2 - k_1 - k_2)^2 - m_3^2 \right) \right] \\
 & I \left[\left((q_2 - k_1 - k_2)^2 - m_3^2 \right) \left((q_1 - q_2 + k_1 + k_2)^2 - m_9^2 \right) \right] \quad I \left[(q_1 - q_2 + k_1 + k_2)^2 - m_9^2 \right] \\
 & I \left[(q_2 - k_1 - k_2)^2 - m_3^2 \right] \quad I \left[\left((q_2 - k_1 - k_2)^2 - m_3^2 \right)^2 \right] \quad I \left[\left((q_2 - k_1 - k_2)^2 - m_3^2 \right)^3 \right] \\
 & I \left[\left((q_1 - k_4)^2 - m_8^2 \right) \left((q_2 - k_1 - k_2)^2 - m_3^2 \right)^2 \right] \quad I \left[\left((q_1 - q_2 + k_1 + k_2)^2 - m_9^2 \right)^2 \right] \\
 & I \left[\left((q_2 - k_1 - k_2)^2 - m_3^2 \right)^2 \left((q_1 - q_2 + k_1 + k_2)^2 - m_9^2 \right) \right]
 \end{aligned}$$

Five-Propagator Topologies





$$I \left[(q_1 - q_2 + k_2)^2 - m_5^2 \right] \quad I \left[\left((q_1 - q_2 + k_2)^2 - m_5^2 \right)^2 \right] \quad I \left[q_1^2 - m_7^2 \right]$$



$$I_L [(q_2 - k_4)^2 - m_2^2]$$

$$I_L [q_1^2 - m_7^2]$$

$$I_L [((q_2 - k_4)^2 - m_2^2)^2]$$


$$I_L [((q_2 - k_4)^2 - m_2^2) (q_1^2 - m_7^2)]$$

$$I_R [q_2^2 - m_1^2]$$

$$I_R [q_1^2 - m_7^2]$$

$$I_R [(q_2^2 - m_1^2)^2]$$

$$I_R [(q_2^2 - m_1^2) (q_1^2 - m_7^2)]$$



$$I [q_1^2 - m_7^2]$$

$$I [(q_1 - k_4)^2 - m_8^2]$$

$$I [(q_2 - k_1 - k_2)^2 - m_3^2]$$



$$I [(q_1 - q_2 + k_1 + k_2)^2 - m_9^2]$$


$$I [((q_2 - k_1 - k_2)^2 - m_3^2) (q_1^2 - m_7^2)]$$

$$I [((q_2 - k_1 - k_2)^2 - m_3^2)^2]$$

$$I [((q_2 - k_1 - k_2)^2 - m_3^2) ((q_1 - k_4)^2 - m_8^2)]$$

Four-Propagator Topologies






$$\begin{aligned}
 & I_L [(q_2 - k_4)^2 - m_2^2] \quad I_L [((q_2 - k_4)^2 - m_2^2)^2] \quad I_L [(q_1 - q_2 + k_2)^2 - m_5^2] \\
 & I_R [(q_2 - k_1 - k_2)^2 - m_3^2] \quad I_R [((q_2 - k_1 - k_2)^2 - m_3^2)^2] \quad I_R [(q_1 - q_2)^2 - m_6^2]
 \end{aligned}$$



The Non-Trivial Vacuum Topology

A. I. Davydychev, Phys. Rev. **D61**, 087701, 2000



The Sunrise Topologies

L. Adams, C. Bogner, and S. Weinzierl, arXiv:1302.7004



$$\begin{array}{lll}
 I_L [(q_2 - k_4)^2 - m_2^2] & I_L [((q_2 - k_4)^2 - m_2^2)^2] & I_L [(q_1 - q_2)^2 - m_6^2] \\
 I_R [q_2^2 - m_1^2] & I_R [(q_2^2 - m_1^2)^2] & I_R [(q_1 - q_2)^2 - m_6^2]
 \end{array}$$

Outlook

Although our code applies to arbitrary scattering processes, limited only by computer time, there is clearly still a very long way to go if the goal is to build a fully automated two-loop program such as those that already exist at one-loop

- Solve the remaining phenomenologically important masters for $2 \rightarrow 2$ processes (*e.g.* those needed for the NNLO wishlist)
- Improve the efficiency of the Reduze 2 IBP relation solver
- Start looking at other situations of potential phenomenological interest where a fully automated approach would be welcome (*e.g.* two-loop $2 \rightarrow 3$)

Trivial Topologies

