



# PROPERTIES OF THE HIGGS BOSON WE CANNOT “UNSEE”

André David (CERN) – Rencontres de Blois 2013



# Things you can't "unsee"

2

[<http://cern.ch/go/Dxh7>]





# Things you can't "unsee"

3

[<http://cern.ch/go/Dxh7>]





# Things you can't "unsee"

4


[<http://cern.ch/go/Dxh7>]





# (self-inflicted) Mission: impossible

5



Channel	Conference note	L	Date
Spin Combination	<a href="#">ATLAS-CONF-2013-040</a>	up to 25 fb <sup>-1</sup>	16/04/2013
Couplings Combination	<a href="#">ATLAS-CONF-2013-034</a>	up to 25 fb <sup>-1</sup>	14/03/2013
Higgs to Diphoton spin	<a href="#">ATLAS-CONF-2013-029</a>	21 fb <sup>-1</sup>	13/03/2013
Higgs to WW(lvlv) spin	<a href="#">ATLAS-CONF-2013-031</a>	21 fb <sup>-1</sup>	11/03/2013
Higgs to WW(lvlv)	<a href="#">ATLAS-CONF-2013-030</a>	25 fb <sup>-1</sup>	11/03/2013
2HDM WW(lvlv)	<a href="#">ATLAS-CONF-2013-027</a>	13 fb <sup>-1</sup>	11/03/2013
Combined of Mass	<a href="#">ATLAS-CONF-2013-014</a>	up to 25 fb <sup>-1</sup>	05/03/2013
Higgs to Diphoton	<a href="#">ATLAS-CONF-2013-012</a>	25 fb <sup>-1</sup>	05/03/2013
Higgs to 4 leptons	<a href="#">ATLAS-CONF-2013-013</a>	25 fb <sup>-1</sup>	05/03/2013
ZH (invisible decays)	<a href="#">ATLAS-CONF-2013-011</a>	18 fb <sup>-1</sup>	05/03/2013
Higgs to dimuon	<a href="#">ATLAS-CONF-2013-010</a>	21 fb <sup>-1</sup>	05/03/2013
Higgs to Zgamma	<a href="#">ATLAS-CONF-2013-009</a>	25 fb <sup>-1</sup>	05/03/2013





May-2013	Full 8 TeV dataset: VBF H, H -> bb	<a href="#">TWiki</a> , <a href="#">PAS</a>
May-2013	Full 8 TeV dataset: ttH, H -> gamma gamma	<a href="#">TWiki</a> , <a href="#">PAS</a>
May-2013	Full 7+8 TeV dataset: VH, H -> bb	<a href="#">TWiki</a> , <a href="#">PAS</a>
May-2013	Full 8 TeV dataset: H -> WW -> lnuJ	<a href="#">TWiki</a> , <a href="#">PAS</a>
May-2013	Full 7+8 TeV dataset: H -> ZZ -> 2l2nu	<a href="#">TWiki</a> , <a href="#">PAS</a>
Apr-2013	Moriond Higgs Combination	<a href="#">TWiki</a> , <a href="#">PAS</a>
Mar-2013	Full 7+8 TeV dataset: H -> gamma gamma	<a href="#">TWiki</a> , <a href="#">PAS</a>
Mar-2013	Full 7+8 TeV dataset: H -> ZZ -> 4l	<a href="#">TWiki</a> , <a href="#">PAS</a>
Mar-2013	Full 7+8 TeV dataset: H -> WW -> 2l2nu	<a href="#">TWiki</a> , <a href="#">PAS</a>
Mar-2013	Full 7+8 TeV dataset: H -> tau tau	<a href="#">TWiki</a> , <a href="#">PAS</a>
Mar-2013	Full 7+8 TeV dataset: H -> Z gamma	<a href="#">TWiki</a> , <a href="#">PAS</a>
Mar-2013	Full 7+8 TeV dataset: H -> WWW -> 3l3nu	<a href="#">TWiki</a> , <a href="#">PAS</a>
Mar-2013	Full 7+8 TeV dataset: VH -> tau tau	<a href="#">TWiki</a> , <a href="#">PAS</a>

□ Present a coherent view of present-day results of Higgs properties from the LHC and Tevatron experiments.

□ Any omission or mistake are the speaker's fault.

# A tribute to those doing SM calculations

“Yesterday’s discovery is today’s calibration, and tomorrow’s background.” – V. L. Telegdi






# Higgs in CMS – ca. 2008

7


[<http://cern.ch/go/dJf7>] [<http://cern.ch/go/Sx8m>]




- **Mass mechanism** – the mexican hat field, first published by Brout and Englert (1964).
- **Higgs boson** – the field's massive radial excitation, tacit to Brout and Englert, massless via approximations in Guralnik et al., and explicitly mentioned by Higgs (1964).
- **Viability** – photons and massive weak bosons can coexist was shown by Kibble (1967).

# How SM Higgses are born


□ **Gluon fusion**






□ **VBF**



□ **VH**




□ **ttH**







# How SM Higgses die




□ Coupling and kinematics drive BR ( $b\bar{b}$ ,  $WW$ ,  $\tau\tau$ ,  $ZZ$ ).

□ Decays with photons ( $\gamma\gamma$ ,  $Z\gamma$ ) only through loops.

Near to maximal  $\Pi BR_i \rightarrow$












# Looking up to a new boson





# Higgsdependence day recap

- **Both experiments at  $5.0\sigma$ .**
  - One above expectations...  
 $\sigma_{\text{ATLAS}}/\sigma_{\text{SM}} = 1.2 \pm 0.3$  (at 126.5 GeV).
  - ...the other one below.  
 $\sigma_{\text{CMS}}/\sigma_{\text{SM}} = 0.80 \pm 0.20$  (at 125 GeV).
- **Mass**
  - ATLAS: min. p-value at 126.5 GeV.
  - CMS:  $m_\chi = 125.3 \pm 0.6$  GeV.
- **“Proto-couplings” compatible with SM.**
- **“More data needed...”**



# From the other side of the pond



- Combination of Tevatron  $VH \rightarrow b\bar{b}$  searches, in July 2012:
  - **$2.8 \sigma$  local significance at  $m_H = 125 \text{ GeV}$ .**



2012 2011 2010 2009 2008

## Who Should Be TIME's Person of the Year 2012? >

As always, TIME's editors will choose the Person of the Year, but that doesn't mean readers shouldn't have their say. Cast your vote for the person you think most influenced the news this year for better or worse. Voting closes at 11:59 p.m. on Dec. 12, and the winner will be announced on Dec. 14.


Like 1.5k Tweet 538 +1 20 Share 7

### THE CANDIDATES

## The Higgs Boson

By Jeffrey Kluger | Monday, Nov. 26, 2012

◀ 18 of 40 ▶



SSPL/GETTY IMAGES

Simulation of a Higgs-Boson decaying into four muons, CERN, 1990.

### What do you think?

Should **The Higgs Boson** be TIME's Person of the Year 2012?

Definitely  No Way

VOTE

Take a moment to thank this little particle for all the work it does, because without it, you'd be just inchoate energy without so much as a bit of mass. What's more, the same would be true for the entire universe. It was in the 1960s that Scottish physicist Peter Higgs first posited the existence of a particle that causes energy to make the jump to matter. But it was not until last summer that a team of researchers at Europe's Large Hadron Collider — Rolf Heuer, Joseph Incandela and Fabiola Gianotti — at last sealed the deal and in so doing finally fully confirmed Einstein's general theory of relativity. The Higgs — as particles do — immediately decayed to more-fundamental particles, but the scientists would surely be happy to collect any honors or awards in its stead.

Photos: Step inside the Large Hadron Collider.

### WHO SHOULD BE TIME'S PERSON OF THE YEAR 2012?

The Candidates

Video

Poll Results

### PAST PERSONS OF THE YEAR



2011: The Protester



2010: Facebook's Mark Zuckerberg



2009: Ben Bernanke



2008: Barack Obama

Most Read

Most Emailed

- 1 Who Should Be TIME's Person of the Year 2012?
- 2 LIFE Behind the Picture: The Photo That Changed the Face of AIDS
- 3 Nativity-Scene Battles: Score One for the Atheists
- 4 The \$7 Cup of Starbucks: A Logical Extension of the Coffee Chain's Long-Term Strategy

2012 2011 2010 2009 2008

## Who Should Be TIME's Person of the Year 2012? >

As always, TIME's editors will choose the Person of the Year, but that doesn't mean readers shouldn't have their say. Cast your vote for the person you think most influenced the news this year for better or worse. Voting closes at 11:59 p.m. on Dec. 12, and the winner will be announced on Dec. 14.

Like 1.5k Tweet 538 +1 20 Share 7


### THE CANDIDATES

## The Higgs Boson

By Jeffrey Kluger | Monday, Nov. 26, 2012

◀ 18 of 40 ▶

Simulation of a Higgs-Boson decaying into four muons, CERN, 1990.



SSPL/GETTY IMAGES

Simulation of a Higgs-Boson decaying into four muons, CERN, 1990.



Take a moment to thank this little particle for all the work it does, because without it, you'd be just inchoate energy without so much as a bit of mass. What's more, the same would be true for the entire universe. It was in the 1960s that Scottish physicist Peter Higgs first posited the existence of a particle that causes energy to make the jump to matter. But it was not until last summer that a team of researchers at Europe's Large Hadron Collider — Rolf Heuer, Joseph Incandela and Fabiola Gianotti — at last sealed the deal and in so doing finally fully confirmed Einstein's general theory of relativity. The Higgs — as particles do — immediately decayed to more-fundamental particles, but the scientists would surely be happy to collect any honors or awards in its stead.

Photos: Step inside the Large Hadron Collider.

## WHO SHOULD BE TIME'S PERSON OF THE YEAR 2012?

The Candidates

Video

Poll Results

### PAST PERSONS OF THE YEAR



2011: The Protester



2010: Facebook's Mark Zuckerberg



2009: Ben Bernanke



2008: Barack Obama

Most Read

Most Emailed

- Who Should Be TIME's Person of the Year 2012?
- LIFE Behind the Picture: The Photo That Changed the Face of AIDS
- Nativity-Scene Battles: Score One for the Atheists
- The \$7 Cup of Starbucks: A Logical Extension of the Coffee Chain's Long-Term Strategy

2012 2011 2010 2009 2008

## Who Should Be TIME's Person of the Year 2012? >


As always, TIME's editors will choose the Person of the Year, but that doesn't mean readers shouldn't have their say. Cast your vote for the person you think most influenced the news this year for better or worse. Voting closes at 11:59 p.m. on Dec. 12, and the winner will be announced on Dec. 14.

Like 1.5k Tweet 538 +1 20 Share 7

### THE CANDIDATES

## The Higgs Boson

By Jeffrey Kluger | Monday, Nov. 26, 2012



### What do you think?

Should The Higgs Boson be TIME's Person of the Year 2012?

Definitely  No Way

VOTE

◀ 18 of 40 ▶

## WHO SHOULD BE TIME'S PERSON OF THE YEAR 2012?

The Candidates

Video

Poll Results

### PAST PERSONS OF THE YEAR



2011: The Protester



2010: Facebook's Mark Zuckerberg




last summer that a team of researchers at Europe's Large Hadron Collider — Rolf Heuer, Joseph Incandela and Fabiola Gianotti — at last sealed the deal and in so doing finally fully confirmed Einstein's general theory of relativity. The









# Timeline of the results





# The build up of a signal

18



- Thanks to the excellent performance of the LHC !
- $> 15 \text{ fb}^{-1}$  delivered after July 2012.



# Where we stand today

19

Significance Obs. (pre-fit exp.)	$H \rightarrow ZZ$	$H \rightarrow \gamma \gamma$	$H \rightarrow WW$	$H \rightarrow b\bar{b}$	$H \rightarrow \tau \tau$
<b>ATLAS</b>	<b>6.6 <math>\sigma</math></b> (4.4 $\sigma$ )	<b>7.4 <math>\sigma</math></b> (4.1 $\sigma$ )	<b>2.5 <math>\sigma</math></b> (1.6 $\sigma$ )	<b>-0.4 <math>\sigma</math></b> (1.0 $\sigma$ )	<b>1.1 <math>\sigma</math></b> (1.7 $\sigma$ )
	124.3 GeV		126.8 GeV	125 GeV	
<b>CMS</b>	<b>6.7 <math>\sigma</math></b> (7.1 $\sigma$ )	<b>3.9 <math>\sigma</math></b> (4.2 $\sigma$ )	<b>3.9 <math>\sigma</math></b> (5.6 $\sigma$ )	<b>2.0 <math>\sigma</math></b> (2.1 $\sigma$ )	<b>2.8 <math>\sigma</math></b> (2.7 $\sigma$ )
	125.7 GeV				

- Combined p-values  $< 10^{-20}$  are telling us to make measurements...



	Test statistic	Profiled?	Test statistic sampling
LEP	$q_\mu = -2 \ln \frac{\mathcal{L}(data \mu, \tilde{\theta})}{\mathcal{L}(data 0, \tilde{\theta})}$	no	Bayesian-frequentist hybrid
Tevatron	$q_\mu = -2 \ln \frac{\mathcal{L}(data \mu, \hat{\theta}_\mu)}{\mathcal{L}(data 0, \hat{\theta}_0)}$	yes	Bayesian-frequentist hybrid
LHC	$\tilde{q}_\mu = -2 \ln \frac{\mathcal{L}(data \mu, \hat{\theta}_\mu)}{\mathcal{L}(data \hat{\mu}, \hat{\theta})}$	yes $(0 \leq \hat{\mu} \leq \mu)$	frequentist

- LEP: nuisance parameters (  $\theta$  ) not profiled.
- Tevatron: Bayesian treatment.
  - ▣ Maximise likelihood against nuisances.
- LHC: profiled likelihood.
  - ▣ Denominator uses **global best-fit likelihood** with **constrained signal strength**.



# Measuring the mass

21



**ATLAS**

**CMS**


$m_X$   $125.5 \pm 0.2 \text{ (stat.) } ^{+0.5}_{-0.6} \text{ (syst.) GeV}$

$125.7 \pm 0.3 \text{ (stat.) } \pm 0.3 \text{ (syst.) GeV}$





**Naïve average:  $125.6 \pm 0.4 \text{ GeV}$**

# Parity: $H \rightarrow ZZ \rightarrow 4\ell$

[CMS-PAS-HIG-13-003] [ATLAS-CONF-2013-013]





- Discriminants built from decay angles and invariant masses.
- Profiled likelihood ratio test statistic.



$CL_s$	0.37%	0.16%
$P(\text{obs.}   0^+)$	0.40	$-0.5 \sigma$
$P(\text{obs.}   0^-)$	0.0022	$3.3 \sigma$

# Other $J^P$ in $H \rightarrow ZZ \rightarrow 4\ell$



		J <sup>P</sup> -MELA analysis			CL <sub>S</sub>
		tested J <sup>P</sup> for an assumed 0 <sup>+</sup>		tested 0 <sup>+</sup> for an assumed J <sup>P</sup>	
		expected	observed	observed*	
0 <sup>-</sup>	p <sub>0</sub>	0.0011	0.0022	0.40	0.004
1 <sup>+</sup>	p <sub>0</sub>	0.0031	0.0028	0.51	0.006
1 <sup>-</sup>	p <sub>0</sub>	0.0010	0.027	0.11	0.031
2 <sup>+</sup> <sub>m</sub>	p <sub>0</sub>	0.064	0.11	0.38	0.182
2 <sup>-</sup>	p <sub>0</sub>	0.0032	0.11	0.08	0.116

J <sup>P</sup>	production	expect (μ=1)	obs. 0 <sup>+</sup>	obs. J <sup>P</sup>	CL <sub>S</sub>
0 <sup>-</sup>	gg → X	2.6σ (2.8σ)	0.5σ	3.3σ	0.16%
0 <sup>+</sup> <sub>h</sub>	gg → X	1.7σ (1.8σ)	0.0σ	1.7σ	8.1%
2 <sup>+</sup> <sub>m(gg)</sub>	gg → X	1.8σ (1.9σ)	0.8σ	2.7σ	1.5%
2 <sup>+</sup> <sub>m(qq)</sub>	qq → X	1.7σ (1.9σ)	1.8σ	4.0σ	<0.1%
1 <sup>-</sup>	qq → X	2.8σ (3.1σ)	1.4σ	>4.0σ	<0.1%
1 <sup>+</sup>	qq → X	2.3σ (2.6σ)	1.7σ	>4.0σ	<0.1%

**ATLAS**

**CMS**

**CL<sub>S</sub> for J≠0**

**< 18.2%**


**< 1.5%**




# ATLAS: focus on $2^+_m$

24

[ATLAS-CONF-2013-040]



- Combined  $H \rightarrow ZZ, WW,$  and  $\gamma\gamma$ .
- Scan for fraction of  $(gg/q\bar{q}) \rightarrow 2^+_m$ :
  - $CL_s < 0.06\% \quad \forall f_{q\bar{q}}$ .







# Oversimplified big picture

★ "seen" ★ "tried" - "impossible"	$H \rightarrow b\bar{b}$			$H \rightarrow \tau\tau$			$H \rightarrow WW$			$H \rightarrow ZZ$			$H \rightarrow \gamma\gamma$			$H \rightarrow Z\gamma$			$H \rightarrow \text{inv.}$			$H \rightarrow \mu\mu$			$H \rightarrow c\bar{c}$ $H \rightarrow HH$					
	T	A	C	T	A	C	T	A	C	T	A	C	T	A	C	T	A	C	T	A	C	T	A	C	T	A	C			
ggH	-	-	-	★	★	★	★	★	★	★	★	★	★	★	★	-	★	★				-	★		-					
VBF			★	★	★	★		★	★		★	★		★	★	-						-			-					
VH	★	★	★	★		★	★		★		★			★	★	-				★		-			-					
ttH		★	★	★			★								★	-						-			-					

T – Tevatron  
A – ATLAS  
C – CMS




□ Still much to measure to the right and to the bottom.



# Relative signal strengths



26

[arXiv:1303.6346] [ATLAS-CONF-2013-034] [CMS-PAS-HIG-13-005]





	Tevatron	ATLAS	CMS
$m_H$	125 GeV	125.5 GeV	125.7 GeV
$\mu = \sigma/\sigma_{\text{SM}}$	$1.44^{+0.59}_{-0.56}$	$1.30 \pm 0.20$	$0.80 \pm 0.14$

**Naïve average:  $0.98 \pm 0.11$**



- Scale fermion-mediated (ggH & ttH) and vector-boson-mediate (VBF & VH) together.

# Production mechanisms



- Ratio of production scaling factors does not depend on decay mode.
- **Combined  $> 3\sigma$  evidence for  $\mu_{VBF,VH} / \mu_{ggH,ttH} > 0$ .**

# Scalar coupling structure


**Yukawa  
sector**

**Gauge sector**

**Mixed  
sector**

Up type

Down type



Quark loop

Loops ( $\gamma$ ,  $g$ ) are sensitive to BSM contributions.

# Interim scalar coupling deviations



## framework

30

[arXiv:1209.0040]

Production modes

$$\frac{\sigma_{ggH}}{\sigma_{ggH}^{SM}} = \begin{cases} \kappa_g^2(\kappa_b, \kappa_t, m_H) \\ \kappa_g^2 \end{cases}$$

$$\frac{\sigma_{VBF}}{\sigma_{VBF}^{SM}} = \kappa_{VBF}^2(\kappa_W, \kappa_Z, m_H)$$

$$\frac{\sigma_{WH}}{\sigma_{WH}^{SM}} = \kappa_W^2$$

$$\frac{\sigma_{ZH}}{\sigma_{ZH}^{SM}} = \kappa_Z^2$$

$$\frac{\sigma_{t\bar{t}H}}{\sigma_{t\bar{t}H}^{SM}} = \kappa_t^2$$

Detectable decay modes

$$\frac{\Gamma_{WW^{(*)}}}{\Gamma_{WW^{(*)}}^{SM}} = \kappa_W^2$$

$$\frac{\Gamma_{ZZ^{(*)}}}{\Gamma_{ZZ^{(*)}}^{SM}} = \kappa_Z^2$$

$$\frac{\Gamma_{b\bar{b}}}{\Gamma_{b\bar{b}}^{SM}} = \kappa_b^2$$

$$\frac{\Gamma_{\tau^-\tau^+}}{\Gamma_{\tau^-\tau^+}^{SM}} = \kappa_\tau^2$$

$$\frac{\Gamma_{\gamma\gamma}}{\Gamma_{\gamma\gamma}^{SM}} = \begin{cases} \kappa_Y^2(\kappa_b, \kappa_t, \kappa_\tau, \kappa_W, m_H) \\ \kappa_Y^2 \end{cases}$$

$$\frac{\Gamma_{Z\gamma}}{\Gamma_{Z\gamma}^{SM}} = \begin{cases} \kappa_{(Z\gamma)}^2(\kappa_b, \kappa_t, \kappa_\tau, \kappa_W, m_H) \\ \kappa_{(Z\gamma)}^2 \end{cases}$$

Currently undetectable decay modes

$$\frac{\Gamma_{t\bar{t}}}{\Gamma_{t\bar{t}}^{SM}} = \kappa_t^2$$

$$\frac{\Gamma_{gg}}{\Gamma_{gg}^{SM}} : \text{ see Section 3.1.2}$$

$$\frac{\Gamma_{c\bar{c}}}{\Gamma_{c\bar{c}}^{SM}} = \kappa_c^2$$

$$\frac{\Gamma_{s\bar{s}}}{\Gamma_{s\bar{s}}^{SM}} = \kappa_s^2$$

$$\frac{\Gamma_{\mu^-\mu^+}}{\Gamma_{\mu^-\mu^+}^{SM}} = \kappa_\mu^2$$

Total width

$$\frac{\Gamma_H}{\Gamma_H^{SM}} = \begin{cases} \kappa_H^2(\kappa_i, m_H) \\ \kappa_H^2 \end{cases}$$

□ Narrow-width approximation,  $(\sigma \times BR) = \sigma \cdot \Gamma / \Gamma_H$

# Interim scalar coupling deviations



## framework

31

[arXiv:1209.0040]

### Production modes

$$\frac{\sigma_{ggH}}{\sigma_{ggH}^{SM}} = \begin{cases} \kappa_b^2(\kappa_b, \kappa_t, m_H) \\ \kappa_g^2 \end{cases}$$

$$\frac{\sigma_{VBF}}{\sigma_{VBF}^{SM}} = \kappa_{VBF}^2(\kappa_W, \kappa_Z, m_H)$$

$$\frac{\sigma_{WH}}{\sigma_{WH}^{SM}} = \kappa_W^2$$

$$\frac{\sigma_{ZH}}{\sigma_{ZH}^{SM}} = \kappa_Z^2$$

$$\frac{\sigma_{t\bar{t}H}}{\sigma_{t\bar{t}H}^{SM}} = \kappa_t^2$$

### Detectable decay modes

$$\frac{\Gamma_{WW^{(*)}}}{\Gamma_{WW^{(*)}}^{SM}} = \kappa_W^2$$

$$\frac{\Gamma_{ZZ^{(*)}}}{\Gamma_{ZZ^{(*)}}^{SM}} = \kappa_Z^2$$

$$\frac{\Gamma_{b\bar{b}}}{\Gamma_{b\bar{b}}^{SM}} = \kappa_b^2$$

$$\frac{\Gamma_{\tau^-\tau^+}}{\Gamma_{\tau^-\tau^+}^{SM}} = \kappa_\tau^2$$

$$\frac{\Gamma_{\gamma\gamma}}{\Gamma_{\gamma\gamma}^{SM}} = \begin{cases} \kappa_\gamma^2(\kappa_b, \kappa_t, \kappa_\tau, \kappa_W, m_H) \\ \kappa_\gamma^2 \end{cases}$$

$$\frac{\Gamma_{Z\gamma}}{\Gamma_{Z\gamma}^{SM}} = \begin{cases} \kappa_{(Z\gamma)}^2(\kappa_b, \kappa_t, \kappa_\tau, \kappa_W, m_H) \\ \kappa_{(Z\gamma)}^2 \end{cases}$$

### Currently undetectable decay modes

$$\frac{\Gamma_{t\bar{t}}}{\Gamma_{t\bar{t}}^{SM}} = \kappa_t^2$$

$$\frac{\Gamma_{gg}}{\Gamma_{gg}^{SM}} : \text{ see Section 3.1.2}$$

$$\frac{\Gamma_{c\bar{c}}}{\Gamma_{c\bar{c}}^{SM}} = \kappa_c^2$$

$$\frac{\Gamma_{s\bar{s}}}{\Gamma_{s\bar{s}}^{SM}} = \kappa_s^2$$

$$\frac{\Gamma_{\mu^-\mu^+}}{\Gamma_{\mu^-\mu^+}^{SM}} = \kappa_\mu^2$$

### Total width

$$\frac{\Gamma_H}{\Gamma_H^{SM}} = \begin{cases} \kappa_H^2(\kappa_i, m_H) \\ \kappa_H^2 \end{cases}$$

- Contributions resolved at NLO QCD and LO EWK.
- Peg things unmeasured to closest of kin.

# Probing custodial symmetry

## Probing custodial symmetry assuming no invisible or undetectable widths

Free parameters:  $\kappa_Z, \lambda_{WZ} (= \kappa_W / \kappa_Z), \kappa_f (= \kappa_t = \kappa_b = \kappa_\tau)$ .

	$H \rightarrow \gamma\gamma$	$H \rightarrow ZZ^{(*)}$	$H \rightarrow WW^{(*)}$	$H \rightarrow b\bar{b}$	$H \rightarrow \tau^-\tau^+$
ggH t $\bar{t}$ H	$\frac{\kappa_f^2 \cdot \kappa_f^2 (\kappa_f, \kappa_f, \kappa_f, \kappa_Z \lambda_{WZ})}{\kappa_H^2 (\kappa_f)}$	$\frac{\kappa_f^2 \cdot \kappa_Z^2}{\kappa_H^2 (\kappa_f)}$	$\frac{\kappa_f^2 \cdot (\kappa_Z \lambda_{WZ})^2}{\kappa_H^2 (\kappa_f)}$	$\frac{\kappa_f^2 \cdot \kappa_f^2}{\kappa_H^2 (\kappa_f)}$	$\frac{\kappa_f^2 \cdot \kappa_f^2}{\kappa_H^2 (\kappa_f)}$
VBF	$\frac{\kappa_{VBF}^2 (\kappa_Z, \kappa_Z \lambda_{WZ}) \cdot \kappa_f^2 (\kappa_f, \kappa_f, \kappa_f, \kappa_Z \lambda_{WZ})}{\kappa_H^2 (\kappa_f)}$	$\frac{\kappa_{VBF}^2 (\kappa_Z, \kappa_Z \lambda_{WZ}) \cdot \kappa_Z^2}{\kappa_H^2 (\kappa_f)}$	$\frac{\kappa_{VBF}^2 (\kappa_Z, \kappa_Z \lambda_{WZ}) \cdot (\kappa_Z \lambda_{WZ})^2}{\kappa_H^2 (\kappa_f)}$	$\frac{\kappa_{VBF}^2 (\kappa_Z, \kappa_Z \lambda_{WZ}) \cdot \kappa_f^2}{\kappa_H^2 (\kappa_f)}$	$\frac{\kappa_{VBF}^2 (\kappa_Z, \kappa_Z \lambda_{WZ}) \cdot \kappa_f^2}{\kappa_H^2 (\kappa_f)}$
WH	$\frac{(\kappa_Z \lambda_{WZ})^2 \cdot \kappa_f^2 (\kappa_f, \kappa_f, \kappa_f, \kappa_Z \lambda_{WZ})}{\kappa_H^2 (\kappa_f)}$	$\frac{(\kappa_Z \lambda_{WZ})^2 \cdot \kappa_Z^2}{\kappa_H^2 (\kappa_f)}$	$\frac{(\kappa_Z \lambda_{WZ})^2 \cdot (\kappa_Z \lambda_{WZ})^2}{\kappa_H^2 (\kappa_f)}$	$\frac{(\kappa_Z \lambda_{WZ})^2 \cdot \kappa_f^2}{\kappa_H^2 (\kappa_f)}$	$\frac{(\kappa_Z \lambda_{WZ})^2 \cdot \kappa_f^2}{\kappa_H^2 (\kappa_f)}$
ZH	$\frac{\kappa_Z^2 \cdot \kappa_f^2 (\kappa_f, \kappa_f, \kappa_f, \kappa_Z \lambda_{WZ})}{\kappa_H^2 (\kappa_f)}$	$\frac{\kappa_Z^2 \cdot \kappa_Z^2}{\kappa_H^2 (\kappa_f)}$	$\frac{\kappa_Z^2 \cdot (\kappa_Z \lambda_{WZ})^2}{\kappa_H^2 (\kappa_f)}$	$\frac{\kappa_Z^2 \cdot \kappa_f^2}{\kappa_H^2 (\kappa_f)}$	$\frac{\kappa_Z^2 \cdot \kappa_f^2}{\kappa_H^2 (\kappa_f)}$

## Probing custodial symmetry without assumptions on the total width

Free parameters:  $\kappa_{ZZ} (= \kappa_Z \cdot \kappa_Z / \kappa_H), \lambda_{WZ} (= \kappa_W / \kappa_Z), \lambda_{FZ} (= \kappa_f / \kappa_Z)$ .

	$H \rightarrow \gamma\gamma$	$H \rightarrow ZZ^{(*)}$	$H \rightarrow WW^{(*)}$	$H \rightarrow b\bar{b}$	$H \rightarrow \tau^-\tau^+$
ggH t $\bar{t}$ H	$\kappa_{ZZ}^2 \lambda_{FZ}^2 \cdot \kappa_f^2 (\lambda_{FZ}, \lambda_{FZ}, \lambda_{FZ}, \lambda_{WZ})$	$\kappa_{ZZ}^2 \lambda_{FZ}^2$	$\kappa_{ZZ}^2 \lambda_{FZ}^2 \cdot \lambda_{WZ}^2$	$\kappa_{ZZ}^2 \lambda_{FZ}^2 \cdot \lambda_{FZ}^2$	$\kappa_{ZZ}^2 \lambda_{FZ}^2 \cdot \lambda_{FZ}^2$
VBF	$\kappa_{ZZ}^2 \kappa_{VBF}^2 (1, \lambda_{WZ}^2) \cdot \kappa_f^2 (\lambda_{FZ}, \lambda_{FZ}, \lambda_{FZ}, \lambda_{WZ})$	$\kappa_{ZZ}^2 \kappa_{VBF}^2 (1, \lambda_{WZ}^2)$	$\kappa_{ZZ}^2 \kappa_{VBF}^2 (1, \lambda_{WZ}^2) \cdot \lambda_{WZ}^2$	$\kappa_{ZZ}^2 \kappa_{VBF}^2 (1, \lambda_{WZ}^2) \cdot \lambda_{FZ}^2$	$\kappa_{ZZ}^2 \kappa_{VBF}^2 (1, \lambda_{WZ}^2) \cdot \lambda_{FZ}^2$
WH	$\kappa_{ZZ}^2 \lambda_{WZ}^2 \cdot \kappa_f^2 (\lambda_{FZ}, \lambda_{FZ}, \lambda_{FZ}, \lambda_{WZ})$	$\kappa_{ZZ}^2 \cdot \lambda_{WZ}^2$	$\kappa_{ZZ}^2 \lambda_{WZ}^2 \cdot \lambda_{WZ}^2$	$\kappa_{ZZ}^2 \lambda_{WZ}^2 \cdot \lambda_{FZ}^2$	$\kappa_{ZZ}^2 \lambda_{WZ}^2 \cdot \lambda_{FZ}^2$
ZH	$\kappa_{ZZ}^2 \cdot \kappa_f^2 (\lambda_{FZ}, \lambda_{FZ}, \lambda_{FZ}, \lambda_{WZ})$	$\kappa_{ZZ}^2$	$\kappa_{ZZ}^2 \cdot \lambda_{WZ}^2$	$\kappa_{ZZ}^2 \cdot \lambda_{FZ}^2$	$\kappa_{ZZ}^2 \cdot \lambda_{FZ}^2$








# Probing custodial symmetry

33

[arXiv:1303.6346] [ATLAS-CONF-2013-034] [CMS-PAS-HIG-13-005]



**Tevatron**

$[\kappa_{W'}, \kappa_{Z'}, \kappa_f]$

**ATLAS**

$[\lambda_{WZ}, \lambda_{FZ}, \kappa_{ZZ}]$

**CMS**

$[\lambda_{WZ}, \kappa_{Z'}, \kappa_f]$

$\lambda_{WZ}$

$1.24^{+2.34}_{-0.42}$

$[0.64, 0.87]$

$0.86 \pm 0.13$

# Weak bosons and fermions






## Boson and fermion scaling assuming no invisible or undetectable widths

Free parameters:  $\kappa_V (= \kappa_W = \kappa_Z)$ ,  $\kappa_f (= \kappa_t = \kappa_b = \kappa_\tau)$ .

	$H \rightarrow \gamma\gamma$	$H \rightarrow ZZ^{(*)}$	$H \rightarrow WW^{(*)}$	$H \rightarrow b\bar{b}$	$H \rightarrow \tau^-\tau^+$
ggH ttH	$\frac{\kappa_f^2 \cdot \kappa_\gamma^2 (\kappa_f, \kappa_f, \kappa_f, \kappa_V)}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_f^2 \cdot \kappa_V^2}{\kappa_H^2 (\kappa_i)}$		$\frac{\kappa_f^2 \cdot \kappa_f^2}{\kappa_H^2 (\kappa_i)}$	
VBF WH ZH	$\frac{\kappa_V^2 \cdot \kappa_\gamma^2 (\kappa_f, \kappa_f, \kappa_f, \kappa_V)}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_V^2 \cdot \kappa_V^2}{\kappa_H^2 (\kappa_i)}$		$\frac{\kappa_V^2 \cdot \kappa_f^2}{\kappa_H^2 (\kappa_i)}$	

# Weak bosons and fermions





	Tevatron	ATLAS	CMS
P(SM)	-	8%	$< 1 \sigma$

# Weak bosons and fermions



36

[ATLAS-CONF-2013-034] [CMS-PAS-HIG-13-005]



	ATLAS	CMS
P(SM)	8%	< 1σ

# Looking for new particles



Probing loop structure <b>assuming no invisible</b> or undetectable widths					
Free parameters: $\kappa_g, \kappa_\gamma$ .					
	$H \rightarrow \gamma\gamma$	$H \rightarrow ZZ^{(*)}$	$H \rightarrow WW^{(*)}$	$H \rightarrow b\bar{b}$	$H \rightarrow \tau^-\tau^+$
ggH	$\frac{\kappa_g^2 \cdot \kappa_\gamma^2}{\kappa_H^2(\kappa_i)}$		$\frac{\kappa_g^2}{\kappa_H^2(\kappa_i)}$		
ttH					
VBF	$\frac{\kappa_\gamma^2}{\kappa_H^2(\kappa_i)}$		$\frac{1}{\kappa_H^2(\kappa_i)}$		
WH					
ZH					

Probing loop structure <b>allowing for invisible</b> or undetectable widths					
Free parameters: $\kappa_g, \kappa_\gamma, BR_{inv.,undet.}$ .					
	$H \rightarrow \gamma\gamma$	$H \rightarrow ZZ^{(*)}$	$H \rightarrow WW^{(*)}$	$H \rightarrow b\bar{b}$	$H \rightarrow \tau^-\tau^+$
ggH	$\frac{\kappa_g^2 \cdot \kappa_\gamma^2}{\kappa_H^2(\kappa_i)/(1-BR_{inv.,undet.})}$		$\frac{\kappa_g^2}{\kappa_H^2(\kappa_i)/(1-BR_{inv.,undet.})}$		
ttH					
VBF	$\frac{\kappa_\gamma^2}{\kappa_H^2(\kappa_i)/(1-BR_{inv.,undet.})}$		$\frac{1}{\kappa_H^2(\kappa_i)/(1-BR_{inv.,undet.})}$		
WH					
ZH					



$$\kappa_i^2 = \Gamma_{ii}/\Gamma_{ii}^{SM}$$



# Looking for new particles in loops



38

[ATLAS-CONF-2013-034] [CMS-PAS-HIG-13-005]




	ATLAS	CMS
$\kappa_\gamma$	$1.23^{+0.16}_{-0.13}$	$0.97 \pm 0.18$
$\kappa_g$	$1.08 \pm 0.14$	$0.83 \pm 0.11$

# Looking for new particles






	ATLAS	CMS
$BR_{BSM}$	<b>&lt; 0.6 (95% CL)</b>	<b>&lt; 0.52 (95% CL)</b>

# ZH → ℓℓ + invisible




- MET > 90 GeV.
- 2D sideband on:
  - $|\text{MET} - p_T^{\ell\ell}| / p_T^{\ell\ell}$
  - $\Delta\phi(\text{MET}, p_T^{\text{miss.}})$
- Not yet sensitive to standard candle:  
ZH → ZZZ → 2ℓ4ν
- **At  $m_H = 125$  GeV,**  
**BR<sub>inv.</sub> < 0.65 (0.84)**  
**(95%CL), obs.(exp.).**









# $H \rightarrow Z \gamma \rightarrow \ell \ell \gamma$



- Loop-mediated decay: sensitive to BSM.
- Both analyses on full 7 and 8 TeV data sets.




Obs. (exp.)

 **$\mu$  at 125 GeV (95% CL)**
**ATLAS**
**< 18.2 (13.5)**
**CMS**
**< 9 (12)**

# A further take on loops

- Resolve the  $H \rightarrow \gamma \gamma$ ,  $H \rightarrow Z \gamma$ , and  $ggH$  loops.



# Probing the fermion sector

	u-type	d-type	lepton		
2HDM	I	$\frac{\cos \alpha}{\sin \beta}$	$\frac{\cos \alpha}{\sin \beta}$	$\frac{\cos \alpha}{\sin \beta}$	SM-like
	I'	$\frac{\cos \alpha}{\sin \beta}$	$\frac{\cos \alpha}{\sin \beta}$	$\frac{-\sin \alpha}{\cos \beta}$	
	II	$\frac{\cos \alpha}{\sin \beta}$	$\frac{-\sin \alpha}{\cos \beta}$	$\frac{-\sin \alpha}{\cos \beta}$	
	II'	$\frac{\cos \alpha}{\sin \beta}$	$\frac{-\sin \alpha}{\cos \beta}$	$\frac{\cos \alpha}{\sin \beta}$	

## Probing up-type and down-type fermion symmetry assuming no invisible or undetectable widths

Free parameters:  $\kappa_V (= \kappa_Z = \kappa_W)$ ,  $\lambda_{du} (= \kappa_d/\kappa_u)$ ,  $\kappa_u (= \kappa_t)$ .



	$H \rightarrow \gamma\gamma$	$H \rightarrow ZZ^{(*)}$	$H \rightarrow WW^{(*)}$	$H \rightarrow b\bar{b}$	$H \rightarrow \tau^-\tau^+$
ggH	$\frac{\kappa_g^2 (\kappa_u \lambda_{du}, \kappa_u) \cdot \kappa_\gamma^2 (\kappa_u \lambda_{du}, \kappa_u, \kappa_u \lambda_{du}, \kappa_V)}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_g^2 (\kappa_u \lambda_{du}, \kappa_u) \cdot \kappa_V^2}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_g^2 (\kappa_u \lambda_{du}, \kappa_u) \cdot (\kappa_u \lambda_{du})^2}{\kappa_H^2 (\kappa_i)}$		
t $\bar{t}$ H	$\frac{\kappa_u^2 \cdot \kappa_\gamma^2 (\kappa_u \lambda_{du}, \kappa_u, \kappa_u \lambda_{du}, \kappa_V)}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_u^2 \cdot \kappa_V^2}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_u^2 \cdot (\kappa_u \lambda_{du})^2}{\kappa_H^2 (\kappa_i)}$		
VBF WH ZH	$\frac{\kappa_V^2 \cdot \kappa_\gamma^2 (\kappa_u \lambda_{du}, \kappa_u, \kappa_u \lambda_{du}, \kappa_V)}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_V^2 \cdot \kappa_V^2}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_V^2 \cdot (\kappa_u \lambda_{du})^2}{\kappa_H^2 (\kappa_i)}$		



## Probing quark and lepton fermion symmetry assuming no invisible or undetectable widths

Free parameters:  $\kappa_V (= \kappa_Z = \kappa_W)$ ,  $\lambda_{lq} (= \kappa_l/\kappa_q)$ ,  $\kappa_q (= \kappa_t = \kappa_b)$ .



	$H \rightarrow \gamma\gamma$	$H \rightarrow ZZ^{(*)}$	$H \rightarrow WW^{(*)}$	$H \rightarrow b\bar{b}$	$H \rightarrow \tau^-\tau^+$
ggH	$\frac{\kappa_q^2 \cdot \kappa_\gamma^2 (\kappa_q, \kappa_q, \kappa_q \lambda_{lq}, \kappa_V)}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_q^2 \cdot \kappa_V^2}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_q^2 \cdot \kappa_q^2}{\kappa_H^2 (\kappa_i)}$		
t $\bar{t}$ H	$\frac{\kappa_q^2 \cdot \kappa_\gamma^2 (\kappa_q, \kappa_q, \kappa_q \lambda_{lq}, \kappa_V)}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_q^2 \cdot \kappa_V^2}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_q^2 \cdot (\kappa_q \lambda_{lq})^2}{\kappa_H^2 (\kappa_i)}$		
VBF WH ZH	$\frac{\kappa_V^2 \cdot \kappa_\gamma^2 (\kappa_q, \kappa_q, \kappa_q \lambda_{lq}, \kappa_V)}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_V^2 \cdot \kappa_V^2}{\kappa_H^2 (\kappa_i)}$	$\frac{\kappa_V^2 \cdot \kappa_q^2}{\kappa_H^2 (\kappa_i)}$		

# Probing the fermion sector



$\lambda_{du}$

$\lambda_{lq}$

**CMS**

**[0.74, 1.95] (95% CL)**


**[0.57, 2.05] (95% CL)**

# Summary of scalar couplings tests

**ATLAS Preliminary**  $\sqrt{s} = 7 \text{ TeV}, \int \text{Ldt} = 4.6\text{-}4.8 \text{ fb}^{-1}$


■  $\pm 1\sigma$    ■  $\pm 2\sigma$

$\sqrt{s} = 8 \text{ TeV}, \int \text{Ldt} = 13\text{-}20.7 \text{ fb}^{-1}$



$m_H = 125.5 \text{ GeV}$

parameter value





# Resolving SM contributions

46

[CMS-PAS-HIG-13-005]

- Individual couplings:

$$\kappa_W, \kappa_Z, \kappa_b, \kappa_t,$$
$$\kappa_\tau.$$

- All loops resolved:


$$\kappa_\gamma(\kappa_W, \kappa_t), \kappa_g(\kappa_t,$$
$$\kappa_b).$$

- SMH width scaled.

- Not explicitly a LHC HXSWG benchmark.

# Resolving SM contributions

- **$P(\text{SM})=0.52$ .**
- “Reduced” couplings as function of “mass”:
  - $\lambda_f = \kappa_f (m_f/\text{vev})$
  - $(g_V/2\text{vev})^{1/2} = \kappa_V^{1/2} (m_V/\text{vev})$



# Birth of a Higgs boson

Results from ATLAS and CMS now provide enough evidence to identify the new particle of 2012 as ‘a Higgs boson’.

In the history of particle physics, July 2012 will feature prominently as the date when the ATLAS and CMS collaborations announced that they had discovered a new particle with a mass near 125 GeV in studies of proton–proton collisions at the LHC. The discovery followed just over a year of dedicated searches for the Higgs boson, the particle linked to the Brout-Englert-Higgs mechanism that endows elementary particles with mass. At this early stage, the phrase “Higgs-like boson” was the recognized shorthand for a boson whose properties were yet to be fully investigated (*CERN Courier* September 2012 p43 and p49). The outstanding performance of the LHC in the second half of 2012 delivered four times as much data at 8 TeV in the centre of mass as were used in the “discovery” analyses. Thus equipped, the experiments were able to present new results at the 2013 Rencontres de Moriond in March, giving the particle-physics community enough evidence to


March, giving the particle-physics community enough evidence to name this new boson “a Higgs boson”.

results that further elucidate the nature of the particle discovered just eight months earlier. The collaborations find that the new particle is looking more and more like a Higgs boson. However, it remains an open question whether this is *the* Higgs boson of the Standard Model of particle physics, or one of several such bosons predicted in theories that go beyond the Standard Model. Finding the answer to this question will require more time and data.

This brief summary provides an update of the measurements





Observed $CL_s$ compared with $J^P=0^+$	$0^-$ (gg) pseudo-scalar	$2_m^+$ (gg) minimal couplings	$2_m^+$ (q $\bar{q}$ ) minimal couplings	$1^-$ (q $\bar{q}$ ) exotic vector	$1^+$ (q $\bar{q}$ ) exotic pseudo-vector	
ZZ <sup>(*)</sup>	ATLAS	2.2%	6.8%	16.8%	6.0%	0.2%
	CMS	0.16%	1.5%	<0.1%	<0.1%	<0.1%
WW <sup>(*)</sup>	ATLAS	–	5.1%	1.1%	–	–
	CMS	–	14%	–	–	–
$\Upsilon\Upsilon$	ATLAS	–	0.7%	12.4%	–	–

Table 1. Summary of preliminary results of the hypothesis tests compared with the Standard Model hypothesis of no spin, positive parity ( $J^P=0^+$ ). All alternatives are disfavoured using the  $CL_s$  ratio of probabilities that takes into account how the observation relates to both the Standard Model and the alternative hypotheses.





# Timeline of the results



# Conclusions



50






- **Went from “a new particle” to “a Higgs Boson”.**
  - $m_\chi \sim 125.6 \pm 0.4$  GeV.
  
- **Big picture unfailingly consistent with SMH:**
  - Per channel, per final state, and per production mode.
  - No significant deviations of scalar couplings.
  - Parity hypothesis tests disfavor  $0^-$ .
  - Other  $J^P$  hypothesis tests disfavor  $J \neq 0$ .
  
- **Working hard to leave no stone unturned.**
  - Look for the Higgs parallel session on Wednesday.
  - **Theoretical progress still needed (ggH).**
  - Many channels in the works:  $t\bar{t}H$ ,  $\mu\mu$ , invisible.
  
- **For when a LHC combination?**

# The beautiful boring 2013 Universe

□ **Up above:** “Simple six-parameter  $\Lambda$ CDM”.

□ **Down below:** (Not-as-simple) Standard Model of Particle Physics.



Looking forward to LHC combination and surprises at higher energy: PeV neutrinos, LHC 13 TeV, ...




# “...and references therein.”

- ATLAS  
<https://twiki.cern.ch/twiki/bin/view/AtlasPublic/HiggsPublicResults>
- CMS  
<https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsHIG>
- Tevatron  
<http://tevnphwg.fnal.gov/>
  - CDF  
<http://www-cdf.fnal.gov/physics/new/hdg/Results.html>
  - D0  
[http://www-d0.fnal.gov/d0\\_publications/d0\\_pubs\\_list\\_bytopic.html#higgs](http://www-d0.fnal.gov/d0_publications/d0_pubs_list_bytopic.html#higgs)




53

# For discussion


# VBF, $H \rightarrow b\bar{b}$






- Neural network event classifier.
- Simultaneous  $m_{b\bar{b}}$  fits to 4 ANN categories.
- At  $m_H = 125$  GeV,  $\mu < 3.6$  (3.0) (95%CL), obs.(exp.) or  $\mu = 0.7 \pm 1.4$ .



# $ttH, H \rightarrow \gamma \gamma$






- Tagging of leptonic and hadronic W decays from top (anti-)quarks.
- At  $m_H = 125$  GeV,  $\mu < 5.4$  (5.3) (95%CL), obs.(exp.).





# $H \rightarrow \mu \mu$

- Clean final state.
- Probe coupling to second-generation fermion.
- $BR < 10^{-4}$  in the search range.
- At  $m_H = 125$  GeV,  $\mu < 9.8$  (8.2) (95%CL), obs.(exp.).





# Measuring the mass



□ Combinations of the high-resolution channels.



# Where we stand today – with cookery

58



Significance	$H \rightarrow ZZ$	$H \rightarrow \gamma \gamma$	$H \rightarrow WW$	$H \rightarrow b\bar{b}$	$H \rightarrow \tau \tau$
Obs. (pre-fit exp.)					
<b>ATLAS</b>	$\mu = 1.5 \pm 0.4$ <b><math>\sim 3.8 \sigma</math></b> ( $\sim 2.5 \sigma$ )	$\mu = 1.6 \pm 0.3$ <b><math>\sim 5.3 \sigma</math></b> ( $\sim 3.3 \sigma$ )	<b><math>2.5 \sigma</math></b> ( $1.6 \sigma$ )	$-0.4 \sigma$ ( $1.0 \sigma$ )	$1.1 \sigma$ ( $1.7 \sigma$ )
	125.5 GeV			125 GeV	
<b>CMS</b>	$6.7 \sigma$ ( $7.1 \sigma$ )	$3.9 \sigma$ ( $4.2 \sigma$ )	$3.9 \sigma$ ( $5.6 \sigma$ )	$2.0 \sigma$ ( $2.1 \sigma$ )	<b><math>2.8 \sigma</math></b> ( $2.7 \sigma$ )
	125.7 GeV				

- Combined p-values  $< 10^{-20}$  are telling us to make measurements...


# Looking well ahead



- 300/fb at 14 TeV:
  - ▣ Vast improvement over present datasets.
  - ▣ Room for theory improvements.
- For (HL-LHC) 3 ab<sup>-1</sup>:
  - ▣ self-coupling seems feasible with  $\lambda_{HH} \sim 3\sigma$  /expt.

CMS Projection



- Slight difference in ATLAS results:
  - $\Delta m = 2.3^{+0.6}_{-0.7}(\text{stat.}) \pm 0.6(\text{syst.}) \text{ GeV}$
  - $2.4\sigma$  ( $p=1.5\%$ )
- Using more conservative energy scale uncertainties:  **$1.8\sigma$  ( $p=8\%$ )**.










# CMS: channel compatibility

62







# Probing possible 2HDM

63

[CMS-PAS-HIG-13-005]



$\lambda_{du}$

$\lambda_{lq}$

CMS

[0.74, 1.95] (95% CL)

[0.57, 2.05] (95% CL)





- The spin-2 amplitude has many (higher-order) terms:

$$\begin{aligned}
 A(X \rightarrow V_1 V_2) = \Lambda^{-1} & \left[ 2g_1^{(2)} t_{\mu\nu} f^{*(1)\mu\alpha} f^{*(2)\nu\alpha} + 2g_2^{(2)} t_{\mu\nu} \frac{q_\alpha q_\beta}{\Lambda^2} f^{*(1)\mu\alpha} f^{*(2)\nu\beta} + g_3^{(2)} \frac{\tilde{q}^\beta \tilde{q}^\alpha}{\Lambda^2} t_{\beta\nu} \left( f^{*(1)\mu\nu} f_{\mu\alpha}^{*(2)} + f^{*(2)\mu\nu} f_{\mu\alpha}^{*(1)} \right) \right. \\
 & + g_4^{(2)} \frac{\tilde{q}^\nu \tilde{q}^\mu}{\Lambda^2} t_{\mu\nu} f^{*(1)\alpha\beta} f_{\alpha\beta}^{*(2)} + m_V^2 \left( 2g_5^{(2)} t_{\mu\nu} \epsilon_1^{*\mu} \epsilon_2^{*\nu} + 2g_6^{(2)} \frac{\tilde{q}^\mu q_\alpha}{\Lambda^2} t_{\mu\nu} (\epsilon_1^{*\nu} \epsilon_2^{*\alpha} - \epsilon_1^{*\alpha} \epsilon_2^{*\nu}) + g_7^{(2)} \frac{\tilde{q}^\mu \tilde{q}^\nu}{\Lambda^2} t_{\mu\nu} \epsilon_1^* \epsilon_2^* \right) \\
 & \left. + g_8^{(2)} \frac{\tilde{q}_\mu \tilde{q}_\nu}{\Lambda^2} t_{\mu\nu} f^{*(1)\alpha\beta} \tilde{f}_{\alpha\beta}^{*(2)} + m_V^2 \left( g_9^{(2)} \frac{t_{\mu\alpha} \tilde{q}^\alpha}{\Lambda^2} \epsilon_{\mu\nu\rho\sigma} \epsilon_1^{*\nu} \epsilon_2^{*\rho} q^\sigma + \frac{g_{10}^{(2)} t_{\mu\alpha} \tilde{q}^\alpha}{\Lambda^4} \epsilon_{\mu\nu\rho\sigma} q^\rho \tilde{q}^\sigma (\epsilon_1^{*\nu} (q\epsilon_2^*) + \epsilon_2^{*\nu} (q\epsilon_1^*)) \right) \right], \quad (18)
 \end{aligned}$$

# Spin is so much more than a number

- The spin-2 amplitude has many (higher-order) terms:

$$\begin{aligned}
 A(X \rightarrow V_1 V_2) = \Lambda^{-1} & \left[ 2g_1^{(2)} t_{\mu\nu} f^{*(1)\mu\alpha} f^{*(2)\nu\alpha} + 2g_2^{(2)} \frac{q_\alpha q_\beta}{\Lambda^2} f^{*(1)\mu\alpha} f^{*(2)\nu\beta} + g_3^{(2)} \frac{\tilde{a}^\beta \tilde{a}^\alpha}{\Lambda^2} (f^{*(1)\mu\nu} f^{*(2)\mu\alpha} - f^{*(1)\mu\alpha} f^{*(2)\mu\nu}) \right. \\
 & + g_4^{(2)} \frac{\tilde{a}^\nu \tilde{a}^\mu}{\Lambda^2} f^{*(1)\mu\alpha} f^{*(2)\nu\alpha} + m_V^2 \left( 2g_5^{(2)} t_{\mu\nu} \epsilon_1^{*\mu} \epsilon_2^{*\nu} + 2g_6^{(2)} \frac{\tilde{a}^\mu q_\alpha}{\Lambda^2} (f^{*(1)\mu\alpha} \epsilon_2^{*\nu} - f^{*(2)\mu\alpha} \epsilon_1^{*\nu}) + g_7^{(2)} \frac{\tilde{a}^\mu \tilde{a}^\nu}{\Lambda^2} \epsilon_1^{*\mu} \epsilon_2^{*\nu} \right) \\
 & \left. + g_8^{(2)} \frac{\tilde{a}_\nu \tilde{a}_\nu}{\Lambda^2} f^{*(1)\mu\alpha} f^{*(2)\mu\alpha} + m_V^2 \left( g_9^{(2)} \frac{t_{\mu\alpha} \tilde{a}^\alpha}{\Lambda^2} \epsilon_1^{*\mu} \tilde{a}^\nu + g_{10}^{(2)} \frac{g^{\mu\nu} t_{\mu\alpha} \tilde{a}^\alpha}{\Lambda^2} \epsilon_1^{*\nu} \epsilon_2^{*\mu} \right) \right], \quad (18)
 \end{aligned}$$


- Keep only dim-4 terms ( $g_1 = g_5 \neq 0$ ):
  - ▣ Graviton-like “couplings” ( $2^+_m$ ).

- **Until there is enough data, perform hypothesis tests against SMH ( $0^+$ ).**
- Select models using simplifying assumptions on amplitudes:
  - ▣  $0^-$  (parity) “from” ZZ.
  - ▣  $2_m^+$  (graviton-like minimal couplings) also “from” WW and  $\gamma\gamma$ .

scenario	$X \rightarrow ZZ$	$X \rightarrow WW$	$X \rightarrow \gamma\gamma$
$0_m^+$ vs background	5.0	5.0	5.0
$0_m^+$ vs $0_h^+$	1.7	1.1	0.0
$0_m^+$ vs $0^-$	2.9	1.2	0.0
$0_m^+$ vs $1^+$	1.9	2.0	–
$0_m^+$ vs $1^-$	2.6	3.2	–
$0_m^+$ vs $2_m^+$	1.5	2.8	2.4
$0_m^+$ vs $2_h^+$	$\sim 5$	1.1	3.1
$0_m^+$ vs $2_h^-$	$\sim 5$	2.5	3.1

# CMS: $2_m^+$ combination

- Combination of  $H \rightarrow ZZ, WW$ :
  - ▣  $p(\text{obs.} | 0^+) = -0.34 \sigma$
  - ▣  $p(\text{obs.} | 2_m^+(gg)) = 2.84 \sigma$
  - ▣  $CL_s = 0.6\%$







# “C6” vs “resolved C6”

## Generic coupling fit

- Assume custodial symmetry ( $\kappa_V = \kappa_W = \kappa_Z$ ).
- Loops treated effectively ( $\kappa_\gamma, \kappa_g$ ).
- Option to allow BSM decays, forcing  $\kappa_V \leq 1$ .

## Resolved coupling fit

- Keep W and Z separate.
- Loops assuming SM structure:
  - $\kappa_g (\kappa_b, \kappa_t)$ .
  - $\kappa_\gamma (\kappa_W, \kappa_b, \kappa_t, \kappa_\tau)$ .
- Only SM-like decays.

# “C6” vs “resolved C6”

